WHOI-78-12

WIND STRESS ON THE OCEAN OVER THE
EASTERN CONTINENTAL SHELF OF NORTH AMERICA

by

Peter M. Saunders

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Woods Hole, Massachusetts 02543

July 1978

TECHNICAL REPORT

*Prepared for the Office of Naval Research
under Contract N00014-74-C-0262; NR 083-004.*

*Reproduction in whole or in part is permitted
for any purpose of the United States Government.
In citing this report in a bibliography, the
reference should be given to the "Journal of
Physical Oceanography, Vol. 7, No. 4, July 1977,
pp. 555-566".*

*Approved for public release; distribution un-
limited.*

Approved for Distribution: Valentine Worthington, Chairman
Department of Physical Oceanography

Wind Stress on the Ocean over the Eastern Continental Shelf of North America¹

PETER M. SAUNDERS²

Woods Hole Oceanographic Institution, Woods Hole, Mass. 02543

(Manuscript received 8 December 1976, in revised form 2 March 1977)

ABSTRACT

Employing one million ship reports gathered in the years 1941–72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1} . Atmospheric stratification is taken into account but its effect is shown to be small.

In the summer season the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons the stress is directed toward the south and east being strongest in winter ($1\text{--}1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25\text{--}0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.

1. Introduction

Dynamical studies of the circulation in shallow coastal regions require knowledge of atmospheric forcing, including pressure, wind stress and heat fluxes. In this paper I have made use of the archived data incorporated in the voluntary surface meteorological messages reported at standard times (0000 and 1200 GMT) by commercial shipping. Such data, made at random positions, seemed most suited for calculations of the climatology of atmospheric forcing and the wind stress was chosen because it is a significant term. It is not clear to the author what properties of the wind stress are most important so that the simplest statistics, namely, mean and standard deviation, have been chosen. However, to provide further data polar histograms of stress frequency have also been computed. It was my aim to condense the million ship data to a somewhat more manageable size: thus this report should be regarded as the description of a data source or a data summary rather than the results of a scientific investigation.

FIG. 1. A comparison of three-month stress averages (both east and north) computed employing two drag coefficients (see text) for 1° squares in the area $70\text{--}80^\circ\text{W}$, $40\text{--}50^\circ\text{N}$. Solid line has a slope of unity.

¹ Contribution No. 3880 from the Woods Hole Oceanographic Institution.

² Present affiliation: Institute of Oceanographic Sciences, Wormley, Godalming, Surrey, GU8 5UB, England.

2. Choice of drag coefficient

The National Climatic Center at Asheville, N.C., furnished magnetic tapes of surface marine messages

FIG. 2. Number of observations in each 1° square for all months of the period 1941-72 for the region studied in this paper. The 200 m isobath is shown as a dotted line.

to Mr. A. F. Bunker of the Woods Hole Oceanographic Institution, who in turn kindly made them available to the author. Each message contains a report of ships position and of wind speed and direction (along with numerous other meteorological variables), the wind U measured at a nominal height of 10 m above the sea surface. For each individual observation the stress τ is estimated from

$$\tau = \rho C_D U |U|, \quad (1)$$

where ρ is the air density (here given a standard value of $1.2 \times 10^{-3} \text{ g cm}^{-3}$) and C_D is the drag coefficient. The drag coefficient is itself a function of wind speed, of air-sea temperature difference and possibly of sea state, and is the subject of considerable controversy. The recent findings of Smith and Banke (1975) have been adopted in this paper (see also Saunders, 1976), which under neutral conditions give a value

for C_D of

$$C_D = 0.63 \times 10^{-3} + 0.66 \times 10^{-6} |U|, \quad (2)$$

where $|U|$ is measured in centimeters per second. This equation yields a value of $10^3 C_D = 1.0$ at 5 m s^{-1} , 1.3 at 10 m s^{-1} , 1.6 at 15 m s^{-1} and 2.3 at 25 m s^{-1} .

It is, of course, unsatisfactory to relate a non-dimensional parameter C_D to the magnitude of a physical variable, viz., $|U|$, and Eq. (2) is employed here merely for computational convenience. Charnock (1955) has suggested that the roughness length z_0 is given by

$$z_0 = U_*^2 / (ag),$$

where U_* is the friction velocity $[=(\tau/\rho)^{1/2}]$ and g the acceleration of gravity. If the constant a is assigned a value near 80 and the logarithmic profile equation is employed, Smith and Banke (1975) show

FIG. 3. Monthly averaged east and north stress components in areas of approximately $3^{\circ} \times 1^{\circ}$ for the period 1941-72. Horizontal bars indicate three-month averages. (a) mid-Atlantic bight (40°N , 70°W ; total number of observations 42 681); (b) tail of Grand Banks (43°N , 50°W ; total number of observations 19 402); (c) northern Florida (30°N , 80°W ; total number of observations 17 008).

FIG. 4. Magnitude and direction of the climatological winter wind stress. The standard error of the magnitude of the stress is also shown (e.g., ± 13).

FIG. 5. Contours of magnitude of climatological winter wind stress; arrows indicate direction in which stress acts.

that the drag coefficients deduced therefrom fit their measurements as well as Eq. (2). To take into account the effect of atmospheric stratification (both stable and unstable conditions) the relations of Deardorff (1968) derived from overland studies were employed. Later it will be shown that the stratification correction is quite small and therefore the exact formulation unimportant compared with the choice of neutral drag coefficient.

In a recent review Pond (1975) has recommended that a value of $C_D = 1.5 \times 10^{-3}$ independent of both stratification and wind speed be employed. In a trial area, Marsden square 152 which is bounded by longitudes 70° and 80° W and latitudes 40° and 50° N (and contains Cape Cod), the climatological east and north stress components were computed for 1° squares and by three-month seasons employing the drag coefficient of Smith and Banke

FIG. 6. Polar histograms of stress frequency (parts per thousand) for the winter months showing the offshore increase. Left for 1° square 37° N, 76° W; right for 37° N, 73° W. The stress boundaries are shown in Table 2. Climatological means are indicated by arrows.

TABLE 1. Summary of the average standard deviation of a component of the wind stress $\sigma = \frac{1}{2}(\sigma_E + \sigma_N)$ [dyn cm⁻²].

Location	Winter		Spring		Summer		Fall		Annual	
	Inshore	Offshore	Inshore	Offshore	Inshore	Offshore	Inshore	Offshore	Inshore	Offshore
South of 42°N	1.5	2.5	1.0	2.0	0.5	1.0	1.0	2.0	1.0	2.0
Gulf of Maine/Nova Scotia	2.5	3.0	1.5	2.5	1.0	1.5	1.5	2.0	1.5	2.0
Gulf of St. Lawrence	2.5	4.5	2.0	3.0	1.0	1.0	2.0	3.0	2.0	3.0
Newfoundland	3.0	4.0	2.0	2.5	1.0	1.5	2.0	2.5	2.0	2.5

and that suggested by Pond. As seen in Fig. 1 the results are indistinguishable which can be interpreted to mean that the rms scalar wind speed is close to 13 m s⁻¹, the value at which the two drag coefficients are identical. This result covers a range of values of average stress from 0 to 1.0 dyn cm⁻²; larger average values, which involve more frequent strong winds (>20 m s⁻¹) may be expected to yield lower values of average stress for $C_D = 1.5 \times 10^{-3}$ than those obtained with Eq. (2)—perhaps 30% less with climatological values from Eq. (2) of 2.0 dyn cm⁻².

Although all the results reported in this paper *have* included the effect of atmospheric stratification, trial computations have been made assuming neutral conditions; the principal difference is a reduction in computer running time by a factor of nearly one-half! The differences in average stress are characteristically 0.02–0.05 dyn cm⁻² in winter (larger with stratification taken into account) and 0.01–0.03 dyn cm⁻² in summer (smaller with stratification taken into account). These comparisons were made for Marsden square 150 which is bounded by longitudes 50° and 60°W and latitudes 40° and 50°N (and contains Newfoundland).

3. Spatial or temporal resolution?

The total number of observations for the period under study (1941–72) in the area between the coast

and the 200 m isobath, which was taken as the limit of the continental shelf, is approximately one million. The number in each of the approximately 200 1° squares shown in Fig. 2 is thus characteristically a few thousand, ranging from approximately 500–30 000. The question to be answered is to what extent can the annual variation in the climatological stress be represented on a small scale: are the data dense enough to permit a description of both? Because of the magnitude of the stress associated with the passage of weather systems the standard deviation σ of the stress in a period of the order of a month or longer is 1–5 dyn cm⁻². If there are N observations in this period, the standard error ϵ in the estimate of the mean is $\sigma/N^{1/2}$ (assuming a Gaussian process). If $\sigma = 3$ dyn cm⁻² and $N = 100$, $\epsilon = 0.3$ dyn cm⁻², whereas if $N = 1000$, $\epsilon = 0.1$ dyn cm⁻². The latter clearly is a better target for either a seasonal description or a discussion of the spatial variation. It will be seen by examining Fig. 2 that almost all 1° squares contain less than 12 000 data values which would then be the minimum required for a *monthly* description of the average stress. Hence we must either group the squares (approximately by 3) or group the months (also by 3). The latter choice has been made here, computing the seasonal climatology by 1° squares so as to maximize spatial resolution.

FIG. 7. As in Fig. 6 but for 44°N, 63°W (left) and 44°N, 55°W (right).

FIG. 8. Magnitude (and standard error) and direction of the climatological spring wind stress.

To illustrate the seasonal variation by months the average stress components have been computed for three small regions ($1^\circ \times 3^\circ$) in the center, at the northeastern margin, and at the southern margin of the area under study; these are shown in Figs. 3a-3c, respectively. The 3 month seasonal averages are shown as horizontal bars. In Fig. 3a it is observed that little information is lost by the seasonal description, and it is believed that this result is representative of most of the area studied. In Figs. 3b and 3c significant peaks are diminished by seasonal averaging but as will be seen later both of these areas exhibit regional peculiarities.

4. Wind stress in the winter months—December, January, February (1941-1972)

Given a wind observation in one of the above months of the 32-year period an instantaneous stress τ is computed from Eqs. (1) and (2) and those formulated by Deardoff (1968) to take account of atmo-

spheric stratification, mainly air-sea temperature difference. The stress is resolved into east τ_E and north τ_N components and accumulated by 1° squares. The number of observations N and squares of stress components are also accumulated so that mean stress components and their variances σ_E^2 and σ_N^2 can be calculated.

In Fig. 4 the resulting average east and north components are combined into magnitude and direction and tabulated. The direction indicated is that in which the average stress acts, a convention appropriate for oceanographic studies. Values of the standard error of the magnitude of the mean ϵ_τ have also been tabulated and the standard error of the direction ϵ_θ can readily be calculated from the approximate expression $\epsilon_\tau/|\bar{\tau}|$. The above quantities are assumed related to the standard error of the east ϵ_E and north ϵ_N component averages by the relations

$$\epsilon_\tau = (\epsilon_E^2 \bar{\tau}_E^2 + \epsilon_N^2 \bar{\tau}_N^2)^{1/2} / |\bar{\tau}|,$$

$$\epsilon_\theta = (\epsilon_E^2 \bar{\tau}_N^2 + \epsilon_N^2 \bar{\tau}_E^2)^{1/2} / |\bar{\tau}|^2,$$

FIG. 9. Contours of magnitude from Fig. 8.

FIG. 10. As in Fig. 8 except for summer.

FIG. 11. Contours of magnitude from Fig. 10.

with $\epsilon_E = \sigma_E / N^{\frac{1}{2}}$ and $\epsilon_N = \sigma_N / N^{\frac{1}{2}}$. When $\sigma_E = \sigma_N = \sigma_{av}$ (their average) the above expressions simplify to $\epsilon_r = \sigma_{av} / N^{\frac{1}{2}}$ and $\epsilon_\theta = \epsilon_r / |\bar{\tau}|$. The latter are the expressions employed here in calculating ϵ_r and suggested for ϵ_θ . (Test calculations show that σ_E and σ_N differ by less than 25% from σ , their average, so that the above expressions are sufficiently close.)

To illustrate the major features of the distribution the magnitude of the stress has been contoured (see Fig. 5) and schematic indication of the direction made in offshore regions. It will be noted that average climatological winter stresses are everywhere to the east and southeast and show the expected increase from low to high latitude. Considerable small-scale structure also exists and there is generally an increase in stress with distance from the coast. This is particularly marked between Long Island and Cape Hatteras (the mid-Atlantic bight). In the Gulf of Maine there is an exception to this with a weak coastal maximum.

North of 42°N regional structures exist which were quite unexpected. Note the maximum in the Gulf of St. Lawrence, the magnitude decreasing downstream as the coast is approached. Probably the most surprising feature of all is the stress minimum appearing to the south and east of the Grand Banks. Charts from other researchers have been consulted, notably those of Bunker and Worthington (1976) who show a stress maxima near 55–60°W, 35–40°N (south of Newfoundland) and another near 40°W, 45–50°N (east of Newfoundland). A trough of lower stress values protrudes southeast of Newfoundland whose western

edge is seen in this analysis. Land-ocean friction differences probably cannot explain such a large-scale feature and it is merely noted here that centers of cyclogenesis (see Meserve, 1974) correspond approximately to these two regions of stress maximum.³

Table 1 summarizes the average standard deviation of a component by region for the four seasons and the annual average: there is a striking similarity in the distribution of σ and the magnitude of the average stress. Thus large stress values arise principally because of frequent strong winds rather than because of the tendency of the winds to blow more frequently in a particular direction. This is demonstrated by the pairs of histograms illustrating the general offshore increase of stress (Figs. 6, 7) which are examples of further calculations that have been made.

It is evident that the climatological mean and standard deviation of the stress do not furnish all of the required information to compute oceanic response. For example, Beardsley and Butman (1974) have found on the shelf south of New England that a westward stress produces larger wind-driven currents than an eastward stress of the same magnitude (the shoreline is oriented east-west). This directional response, which will be different according to the direction of the shoreline, suggests a need for directional

³ These two centers of high stress separated by a trough of lower stress are probably associated with the "two-gyre" nature of the North Atlantic circulation inferred by Worthington (1962).

stress data. Accordingly, polar histograms have been constructed for each 1° square and season for eight principal directions. The stress boundaries chosen, which are described in Table 2, correspond in the case of neutral stratification to 10 kt wind intervals and have average values which are also tabulated. From the frequency distributions the average stress may be deduced by octants, quadrants, etc. These polar histograms (4 seasons $\times$ 8 directions $\times$ 5 magnitudes $\times$ 200 1° squares = 32 000) represent a reduction by a factor of only 30 on the original data set and cannot be reproduced here.

5. Wind stress in the spring months—March, April, May (1941–72)

The results derived for the spring months (Figs. 8, 9) are generally very similar to those obtained in the winter months. Directions of the climatological stress

TABLE 2. Stress boundaries for polar histograms.

Identification	1	2	3	4	5
Stress range (dyn cm ⁻²)	< 1.5	1.5–4.4	4.4–9.4	9.4–17.0	> 17.0
Mean stress (dyn cm ⁻²)	1	3	7	13	25–30
Wind range* (kt)	< 20	20–30	30–40	40–50	> 50

* For neutral stratification

are similar to those of Fig. 4 but magnitudes are between one-fourth and one-half of the winter months and correspondingly gradients are weaker. In the region of simple offshore increase, south of 42°N , the inshore stress, besides being smaller, is more nearly easterly, i.e., it is $20\text{--}30^\circ$ to the left of the offshore stress. Such an effect might be ascribed to friction, but since it is absent in winter (Figs. 4, 5) it is more likely to result from the temperature contrast between land being warmer than the sea in the spring and colder in the winter. It will be noted that offshore

FIG. 12. As in Fig. 8 except for fall.

FIG. 13. Contours of magnitude from Fig. 12.

FIG. 14. As in Fig. 8 except annual climatological wind stress.

FIG. 15. Contours of magnitude from Fig. 14.

directions are very similar for these two seasons, as are the total number of observations.

6. Wind stress in the summer months—June, July, August (1941-72)

The results derived for the summer month (Figs. 10, 11) are quite different from those of the three other seasons of the year. The direction of the climatological stress is everywhere north and east rather than south and east. Magnitudes are lower, reaching values of 0.3–0.4 dyn cm^{-2} in the region south of Newfoundland and quite generally being only 0.1–0.2 dyn cm^{-2} . Offshore gradients are very weak. It was anticipated that a nearshore intensification, resulting from sea breezes, might be found. Since the times of reporting are 0700 and 1900 locally its absence must rather be expected. This is believed to be the only evidence for bias in the data, a topic returned to in Section 8.

7. Wind stress in the fall months—September, October, November (1941-72)

In the fall months while the climatological stress remains small almost everywhere its direction reverts to that of the winter and spring (Figs. 12, 13). A remarkable exception is the region from northern Florida to Cape Hatteras where the stress is toward the south-southwest and shows a narrow stress maximum

100 km off the coast. We know of no explanation of this regional peculiarity which is seen in monthly detail in Fig. 3c. Again a minimum in the stress magnitude occurs southeast of the Grand Banks and a weak maximum in the northern Gulf of Maine (as is also found in winter and spring).

8. Annual average of wind stress (1941-72)

Figs. 14 and 15 show the annual average data and a contoured magnitude map, respectively. Because of the strength of the winter pattern and the similarity of the fall and spring maps it closely resembles all three seasons, showing little influence of the summer pattern. The results are quite similar to those of Bunker and Worthington (1976). The total number of observations (just over one million) is equally distributed among the four seasons so that no bias should be apparent. When this investigation was begun concern was expressed that bias might appear because a storm avoidance procedures. Ships returning to port have no option but to encounter and report storms. Ships leaving port might delay until the worst of a storm was passed before venturing out, but this would artificially lower both the nearshore and offshore frequency of strong winds (since the storm would generally outrun ships) and would also lower the mean. So far no practicable way has been found of evaluating this suggestion, but the nearshore maxi-

imum in the Gulf of Maine lends support to the conjecture that the bias is not important.

It is not our intention to explore the oceanographic significance of this data summary but the following comments are offered. From Nova Scotia south the climatological stress is predominantly a vector directed offshore. In shallow water such a stress might be expected to drive the water away from the beach lowering mean sea level: in deeper water where bottom friction no longer supports the stress an Ekman-like flow might result with vertically averaged transport to the south and southwest. If a depth of 30 m is assumed with a stress of 0.5 dyn cm^{-2} , the (vertically averaged) wind induced drift is 2 cm s^{-1} . Such a current is in the direction of the long-term drift along the shelf but of somewhat smaller magnitude (Bumpus, 1973) although this generalization does less than justice to the complexity of the observed flow.

Acknowledgments. I wish to acknowledge the programming assistance of Roger Goldsmith and the support of the U.S. Office of Naval Research under Contract N00014-74-C0262.12 NR 083-004.

REFERENCES

- Beardsley, R. C., and B. Butman, 1974: Circulation on the New England continental shelf: Response to strong winter storms. *Geophys. Res. Lett.*, **1**, 181-184.
- Bumpus, D. F., 1973: A description of the circulation on the continental shelf of the East Coast of the United States. *Progress in Oceanography*, Vol. 6, Pergamon Press, 111-157.
- Bunker, A. F., and L. V. Worthington, 1976: Energy exchange charts of the North Atlantic Ocean. *Bull. Amer. Meteor. Soc.*, **57**, 670-678.
- Charnock, H., 1955: Wind stress on a water surface. *Quart. J. Roy. Meteor. Soc.*, **81**, 639-640.
- Deardorff, J. W., 1968: Dependence of air-sea transfer coefficients on bulk stability. *J. Geophys. Res.*, **73**, 2549-2557.
- Meserve, J. M., 1974: *U.S. Navy Marine Climatic Atlas of the World. North Atlantic Ocean*. Vol. 1, Naval Weather Service Command, 369 pp.
- Pond, S., 1975: The exchange of momentum, heat and moisture at ocean-atmosphere interface. *Numerical Models of Ocean Circulation*, Nat. Acad. Sci., 26-38.
- Saunders, P. M., 1976: On the uncertainty of wind stress curl calculations. *J. Mar. Res.*, **34**, 155-160.
- Smith, S. D., and E. G. Banke, 1975: Variation of sea surface drag coefficient with wind speed. *Quart. J. Roy. Meteor. Soc.*, **101**, 665-673.
- Worthington, L. V., 1962: Evidence for a two-gyre circulation system in the North Atlantic. *Deep-Sea Res.*, **9**, 51-67.

MANDATORY DISTRIBUTION LIST

FOR UNCLASSIFIED TECHNICAL REPORTS, REPRINTS, & FINAL REPORTS
PUBLISHED BY OCEANOGRAPHIC CONTRACTORS
OF THE OCEAN SCIENCE AND TECHNOLOGY DIVISION
OF THE OFFICE OF NAVAL RESEARCH

(REVISED FEBRUARY 1978)

1	Director of Defense Research
and Engineering
Office of the Secretary of Defense
Washington, D.C. 20301
ATTN: Office Assistant Director
(Research)	1	National Oceanographic Data
Center
National Oceanic & Atmospheric
Administration
3300 Whitehaven St., N.W.
Washington, D.C. 20235
	Office of Naval Research
Arlington, VA 22217	12	Defense Documentation
Center
Cameron Station
Alexandria, VA 22314
1	ATTN: (Code 460)		Commander
Naval Oceanographic
Office
Washington, D.C. 20373
1	ATTN: (Code 102-OS)	1	ATTN: Code 1640
1	ATTN: (Code 200)	1	ATTN: Code 70
1	CDR J. C. Harlett, (USN)
ONR Representative
Woods Hole Oceanographic Inst.
Woods Hole, MA 02543	3	NORDA 430
NSTL Station, MS 39529
1	Office of Naval Research
Branch Office
495 Summer Street
Boston, MA 02210	2	CO NORDA
NSTL Station, MS 39529
	Director
Naval Research Laboratory
Washington, D.C. 20375		
6	ATTN: Library, Code 2620		

UNCLASSIFIED

7/78

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER WH01-78-12	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) WIND STRESS ON THE OCEAN OVER THE EASTERN CONTINENTAL SHELF OF NORTH AMERICA		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER WH01 CON. NO. 3880
7. AUTHOR(s) Peter M. Saunders		8. CONTRACT OR GRANT NUMBER(s) N00014-74-C-0262;
9. PERFORMING ORGANIZATION NAME AND ADDRESS Woods Hole Oceanographic Institution Woods Hole, MA 02543		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 083-004
11. CONTROLLING OFFICE NAME AND ADDRESS NORDA National Space Technology Laboratory Bay St. Louis, MS 39529		12. REPORT DATE July 1978
		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Reprinted from the "Journal of Physical Oceanography, Vol. 7, No. 4, July 1977, pp. 555-566".		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) 1. Wind Stress 2. American Continental Shelf Waters (Atlantic) 3. Drag Coefficient		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Employing one million ship reports gathered in the years 1941-72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1} . Atmospheric stratification is taken into account but its effect is shown to be small. (Cont. on back)		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

In the summer season the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons the stress is directed toward the south and east being strongest in winter ($1-1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25-0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.

Woods Hole Oceanographic Institution WHOI-78-12	<p>1. Wind Stress</p> <p>2. American Continental Shelf Waters (Atlantic)</p> <p>3. Drag Coefficient</p> <p>I. Saunders, Peter M.</p> <p>II. N00014-74-C-0262; NR 083-004</p>	Woods Hole Oceanographic Institution WHOI-78-12	<p>1. Wind Stress</p> <p>2. American Continental Shelf Waters (Atlantic)</p> <p>3. Drag Coefficient</p> <p>I. Saunders, Peter M.</p> <p>II. N00014-74-C-0262; NR 083-004</p>
<p>WIND STRESS ON THE OCEAN OVER THE EASTERN CONTINENTAL SHELF OF NORTH AMERICA by Peter M. Saunders. Reprinted from the "Journal of Physical Oceanography, Vol. 7, No. 4, July 1977, pp. 555-566", July 1978. Prepared for the Office of Naval Research under Contract N00014-74-C-0262; NR 083-004.</p> <p>Employing one million ship reports gathered in the years 1941-72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1}. Atmospheric stratification is taken into account but its effect is shown to be small.</p> <p>In the summer season, the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons, the stress is directed toward the south and east being strongest in winter ($1-1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25-0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter, the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in</p> <p>(Cont. on back)</p>	<p>WIND STRESS ON THE OCEAN OVER THE EASTERN CONTINENTAL SHELF OF NORTH AMERICA by Peter M. Saunders. Reprinted from the "Journal of Physical Oceanography, Vol. 7, No. 4, July 1977, pp. 555-566", July 1978. Prepared for the Office of Naval Research under Contract N00014-74-C-0262; NR 083-004.</p> <p>Employing one million ship reports gathered in the years 1941-72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1}. Atmospheric stratification is taken into account but its effect is shown to be small.</p> <p>In the summer season, the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons, the stress is directed toward the south and east being strongest in winter ($1-1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25-0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter, the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in</p> <p>(Cont. on back)</p>	<p>WIND STRESS ON THE OCEAN OVER THE EASTERN CONTINENTAL SHELF OF NORTH AMERICA by Peter M. Saunders. Reprinted from the "Journal of Physical Oceanography, Vol. 7, No. 4, July 1977, pp. 555-566", July 1978. Prepared for the Office of Naval Research under Contract N00014-74-C-0262; NR 083-004.</p> <p>Employing one million ship reports gathered in the years 1941-72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1}. Atmospheric stratification is taken into account but its effect is shown to be small.</p> <p>In the summer season, the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons, the stress is directed toward the south and east being strongest in winter ($1-1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25-0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter, the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in</p> <p>(Cont. on back)</p>	<p>WIND STRESS ON THE OCEAN OVER THE EASTERN CONTINENTAL SHELF OF NORTH AMERICA by Peter M. Saunders. Reprinted from the "Journal of Physical Oceanography, Vol. 7, No. 4, July 1977, pp. 555-566", July 1978. Prepared for the Office of Naval Research under Contract N00014-74-C-0262; NR 083-004.</p> <p>Employing one million ship reports gathered in the years 1941-72 seasonal averages of the wind stress and its standard deviation have been computed for the shelf region of the eastern North American continent (out to a depth of 200 m). A drag coefficient is assumed which increases with wind speed, from 1.0×10^{-3} at 5 m s^{-1} to 2.3×10^{-3} at 25 m s^{-1}. Atmospheric stratification is taken into account but its effect is shown to be small.</p> <p>In the summer season, the 32-year climatological wind stress is toward the northeast, having a magnitude close to 0.25 dyn cm^{-2} throughout the entire shelf region. In the three other seasons, the stress is directed toward the south and east being strongest in winter ($1-1.5 \text{ dyn cm}^{-2}$) and weakest in fall ($0.25-0.5 \text{ dyn cm}^{-2}$). In addition to the expected increase in magnitude with increasing latitude remarkable small-scale variability occurs. An offshore increase in stress is widespread and dominates the mid-Atlantic Bight; in winter, the stress there increases from 0.5 to 1.0 dyn cm^{-2} in going 200 km offshore. In the Gulf of Maine and especially in the Gulf of St. Lawrence local maxima occur; the tail of the Grand Banks 500 km from shore shows a minimum. Probably much of this variation is associated with the intensity (and frequency) of cyclonic activity rather than directly with changes in</p> <p>(Cont. on back)</p>

friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.

friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.

friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.

friction at the underlying surface. Some oceanographic consequences are commented on but the computations are principally intended as a data source for further research.