WHOI-78-65

RAY CALCULATIONS OF OCEAN SOUND CHANNELS
USING A POCKET PROGRAMMABLE CALCULATOR
AND EXTENDED FORMS OF THE HIRSCH-CARTER
MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE
BETA FUNCTION

by

L. Baxter, II

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Woods Hole, Massachusetts 02543

October 1978

TECHNICAL REPORT

Prepared for the Office of Naval Research under
Contract N00014-77-C-0196.

Reproduction in whole or in part is permitted
for any purpose of the United States Government.
This report should be cited as: Woods Hole Oceano-
graphic Institution Technical Report WHOI-78-65.

Approved for public release; distribution
unlimited.

Approved for Distribution

Earl E. Hays

Earl E. Hays, Chairman
Department of Ocean Engineering

TABLE OF CONTENTS

I.	Introduction	<u>Page</u> 1
II.	Incomplete Beta-function and Calculator Programs for Acoustic Ray Computations	2
III.	The Geometry of Sound Speed Profile Layers in Which $c^2 = c_0^2 (1 - (\alpha z)^\beta)^{-1}$	3
IV.	Fitting Ocean Sound Speed Profiles Using Hirsch-Carter Type Layers	4
V.	Solutions for General Ray Segments in the Hirsch-Carter Model	7
VI.	Calculation vs Axial Angle of Range and Travel Time at the End of Loops Above and Below the Sound Channel Axis and at the End of a Complete Cycle	9
VII.	Calculation of Arrival Times for the Eigen Rays for a Source and Receiver	11
TABLE I	Travel Time at 705 km of rays of order 14, 15 and 16 in Sargasso Sea Profile	12
VIII.	Calculation of the Relative Intensity or Focusing Factor	13
IX.	Calculation of New Axial Angles of a Ray that Propagates from one Profile to Another	
X.	Calculation of Range Annotated Ray Angle Diagrams	
XI.	Notes on the Values of β in Asymmetric Profiles Based on the Hirsch-Carter Model	17
XII.	Acknowledgements	17

Supplement follows main report in this volume

ABSTRACT

Formulas for curve fitting and ray computation using compound models made up of several different layers of form $C^2 = C_0^2 (1 - |\alpha z|^\beta)^{-1}$ are presented. Examples of computation by pocket programmable calculator on two Sargasso Sea profiles, one from the center of a cold ring eddy are given. Necessary tables of the incomplete beta-function and calculator programs are included in a supplement.

RAY CALCULATIONS OF OCEAN SOUND CHANNELS
USING A POCKET PROGRAMMABLE CALCULATOR
AND EXTENDED FORMS OF THE HIRSCH-CARTER
MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE
BETA FUNCTION

L. Baxter, II

I. Introduction

Hirsch and Carter¹ have given closed form expressions for range and travel time of integral numbers of cycles of ray paths in the family of symmetrical profiles given by:

$$c^2 = c_o^2 \left(1 - |\alpha Z|^\beta \right)^{-1} \quad (1)$$

where c is the speed of sound at the vertical distance Z from the depth at which the speed is c_o and α and β are parameters. Pedersen and Gordon², Weinberg³, Stewart⁴, and others have developed the concept of fitting realistic acoustic profiles with layers of various curved profile segments while matching the speed and slope of the speed at the layer interfaces. This technique prevents the calculation of "false caustics" and other artifacts associated with less sophisticated profile fits and minimizes the number of layers needed to represent a natural sound speed profile realistically.

Equation 1 can be used with different parameters in each layer of a multilayer profile fit. The geometry of a ray in a layer may be understood by referring to Figure 1 in which a ray from the reference level ($c = c_o$) is refracted as the sound propagates through higher speed levels, and Z (always positive) is the absolute value of the depth difference from the reference level. With Z defined in this way and α always positive, Equation 1 may be rewritten as:

$$c^2 = c_o^2 \left(1 - (\alpha Z)^\beta \right)^{-1} \quad (1A)$$

The closed form expressions given by Hirsch and Carter¹ for range and travel time apply only to integral multiples of rays from the reference level

to the vertex, z_v , where the ray becomes horizontal. The portions of this path which we need to compute for rays that traverse several layers can however be expressed almost as simply in terms of incomplete beta-functions which have been tabulated⁵. Convergent series for computing them directly have also been published⁶.

For profile models consisting of no more than four layers of this type with no more than two layers on each side of the sound channel axis, it is not too difficult or tiresome to do ray computations with a medium capacity pocket programmable calculator and tables. I have done such calculations, fitting various natural asymmetric profiles with approximations consisting of two or three layers, using different parameters of Equation 1 in each layer and matching speed of sound and its derivative at the interfaces between layers. In this paper I outline the methods and give sample results for two profiles from the Sargasso Sea, one from the center of a cold eddy and one outside of any eddies. I also outline an approximate method for calculating rays that propagate through a small horizontal gradient of sound speed and a method of calculating range annotated ray angle diagrams.

II. Incomplete Beta-function and Calculator Programs for Acoustic Ray Computations

Although an extensive table⁶ of the incomplete beta-function is available, the most important range of the variables for our purpose is too sparsely covered. A supplement* to the present paper tabulates the necessary detail. The supplement also contains a Fortran program for generating any other values that may be required, and operating instructions and listings of the curve fitting and ray computation programs for the Texas Instruments SR56 calculator which I used. The calculator programs could be applied with little change to any equivalent or larger calculator

using algebraic operating system.

III. The Geometry of Sound Speed Profile Layers in Which $c^2 = c_o^2 (1 - (\alpha z)^\beta)^{-1}$

We need the slope dc/dz in order to match different layers at the interfaces. Differentiating Equation 1A, we have:

$$\frac{dc}{dz} = \frac{\alpha^\beta c_o (\alpha z)^{\beta-1}}{2 (1 - (\alpha z)^\beta)^{3/2}} \quad (2)$$

As we shall show later, the ray computations are simpler if we can fit the profile with layers in which the minimum speed of sound is equal to c_o and occurs at one interface. Therefore the limit of the slope, dc/dz , as z approaches zero is an important parameter. Remembering that

$\alpha > 0$ and $z \geq 0$ we have three cases:

Case 1. If $\beta > 1$, $dc/dz \rightarrow 0$ as $z \rightarrow 0$, regardless of the values of c_o and α .

Case 2. If $\beta = 1$, $dc/dz \rightarrow \alpha c_o / 2$ as $z \rightarrow 0$.

Case 3. If $\beta < 1$, $dc/dz \rightarrow \infty$ as $z \rightarrow 0$ regardless of the values of c_o and α .

For realistic sound speed profiles of ocean sound channels, Case 1 layers should be used to interface at the axis or minimum of the sound speed profile; the outer layers may belong to Case 2 or Case 3. This statement will be clarified by the following more detailed discussion of layer geometry for realistic values of α , β , c_o and z .

For refracted rays, the sound speed does not usually exceed about 102% of the axial speed of about 1.493 km/sec. The shape of the curves of Equations 1 and 2 in the range of the parameters for a 2% change in sound speed is most critically dependent on β , and reasonable changes in β may call for changes in α over a range of 10^{28} while changes in units

of Z , or depth variations of actual profiles, change α by much smaller ratios. To show the shape changes due to β on the same axes for various values (Figures 2 and 3), I use arbitrary units for Z with α adjusted to produce a maximum sound speed change of about 2% at $Z=1$. With these conventions, the order of magnitude of α is approximately that which would be realistic for Z kilometers.

Figure 2 shows the geometry of Equation 1 while Figure 3 shows that of Equation 2, i.e. the slope for the same values of the parameters. In these figures curves 1-6 belong to Case 3, curve 7 is Case 2 and curves 8-11 are Case 1. For Case 1 and Case 2 dc/dz increases with increasing Z , but in Case 2 the increase is not significant within the 2% change in sound speed. Case 2 approximates to a straight line and is the only case for which dc/dz at C_o is controlled by the parameter α . Case 3 layers are the only ones in which dc/dz decreases with increasing Z . They are somewhat more difficult to use because matching dc/dz to a lower velocity adjoining layer requires an interface at $Z > 0$. The process will be explained in the next section of this report.

IV. Fitting Ocean Sound Speed Profiles Using Hirsch-Carter Type Layers

We can now see that the conditions of Pedersen and Gordon², matching sound speed and slope, are met by asymmetrical profiles (see Figures 4 and 5) consisting of a Case 1 Hirsch-Carter type upper layer (designated by U) meeting a Case 1 lower layer (designated by L) at the sound channel axis. If the designations are used as subscripts and the subscript A refers to the axis of the sound channel $(C_o)_U = (C_o)_L = C_A$, but $\alpha_U \neq \alpha_L$ and $\beta_U \neq \beta_L$. The U and L layers must belong to Case 1 because only a zero value of dc/dz at $C_o = C_A$, or minimum sound velocity, can give a common tangent at $Z=0$.

Figures 4 and 5 illustrate sound speed profiles from the Sargasso Sea. The profile in Figure 4 is from the center of a cold ring eddy; that in Figure 5 is from a location undisturbed by the eddy. To fit each of these with a U and L layer I used a calculator program which iterates from a trial value of β to place a Hirsch-Carter type curve through C_0 at the axis and points 1 (C_1, Z_1) and 2 (C_2, Z_2) each marked with an X. The dots in Figures 4 and 5 indicate the resulting fit.

Where the fit is not perfect the exact values of α and β depend of course on the chosen points 1 and 2, and an equally good or better fit might appear from a different choice. It simplifies the calculations if β corresponds exactly to a tabulated value in the supplement or in Pearson⁵. Therefore it is worthwhile to try such a value, as close as possible to the calculated β . α is then recalculated to fit a point near the middle of the layer. The fit of the new values of α and β is checked over the measured profile. The values are adopted if the fit seems good enough.

The fit is improved if one does not try to cover too great a range of depths with one layer. The depth range can be subdivided into additional layers but meeting the conditions of equality of sound speed and its derivative are somewhat more complicated when the interface is not at the sound channel axis. The sound ray calculations also become more complicated because the rays must be divided into segments that traverse the various layers. Inspecting Figures 4 and 5 we see that the fit above the axis would be much improved by another layer. In Figures 6 and 7 an "M" layer above the U layer has been added to each of these profiles.

Both in the curve fitting and in the calculations, to be discussed later it will be useful to think of a separate space for each layer.

In Figure 1 the layer interfaces with adjoining layers may occur at Z_Q and Z_S , but the layer space and its coordinate system extends beyond the portion $Z_Q - Z_S$ that actually fits approximately to the real profile. The ray segments OZ_Q and $Z_S Z_V$ in the layer space are only auxilliary constructs for computing the segment $Z_Q Z_S$ which corresponds to the real ray in the layer. In the following discussion the subscripts U, M, or L are intended to indicate the space in which a coordinate is measured.

Figure 6 is an example of a profile that can be fitted with $\beta_M = 1$. $(c_o)_M$ is set equal to C_U at the chosen interface. $(dc/dz)_U$ at the interface is calculated from Equation 2. Then:

$$\alpha_M = 2 (dc/dz)_U / (c_o)_M \quad (3)$$

In Figure 7 $\beta_M < 1$. Since $(dc/dz)_M \rightarrow \infty$ as $Z_M \rightarrow 0$, slopes can be matched only if the reference velocity, $(c_o)_M$, is less than the velocity C_U at the interface and $Z_M > 0$ at the same place.

The fit was carried out as follows: The layer interface in U was chosen near a point of inflection of the empirical profile. C_U and $(dc/dz)_U$ at the interface were calculated. With some trial and error a layer thickness $(Z_M)_{max}$, and layer parameters β_M, α_M and $(c_o)_M$ were selected to approximate the curvature and maximum sound speed in the empirical profile. (Z_M) interface was then computed from $(dc/dz)_U = (dc/dz)_M$ interface using a program based on Equation 2. The program iterates from a trial value Z_t to a more accurate value of (Z_M) interface.

V. Solutions for General Ray Segments in the Hirsch-Carter Model

For our purposes it is useful to put the equations of Hirsch and Carter¹ in slightly different form. Within a layer described by Equation 1, a ray is designated by the angle θ_0 at velocity c_0 (see Figure 1). It vertexes at $Z = Z_v$ where

$$Z_v = (\sin \theta_0)^{2/\beta} / \alpha \quad (4)$$

The variable ξ defined by Hirsch and Carter¹ as

$$\xi = (\alpha Z)^\beta / \sin^2 \theta_0 \quad (5)$$

may also be expressed at Z by

$$\xi = \left(\frac{Z}{Z_v} \right)^\beta \quad (6)$$

The range, R_{0Z} , covered by the ray segment from 0 to Z, can be written:

$$\begin{aligned} R_{0Z} &= \frac{Z_v}{\beta \tan \theta_0} \int_0^\xi (1-x)^{\frac{1}{2}} x^{\frac{1}{\beta}-1} dx \\ &= \frac{Z_v}{\beta \tan \theta_0} \cdot B\left(\frac{1}{\beta}, \frac{1}{2}\right) \cdot I_\xi\left(\frac{1}{\beta}, \frac{1}{2}\right) \end{aligned} \quad (7)$$

where B is the complete beta function and I is the ratio of the incomplete to complete beta function. Let

$$B_1 = B\left(\frac{1}{\beta}, \frac{1}{2}\right) \quad \text{and} \quad I_1 = I_\xi\left(\frac{1}{\beta}, \frac{1}{2}\right)$$

then

$$R_{0Z} = Z_v \cdot B_1 \cdot I_1 / \beta \tan \theta_0 \quad (8)$$

The range, R_{ZZ_v} , can be written

$$R_{ZZ_v} = Z_v \cdot B(1-I_1) / \beta \tan \theta_o \quad (9)$$

Let $B_2 = B(1+\frac{1}{\beta}, \frac{1}{2})$ and $I_2 = I_\varphi(1+\frac{1}{\beta}, \frac{1}{2})$.

The travel times that correspond to the ranges of Equations 8 and 9 may be written

$$T_{CZ} = \frac{R_{CZ}}{c_o \cos \theta_o} \left\{ 1 - \sin^2 \theta_o \frac{I_2 B_2}{I_1 B_1} \right\} \quad (10)$$

and

$$T_{ZZ_v} = \frac{R_{ZZ_v}}{c_o \cos \theta_o} \left\{ 1 - \sin^2 \theta_o \frac{(1-I_2) B_2}{(1-I_1) B_1} \right\} \quad (11)$$

The values of the complete beta function, B_1 and B_2 are constants for a given layer of a profile. They may be calculated or taken at once from the tables. The relative values, I_1 and I_2 , of the incomplete beta function depend on θ_o and Z through Equations 4 and 6, and the tables. The range and travel time of any segment such as Q-S (Figure 1) is easily obtained as a difference between values computed by the above equations.

Programs for Equations 4, 6, 8, 9, 10 and 11 fit easily in the SR56 calculator when I_1 and I_2 are entered from tables. Note that if $Z = Z_v$, I_1 and $I_2=1$ and Equations 8 and 10 reduce to those given by Hirsch and Carter¹. To obtain total ranges for N axis crossings the values can of course be multiplied by $2N$ as is done by Hirsch and Carter.

VI. Calculation vs Axial Angle of Range and Travel Time at the End of Loops Above and Below the Sound Channel Axis and at the End of a Complete Cycle

Since the classical ray acoustics paper of Ewing and Worzell⁷, sound channel computations have often been presented by plots of range and travel time of loops above and below the axis and of a full ray cycle, all vs axial angle as the independent variable. These data are presented for the three-layer fits to the eddy and Sargasso Sea profiles in Figures 8, 9, 10 and 11. The procedure for calculating these plots is described: first for the simpler case of Figures 8 and 9 where $\beta_M = 1$, and then the modifications for Figures 10 and 11 where $\beta_M < 1$.

Axis to axis loops that do not penetrate into a second layer are computed by straight-forward application of Equations 4, 8 and 10 with the factor $2N$ equal to 2, I_1 and I_2 equal to 1, and θ_0 equal to the axial angle, θ_A . On the lower side of the axis where the profile fit has no second layer, the full range of axial angles may be covered this simply.

The Z coordinate of the interface in U layer space can be written $(Z_i)_U$. When $(Z_v)_U$ becomes greater than $(Z_i)_U$, the calculations can be simplified if θ_A is chosen so that γ is exactly equal to a value of X printed in the tables of the incomplete beta-function. Omitting the subscript U :

$$Z_v = Z_i / X^{1/\beta} \quad (12)$$

The axial angle, θ_A , for this ray is given by:

$$\theta_A = \arcsin \left[(\alpha Z_v)^{\beta/2} \right] \quad (13)$$

When $\beta = 1$, the reference sound speed, $(c_0)_M$, for the M layer is

equal to $(c_i)_U$, the sound speed at the interface. The reference angle in the M layer is calculated by

$$(\theta_o)_M = \arccos \left[\frac{(c_o)_M}{c_A} \cos \theta_A \right] \quad (14)$$

With θ_A and $(\theta_o)_M$ tabulated, one returns to the program for Equations 4, 8 and 10. Taking I_1 and I_2 directly without interpolation from the tables, range and travel time for the portion of the loop in the U layer is computed using the θ_A just found. I_1 and I_2 in the layer equal 1 in this case. The portion of the ray in the M layer is computed using the $(\theta_o)_M$ equivalent to θ_A from Equation 14. The values of range, travel time, and distance, Z_V , in the U and M layers are added to obtain the values plotted for the ray loop above the axis. These range and travel time values are added to those computed for the same θ_A below the axis to obtain the values for the full ray cycle.

When $\beta_M < 1$, as in Figures 10 and 11, the segments in M must be computed differently. $(c_o)_M \neq (c_i)_U$ and Equations 9 and 11 must be used instead of 8 and 10. I_1 and I_2 in the M layer are not equal to 1. One could use directly the $(\theta_o)_M$ that correspond to θ_A by Equation 14, but one would have to interpolate in the tables for I_1 and I_2 .

It is easier to defer the interpolation, doing it in $(\theta_o)_M$ at a later stage. This is done by using Equations 12 and 13 on the M layer after they have been used on the U layer. Equation 13 however is understood as:

$$(\theta_o)_{Mx} = \arcsin \left[(\alpha Z_V)^{\beta/2} \right]_M \quad (15)$$

where $(\theta_o)_{Mx}$ is the value of $(\theta_o)_M$ that corresponds to a tabulated value of X and not to θ_A . Range, travel time, and Z_v computed in the M layer for $(\theta_o)_{Mx}$ are interpolated to find the values for $(\theta_o)_M$ that do correspond to θ_A . The results are added to those for the U layer as before.

VII. Calculation of Arrival Times for the Eigen Rays for a Source and Receiver

This problem is merely an extension of the techniques used in Section VI. First one adds the appropriate segments to obtain a plot of range vs axial angle for the source and receiver depths and the possible types of path. Figure 12 is an example of this step. One interpolates to find the axial angles of each path at a given range. Then range and travel time are computed for these axial angles. Due to limited precision in the first interpolation the ranges will differ slightly but the average sound speeds will be correct for each path at the desired range. A second linear interpolation will adjust all the travel times to the correct range. Table I illustrates the result at a range of 705 km in Figure 12 and rays of order 14 through 16.

TABLE I

Travel Time at 705 km of rays of order 14, 15 and 16 in
Sargasso Sea Profile

	T	ΔT	N
7.16964	471.65950	0.0000000	16
7.21127	471.65254	0.0069642	16
7.40994	471.604778	0.054726	16
7.46963	471.59383	0.1001213	16
8.29557	471.438324	0.2211799	15
8.40314	471.414127	0.2453772	15
8.57449	471.36485	0.2946512	15
8.69705	471.33525	0.324255	15
9.70808	471.11683	0.54267	14
9.87478	471.068211	0.59129	14
10.03846	471.015474	0.64403	14
10.21703	470.959929	0.69958	14

VIII. Calculation of the Relative Intensity or Focusing Factor

a. Relative Intensity Except at Caustics

Brekhovskikh⁸ defines a "focusing factor" $f=I/I_0$, the ratio of the acoustic intensity I at a given point in the homogeneous medium to the acoustic intensity I_0 in a homogeneous medium at the same distance. He shows that when $R \gg Z$ and the point is not a caustic.

$$f = R / \sin \theta_p (dR/d\theta_A)_p \quad (16)$$

where θ_p is the horizontal angle at the given point and the derivative is evaluated for the ray that passes through the point.

$$\theta_p = \arccos (c(\cos \theta_A)/c_o) \quad (17)$$

$(dR/d\theta_A)_p$ may be obtained graphically as the slope from a plot like Figure 12.

b. Relative Intensity at a Caustic

In Figure 12 the four rays of a given order appear in two pairs. Each pair appears to join at a point for an axial angle slightly less than 7° . The scale is too coarse to show the detail in the neighborhood of the supposed point which is really the location of a caustic. Figure 13 shows the "point" of the lower pair of order 17 on a greatly expanded scale. The method of calculating this detail will be discussed after I outline its application.

We have been interested in comparing the relative intensity of caustics in differing profiles but at a given range. Although ordinary ray theory fails at a caustic, Brekhovskikh⁹ discusses a method of calculating intensity at a caustic from ray parameters. The full expression involves an Airy function and is rather complicated, but to compare the maxima of caustics under different conditions without computing the true relative

intensity at any point the expressions can be shortened. In the notation of this paper, and discarding factors that don't vary much in actual sound velocity profiles, relative intensity at a given large range and a given acoustic frequency is inversely proportional to $\tan \theta_A \sin \theta_P \left(d^2 R / d\theta_A^2 \right)_P^{2/3}$ where θ_P (see Equation 17) is the angle

with the horizontal of a ray tangent to the caustic. The method of computing the data for Figure 13 enables us to evaluate the derivative $\left(d^2 R / d\theta_A^2 \right)_P$; the other factors are obvious.

To calculate the range of a ray near the caustic, we measure, in Figure 10, the slope, S_f , and intercept I_f , of the full cycle ray that vertexes at the receiver depth. The values are: $S_f=3$ km/degree, $I_f=22$ km. We measure also the slope, S_u , and intercept I_u of the upper branch. $S_u=0$ km/degree. $I_u=10$ km. We then calculate range from the vertex vs axial angle for segments to the receiver depth of rays that vertex slightly shallower. The result appears as Figure 14. We note that the range increment, r_x , due to this segment is approximately the parabola

$$r_x^2 = K \left(\theta_A - (\theta_A)_P \right)^2 \quad (18)$$

where, in the given example, $K=2.9781$ and $(\theta_A)_P$, the axial angle of the ray that vertexes at the receiver depth, $=6.892$ degrees.

Let the angular difference, $[\theta_A - (\theta_A)_P] = \varphi$. Total range of a ray of order N in the vicinity of the vertex can be written as follows:

$$R = Q(I_u + S_u \cdot \varphi) + N(I_f + S_f \cdot \varphi) \pm K^{1/2} \varphi^{3/2} \quad (19)$$

where $Q = 3/2$ or $1/2$ depending on whether there is or is not an extra upper loop in the group of rays under consideration. Figure 13 is a plot

of Equation 19. As indicated by Brekhovskikh⁹ the caustic occurs where

$dR/d\theta_A = 0$, on the branch of the curve with the minus sign. Now:

$$dR/d\theta_A = dR/d\varphi = Q \cdot S_u + N \cdot S_f - K^{1/2} / 2 \varphi^{1/2} \quad (20)$$

Therefore, at the caustic

$$\varphi = K / 4 (Q \cdot S_u + N \cdot S_f)^2 \quad (21)$$

Differentiating Equation 20, we have

$$d^2R/d\theta_A^2 = K^{1/2} \varphi^{-3/2} / 4 \quad (22)$$

Evaluating Equation 22 at the caustic by substitution of Equation 21, we find that

$$(d^2R/d\theta_A^2) = 2 (Q \cdot S_u + N \cdot S_f)^3 / K \quad (23)$$

IX. Calculation of New Axial Angles of a Ray that Propagates from one Profile to Another

Milder¹⁰ has shown that if the change from one profile to another is sufficiently gradual, there is an invariant called the characteristic time. This invariant can be calculated by the equation:

$$J = (T - X \cdot \cos \theta_A / c_A) / 2 \pi \quad (24)$$

where J is the characteristic time, T is the full cycle travel time, X is the full cycle range, θ_A is the axial angle of the ray and c_A is the speed of sound at the axis. The conditions on the horizontal gradient for validity are given in detail by Milder for both wave and ray theory.

In ocean sound channels a horizontal gradient as small as .03 m/sec/km is safe.

Figure 15 is a plot of characteristic time vs axial angle for the

profiles that we have been considering. To find the angle in one profile that is equivalent to an angle in another one finds the value J corresponding to the angle, θ_{A_1} , in the first profile moves horizontally to the curve for the second profile and under the same J one finds the value, θ_{A_2} , in the second profile.

X. Calculation of Range Annotated Ray Angle Diagrams

Flatte¹¹ and Cox¹² have discussed the range annotated ray angle diagram and its applications. A program (for the pocket programmable calculator) adapted to generating data for such a diagram is included in the supplement. The depth difference Z from the reference level and the angle θ are computed from the range on a segment shorter than that from reference level to vertex. Longer paths are plotted by symmetry and addition. There are two cases: One where the given range is R_{OZ} in Equation 8, the other where the given range is R_{ZZ_v} in Equation 7. In the first case:

$$I_1 = R_{OZ} \beta (\tan \theta_o) / B_1 Z_v \quad (25)$$

In the second case:

$$I_1 = 1 - R_{ZZ_v} \beta (\tan \theta_o) / B_1 Z_v \quad (26)$$

The value of I_1 is used to obtain X from the I_x tables of the incomplete beta-function. Then

$$Z = Z_v X^{1/\beta} \quad (27)$$

and from Equation 1

$$c/c_o = (1 - |\alpha Z|^\beta)^{-1/2} \quad (28)$$

but

$$\theta = \arccos (c \cos \theta_0 / c_0)$$

or

$$\theta = \arccos \left(\cos \theta_0 / \sqrt{1 - |\alpha z|^\beta} \right) \quad (29)$$

XI. Notes on the Values of β in Asymmetric Profiles Based on the Hirsch-Carter Model

Hirsch and Carter have pointed out that, in symmetric models of the near axis sound transmission, the observed time dispersion of arrivals occurs only in that subset of the β family for which $1 < \beta < 2$. The actual sound channel, however, is grossly asymmetrical. Because the refraction below the axis is so much weaker than that above, rays at more than a very small axial angle will spend much more time below the axis than above it so that the overall dispersion pattern is like that of a symmetric channel with β near the below axis value of 1.25 or 1.26, the profiles above the axis are fitted, however, with values of β between 2 and 3. This would tend to reduce the dispersion below what one would get by reflection of the lower half of the channel.

XII. Acknowledgements

This work was supported by ONR Contract N00014-74-C-0262; NR 083-004. John C. Beckerle suggested the use of the Hirsch-Carter profile and did some preliminary calculations with it. P. Hirsch suggested use of the incomplete beta-function for computing ray segments. Additional helpful discussions with J. C. Beckerle, Earl E. Hays and George V. Frisk, are gratefully acknowledged.

REFERENCES

1. Peter Hirsch and Ashley H. Carter, "Mathematical Models for the Prediction of SOFAR Propagation Effects". J.Acoust.Soc.Am. 37, pp 90-94 (1965).
2. Melvin A. Pedersen and David F. Gordon, "Comparison of Curvilinear and Linear Profile Approximation in the Calculation of Underwater Sound Intensities by Ray Theory". J.Acoust.Soc.Am. 41, pp 419-438 (1967).
3. Henry Weinberg, "A Continuous Gradient Curve Fitting Technique for Acoustic Ray Analysis". J.Acoust.Soc.Am. 50, pp 975-984 (1971).
4. Kenneth R. Stewart, "Ray Acoustical Model of the Ocean Using a Depth/Sound-Speed Profile with a Continuous First Derivative". J.Acoust.Soc.Am. 38, pp 339-347 (1965).
5. E. S. Pearson and N.L. Johnson, "Tables of the Incomplete Beta-Function" 2nd Edition. Cambridge University Press (1968).
6. M. Abramowitz and I. A. Stegun, "Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables". U. S. Department of Commerce, National Bureau of Standards Applied Mathematical Series 55. Ninth printing, U. S. Government Printing Office, Washington, D.C. (1970).
7. Maurice Ewing and J. Lamar Worzel, "Long-Range Sound Transmission". Geological Soc. of Am., Memoir 27 (1948).
8. L. M. Brekhovskikh, "Waves in Layered Media". Academic Press, Inc., New York (1960), p.474.
9. L. M. Brekhovskikh. op.cit. pp 483-496.
10. D. Michael Milder, "Ray and Wave Invariants for SOFAR Channel Propagation". J.Acoust.Soc.Am. 46, -- 1259-1263 (1969).
11. Stanley M. Flatte, "Angle-depth diagram for use in underwater acoustics". J.Acoust.Soc.Am. 60, pp 1020-1023 (1976).
12. Henry Cox, "Approximate ray angle diagram". J.Acoust.Soc.Am. 61, pp 353-359 (1977).

CAPTIONS TO FIGURES

Figure 1 Path of a ray from reference level to vertex within a layer. Calculation of range and travel time of the segment from 0 to Z or that from Z to Z_v is discussed in the text. The general segment Q to S can be expressed as a difference of segments of either of the above types.

Figure 2 Geometry of sound speed profiles described by Equation 1A. $C_0=1.49275$ in all the curves. The other parameters are listed after the indicated number of each curve as follows:

1. $\alpha = 10^{-28}$, $\beta = .05$;
2. $\alpha = 10^{-14}$, $\beta = .10$;
3. $\alpha = 3 \times 10^{-6}$, $\beta = .25$;
4. $\alpha = 5.6 \times 10^{-5}$, $\beta = .33$;
5. $\alpha = 1.6 \times 10^{-3}$, $\beta = .50$;
6. $\alpha = 7.0 \times 10^{-3}$, $\beta = .65$;
7. $\alpha = .04$, $\beta = 1.0$;
8. $\alpha = .117$, $\beta = 1.5$;
9. $\alpha = .20$, $\beta = 2.0$;
10. $\alpha = .275$, $\beta = 2.5$;
11. $\alpha = .343$, $\beta = 3.0$.

Figure 3 Slopes of the sound speed profiles of Figure 2 on a logarithmic plot.

Figure 4 A sound speed profile from the center of a cold ring eddy is indicated by the continuous line. Two layers according to Equation 1 have been fitted at the axis and the points marked by X. The calculated sound speeds at other points are indicated by dots. The parameters are: $\alpha_v = 0.428668$, $\beta_v = 2.87968$, $C_0 = 1.48867$, $\alpha_L = .0325903$, and $\beta_L = 1.25033$

Figure 5 A sound speed profile in the Sargasso Sea outside of the eddy is indicated by the continuous line. Two layers according to Equation 1 have been fitted at the axis and at the points marked by X. The calculated sound speeds at other points are indicated by dots. The parameters are: $\alpha_v = .335904$, $\beta_v = 2.05506$, $C_0 = 1.49275$, $\alpha_L = .0321452$, and $\beta_L = 1.25931$

- Figure 6 The upper portion of the eddy profile, Figure 4, is fitted by two layers. The previous fit is retained below the sound channel axis. The new parameters are as follows:
 $\alpha_u = 0.44$, $\beta_u = 2.857$, $(c_o)_u = 1.48867$, $\alpha_m = 0.061$
 $\beta_m = 1.0000$, $(c_o)_m = 1.49600$
- Figure 7 The upper portion of the Sargasso Sea profile, Figure 5, is fitted by two layers. The previous fit is retained below the sound channel axis. The new parameters are as follows:
 $\alpha_u = 0.315$, $\beta_u = 2.000$, $(c_o)_u = 1.49275$, $\alpha_m = 4.01 \times 10^{-15}$,
 $\beta_m = 0.100$, $(c_o)_m = 1.4989$; $(z_i)_m$ at the interface = 0.0216,
 $(z_i)_u = 0.575$
- Figure 8 Range and vertex depth vs axial angle for the three layer fit of the eddy profile (Figures 4, 6). Range of a loop above the axis, one below the axis, and a full ray cycle are shown. Vertex depth is for the upper loop and is therefore the shallowest point reached by the ray.
- Figure 9 Travel time vs axial angle for the same profile, fit, and paths as Figure 8.
- Figure 10 Range and vertex depth vs axial angle for the three layer fit of the Sargasso Sea profile (Figures 5 and 7). The data presented corresponds to that presented in Figure 8. The constancy of range of the upper loop over the axial angles 0-10.4 is a property of the fit with $\beta = 2.0$ as noted in the Hirsch-Carter¹ paper.
- Figure 11 Travel time vs axial angle for the same profile, fit, and paths as Figure 10.
- Figure 12 Range vs axial angle of high order rays in the Sargasso Sea profile (Figures 5 and 7). The order is, of course, the number of loops below the axis. The receiver depth is .85 km. The source is on the axis. There are four rays belonging to each order. This figure may be used as described in the text to find axial angles of eigen rays at a given range.
- Figure 13 Range vs axial angle for two rays of the 17th order. This figure demonstrates the formation of a caustic as discussed in the text.

Figure 14 Range vs axial angle of a ray segment from vertex to receiver depth (.85 km, i.e. .4 km above the axis) in the Sargasso Sea profile (Figures 5 and 7). The solid line shows values calculated using the Hirsch-Carter model with tables of the incomplete beta-function. The circles show values from the parabolic fit, $r_x^2 = K (\theta_A - (\theta_A)_P)^2$, with $K = 2.9781$ and $(\theta_A)_P = 6.842^\circ$.

Figure 15 Characteristic time vs axial angle. This figure can be used for estimating the change in axial angle when a ray propagates through a transition region from one sound velocity profile to another.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Fig. 10

Fig. 11

Fig. 12

Fig. 13

Fig. 14

Fig. 15

Supplement to

RAY CALCULATIONS OF OCEAN SOUND CHANNELS
 USING A POCKET PROGRAMMABLE CALCULATOR
 AND EXTENDED FORMS OF THE HIRSCH-CARTER
 MATHEMATICAL MODEL WITH TABLES
 OF THE INCOMPLETE BETA-FUNCTION

L. Baxter, II

A. Tables of the Incomplete Beta-function

The complete beta-function, $B(p, q)$ of the variables p and q is given by

$$B(p, q) = \Gamma(p) \cdot \Gamma(q) / \Gamma(p+q) \quad (1)$$

or alternatively by:

$$B(p, q) = \int_0^1 x^{(p-1)} (1-x)^{(q-1)} dx \quad (2)$$

The incomplete beta-function, $B_x(p, q)$ is given by

$$B_x(p, q) = \int_0^x y^{(p-1)} (1-y)^{(q-1)} dy, \quad 0 < x < 1 \quad (3)$$

The function $I_x(p, q)$ sometimes called the relative incomplete beta-function is given by

$$I_x(p, q) = B_x(p, q) / B(p, q) \quad (4)$$

In these tables and in those of Pearson¹ I_x is given as a function of p, q and x , while $B(p, q)$ is tabulated at the top of each column of $I_x(p, q)$. In the present tables q is taken equal to .5, the only value it assumes in the Hirsch-Carter model equations.

In the paper to which this supplement is appended the Hirsch-Carter model equation is written

$$C^2 = C_0^2 (1 - |\alpha z|^\beta)^{-1} \quad (5)$$

and it is shown that range and travel time can be written for sound ray segments in terms of the following quantities

$$B_1 = B\left(\frac{1}{\beta}, .5\right) \quad (6)$$

$$I_1 = I_x\left(\frac{1}{\beta}, .5\right) \quad (7)$$

$$B_2 = B\left(1 + \frac{1}{\beta}, .5\right) \quad (8)$$

$$I_2 = I_x\left(1 + \frac{1}{\beta}, .5\right) \quad (9)$$

where $\chi = \left(z/z_v\right)^\beta \quad (10)$

and z_v is the value of z for which a sound ray vertexes (i.e. becomes horizontal at its maximum or minimum depth of excursion). To use these tables for sound ray calculations, B_1 is taken from the top of the column with $p = 1/\beta$; I_1 is taken opposite χ from the same column; B_2 is taken from the top of the column with $p = 1 + 1/\beta$; and I_2 is taken opposite χ from that column.

For some computations it may be necessary to enter the tables with a value of I_1 and interpolate to find a value of χ .

The $I_x(p, q)$ Function
 $q = 0.5$ $p = .20$ to $.30$

42-384
 MAR-1964

x	$p = .2$	$.25$	$.30$	x	$p = .2$	$.25$	$.30$
$B(p, q)$	6.268655		4.554444		6.268655		4.554444
.01	.3178035		.184054	.51	.7357336		.6443241
.02	.3653681		.2268620	.52	.7396286		.6493406
.03	.3965678		.2565072	.53	.7435044		.6543426
.04	.4204138		.2799599	.54	.7473630		.6593315
.05	.4399803		.2997001	.55	.7512063		.6643100
.06	.4567164		.3169302	.56	.7550366		.6692804
.07	.4714288		.3323333	.57	.7588556		.6742451
.08	.4846165		.3463399	.58	.7626655		.6792067
.09	.4966091		.3592383	.59	.7664682		.6841673
.10	.5076395		.3712348	.60	.7702656		.6891297
.11	.5178767		.3824811	.61	.7740599		.6940961
.12	.5274482		.3930920	.62	.7778531		.6990691
.13	.5364528		.4031581	.63	.7816472		.7040516
.14	.5449675		.4127504	.64	.7854444		.7090461
.15	.5530558		.4219277	.65	.7892470		.7140552
.16	.5607692		.4307382	.66	.7930571		.7190821
.17	.5681497		.4392212	.67	.7968770		.7241296
.18	.5752329		.4474108	.68	.8007092		.7292010
.19	.5820503		.4553367	.69	.8045565		.7342995
.20	.5886254		.4630221	.70	.8084209		.7394283
.21	.5949836		.4704901	.71	.8123058		.7445919
.22	.6011419		.4777586	.72	.8162140		.7497936
.23	.6071190		.4848451	.73	.8201486		.7550378
.24	.6129291		.4917640	.74	.8241130		.7603290
.25	.6185856		.4985279	.75	.8281112		.7656721
.26	.6241008		.5051492	.76	.8321468		.7710727
.27	.6294845		.5116379	.77	.8362240		.7765365
.28	.6347481		.5180048	.78	.8403481		.7820698
.29	.6398979		.5242568	.79	.8445238		.7876795
.30	.6449436		.5304035	.80	.8487573		.7933743
.31	.6498911		.5364509	.81	.8530544		.7991621
.32	.6547480		.5424062	.82	.8574232		.8050534
.33	.6595198		.5482757	.83	.8618715		.8110591
.34	.6642126		.5540657	.84	.8664085		.8171923
.35	.6688304		.5597800	.85	.8710454		.8234678
.36	.6733793		.5654249	.86	.8757945		.8299029
.37	.6778630		.5710042	.87	.8806710		.8365184
.38	.6822851		.5765225	.88	.8856927		.8433383
.39	.6866506		.5819842	.89	.8908808		.8503926
.40	.6909627		.5873926	.90	.8962620		.8577176
.41	.6952244		.5927516	.91	.9018698		.8653592
.42	.6994404		.5980653	.92	.9077468		.8733766
.43	.7036116		.6033359	.93	.9139497		.8818481
.44	.7077423		.6085674	.94	.9205669		.8908814
.45	.7118349		.6137625	.95	.9276810		.9006315
.46	.7158926		.6189235	.96	.9354948		.9113368
.47	.7199167		.6240546	.97	.9442899		.9233995
.48	.7239106		.6291571	.98	.9546366		.9376046
.49	.7278763		.6342343	.99	.9662007		.9559839
.50	.7318162		.6392893	1.00	1.0000000		1.0000000

$$B = 4.0566228$$

A	B
.350000	.5000000
N	K
25	9
X	I
.000100	.0280396
.000200	.0357387
.000300	.0411886
.000400	.0455524
.000500	.0492533
.000600	.0524994
.000700	.0554103
.000800	.0580623
.000900	.0605066

A	B
.350000	.5000000
N	K
25	9
X	I
.001000	.0627804
.002000	.0800277
.003000	.0922420
.004000	.1020267
.005000	.1103288
.006000	.1176139
.007000	.1241499
.008000	.1301070
.009000	.1356003

The $I_x(p, q)$ Function
 $q = 0.5$ $p = .35$ to $.45$

x	$p = .35$	$.40$	$.45$	x	$p = .35$	$.40$	$.45$
$B(p, q)$	4.0566228	3.6790923	3.3820539		4.0566228	3.6790923	3.3820539
.01	.1407124	.1078504	.0828482	.51	.6050637	.5692794	.5365068
.02	.1795816	.1425152	.1133517	.52	.6105126	.5750914	.5426226
.03	.2072370	.1678532	.1362559	.53	.6159500	.5808963	.5487374
.04	.2294953	.1886002	.1553352	.54	.6213784	.5866976	.5548538
.05	.2484703	.2065132	.1720190	.55	.6268009	.5924978	.5609750
.06	.2652006	.2224676	.1870295	.56	.6322194	.5982989	.5671022
.07	.2802834	.2369720	.2007909	.57	.6376363	.6041037	.5732393
.08	.2940971	.2503520	.2135768	.58	.6430547	.6099150	.5793885
.09	.3068985	.2628293	.2255753	.59	.6484768	.6157355	.5855528
.10	.3188713	.2745643	.2369230	.60	.6539054	.6215677	.5917344
.11	.3301514	.2856761	.2477220	.61	.6593431	.6274146	.5979367
.12	.3408430	.2962561	.2580513	.62	.6647923	.6332787	.6041629
.13	.3510282	.3063771	.2679737	.63	.6702564	.6391632	.6104158
.14	.3607711	.3160962	.2775390	.64	.6757377	.6450714	.6166986
.15	.3701264	.3254619	.2867891	.65	.6812396	.6510065	.6230150
.16	.3791373	.3345131	.2957588	.66	.6867654	.6569715	.6293678
.17	.3878407	.3432824	.3044766	.67	.6923178	.6629704	.6357617
.18	.3962678	.3517990	.3129677	.68	.6979008	.6690063	.6422001
.19	.4044461	.3600866	.3212536	.69	.7035174	.6750832	.6486871
.20	.4123976	.3681658	.3293527	.70	.7091722	.6812059	.6552272
.21	.4201434	.3760556	.3372816	.71	.7148689	.6873782	.6618254
.22	.4277001	.3837714	.3450539	.72	.7206117	.6936052	.6684865
.23	.4350841	.3913282	.3526835	.73	.7264055	.6998915	.6752161
.24	.4423097	.3987384	.3601815	.74	.7322553	.7062432	.6820199
.25	.4493878	.4060128	.3675573	.75	.7381666	.7126657	.6889045
.26	.4563307	.4131626	.3748212	.76	.7441452	.7191660	.6958767
.27	.4631477	.4201962	.3819810	.77	.7501978	.7257510	.7029445
.28	.4698485	.4271229	.3890450	.78	.7563317	.7324284	.7101169
.29	.4764413	.4339495	.3960196	.79	.7625540	.7392069	.7174021
.30	.4829331	.4406840	.4029116	.80	.7688748	.7460967	.7248112
.31	.4893316	.4473321	.4097267	.81	.7753026	.7531077	.7323560
.32	.4956425	.4539000	.4164705	.82	.7818495	.7602529	.7400497
.33	.5018722	.4603935	.4231485	.83	.7885278	.7675460	.7477973
.34	.5080266	.4668183	.4297656	.84	.7953520	.7750028	.7559460
.35	.5141098	.4731785	.4363254	.85	.8023385	.7826417	.7641864
.36	.5201274	.4794785	.4428332	.86	.8095069	.7904840	.7726508
.37	.5260835	.4857231	.4492924	.87	.8168805	.7985555	.7813678
.38	.5319828	.4919162	.4557069	.88	.8244864	.8068863	.7903700
.39	.5378288	.4980616	.4620807	.89	.8323582	.8155129	.7996970
.40	.5436252	.5041628	.4684167	.90	.8405365	.8244806	.8093985
.41	.5493761	.5102239	.4747186	.91	.8490732	.8338465	.8195361
.42	.5550854	.5162477	.4809893	.92	.8580347	.8436838	.8301899
.43	.5607549	.5222378	.4872322	.93	.8675088	.8540894	.8414656
.44	.5663887	.5281965	.4934502	.94	.8776165	.8651971	.8535082
.45	.5719893	.5341268	.4996456	.95	.8885322	.8771990	.8665273
.46	.5775608	.5400325	.5058219	.96	.9005239	.8903706	.8806948
.47	.5831097	.5459157	.5119815	.97	.9140428	.9052704	.8970028
.48	.5886244	.5517796	.5181271	.98	.9299712	.9228111	.9160601
.49	.5941222	.5576259	.5242614	.99	.9505906	.9455295	.9407555
.50	.5996014	.5634587	.5303868	1.00	1.0000000	1.0000000	1.0000000

The $I_x(p, q)$ Function
 $q = 0.5$ $p = .50$ to $.70$

42-384
 NATIONAL
 Bureau of

x	$p = .50$	$.60$	$.70$	x	$p = .50$	$.60$	$.70$
$B(p, q)$	3.141593	2.7745031	2.5057947		3.141593	2.7745031	2.5057947
.01	.0637686	.0379735	.0227433	.51	.5063666	.4527878	.4065981
.02	.0903345	.0576663	.0370236	.52	.5127358	.4595364	.4135910
.03	.1108247	.0736896	.0492781	.53	.5191101	.4663035	.4206160
.04	.1281884	.0877416	.0603989	.54	.5254920	.4730921	.4276764
.05	.1435663	.1005064	.0707605	.55	.5318843	.4799036	.4347748
.06	.1575424	.1123443	.0805656	.56	.5382895	.4867417	.4419128
.07	.1704634	.1234739	.0899398	.57	.5447103	.4936084	.4490941
.08	.1825549	.1340395	.0989678	.58	.5511494	.5005074	.4563214
.09	.1939734	.1441428	.1077098	.59	.5576098	.5074403	.4635974
.10	.2048328	.1538592	.1162111	.60	.5640942	.5144110	.4709253
.11	.2152190	.1632460	.1245067	.61	.5706057	.5214230	.4783086
.12	.2251989	.1723479	.1326241	.62	.5771474	.5284784	.4857498
.13	.2348255	.1812013	.1405862	.63	.5837226	.5355818	.4932539
.14	.2441418	.1898354	.1484110	.64	.5903345	.5427362	.5008236
.15	.2531833	.1982753	.1561146	.65	.5969867	.5499452	.5084635
.16	.2619798	.2065412	.1637101	.66	.6036829	.5572135	.5161774
.17	.2705563	.2146510	.1712089	.67	.6104271	.5645447	.5239704
.18	.2789343	.2226199	.1786208	.68	.6172233	.5719434	.5318468
.19	.2871326	.2304615	.1859550	.69	.6240760	.5794148	.5398124
.20	.2951672	.2381868	.1932187	.70	.6309899	.5869637	.5478718
.21	.3030525	.2458069	.2004193	.71	.6379699	.5945959	.5560322
.22	.3108011	.2533303	.2075625	.72	.6450216	.6023171	.5642993
.23	.3184242	.2607658	.2146543	.73	.6521506	.6101336	.5726806
.24	.3259319	.2681208	.2217001	.74	.6593633	.6180531	.5811834
.25	.3333333	.2754016	.2287039	.75	.6666667	.6260829	.5898160
.26	.3406367	.2826152	.2356706	.76	.6740681	.6342311	.5985883
.27	.3478494	.2897665	.2426040	.77	.6815758	.6425074	.6075099
.28	.3549784	.2968611	.2495079	.78	.6891989	.6509222	.6165923
.29	.3620301	.3039036	.2563858	.79	.6969475	.6594862	.6258479
.30	.3690101	.3108989	.2632412	.80	.7048328	.6682124	.6352907
.31	.3759240	.3178509	.2700768	.81	.7128674	.6771146	.6449362
.32	.3827767	.3247737	.2768956	.82	.7210657	.6862097	.6548027
.33	.3895729	.3316408	.2837010	.83	.7294437	.6955155	.6649100
.34	.3963171	.3384864	.2904952	.84	.7380202	.7050531	.6752819
.35	.4030133	.3453030	.2972809	.85	.7468167	.7148473	.6859450
.36	.4096655	.3520944	.3040609	.86	.7558582	.7249259	.6969310
.37	.4162774	.3588630	.3108370	.87	.7651745	.7353230	.7082773
.38	.4228526	.3656125	.3176118	.88	.7748011	.7460788	.7200286
.39	.4293943	.3723454	.3243879	.89	.7847810	.7572420	.7322387
.40	.4359058	.3790642	.3311675	.90	.7951672	.7688726	.7449745
.41	.4423902	.3857726	.3379527	.91	.8060266	.7810467	.7583203
.42	.4488506	.3924716	.3447459	.92	.8174451	.7938616	.7723840
.43	.4552897	.3991648	.3515493	.93	.8295366	.8074467	.7873092
.44	.4617105	.4058544	.3583647	.94	.8424576	.8219792	.8032923
.45	.4681157	.4125430	.3651944	.95	.8564337	.8377155	.8206178
.46	.4745080	.4192328	.3720406	.96	.8718116	.8550479	.8397213
.47	.4808899	.4259264	.3789058	.97	.8891753	.8746394	.8613367
.48	.4872642	.4326264	.3857918	.98	.9076655	.8977824	.8868971
.49	.4936334	.4393345	.3927010	.99	.9362314	.9278185	.9201049
.50	.5000000	.4460540	.3996360	1.00	1.0000000	1.0000000	1.0000000

The $I_x(p, q)$ Function
 $q = 0.5$
 $p = .80$ to 1.0

42-384
 NATIONAL
 Bureau of Standards

x	$p = .80$	$.90$	1.0	x	$p = .80$	$.90$	1.0
$B(p, q)$	2.2992875	2.1347606	2.0000000		2.2992875	2.1347606	2.0000000
.01	.0136863	.0082688	.0050126	.51	.3663955	.3311312	.3000000
.02	.0238828	.0154671	.0100505	.52	.3735271	.3383188	.3071797
.03	.0331087	.0223325	.0151142	.53	.3807054	.3455681	.3144345
.04	.0417718	.0290028	.0202041	.54	.3879334	.3528807	.3217670
.05	.0500507	.0355405	.0253206	.55	.3952132	.3602604	.3291796
.06	.0580445	.0419816	.0304640	.56	.4025477	.3677084	.3366750
.07	.0658163	.0483495	.0356349	.57	.4099399	.3752282	.3442561
.08	.0734091	.0546607	.0408337	.58	.4173921	.3828230	.3519259
.09	.0808541	.0609272	.0460608	.59	.4249077	.3904452	.3596876
.10	.0881753	.0671584	.0513167	.60	.4324898	.3982482	.3675445
.11	.0953912	.0733620	.0566019	.61	.4401416	.4060860	.3755002
.12	.1025169	.0795439	.0619168	.62	.4478667	.4140115	.3835586
.13	.1095647	.0857094	.0672621	.63	.4556689	.4220294	.3917237
.14	.1165446	.0918629	.0726382	.64	.4635524	.4301435	.4000000
.15	.1234656	.0980081	.0780456	.65	.4715214	.4383522	.4083920
.16	.1303352	.1041486	.0834849	.66	.4795797	.4466783	.4169048
.17	.1371599	.1102871	.0889566	.67	.4877332	.4551088	.4255437
.18	.1439455	.1164266	.0944615	.68	.4959861	.4636554	.4343146
.19	.1506971	.1225693	.1000000	.69	.5043447	.4723238	.4432236
.20	.1574192	.1287174	.1055728	.70	.5128148	.4811201	.4522774
.21	.1641163	.1348737	.1111806	.71	.5214022	.4900523	.4614835
.22	.1707919	.1410391	.1168239	.72	.5301144	.4991264	.4708497
.23	.1774496	.1472163	.1225036	.73	.5389596	.5083517	.4803848
.24	.1840928	.1534067	.1282202	.74	.5479448	.5177364	.4900980
.25	.1907242	.1596119	.1339746	.75	.5570799	.5272899	.5000000
.26	.1973468	.1658340	.1397675	.76	.5663755	.5370244	.5101021
.27	.2039632	.1720740	.1455996	.77	.5758410	.5469501	.5204168
.28	.2105759	.1783337	.1514719	.78	.5854903	.5570816	.5309524
.29	.2171872	.1846145	.1573850	.79	.5953357	.5674319	.5417424
.30	.2237997	.1909180	.1633400	.80	.6053935	.5780193	.5527864
.31	.2304150	.1972454	.1693376	.81	.6156796	.5888605	.5641101
.32	.2370357	.2035984	.1753789	.82	.6262141	.5999773	.5757359
.33	.2436638	.2099783	.1814647	.83	.6370200	.6113940	.5876894
.34	.2503012	.2163866	.1875962	.84	.6481212	.6231374	.6000000
.35	.2569497	.2228247	.1937742	.85	.6595480	.6352395	.6127017
.36	.2636115	.2292939	.2000000	.86	.6713347	.6477369	.6258343
.37	.2702882	.2357957	.2062746	.87	.6835220	.6606746	.6394449
.38	.2769818	.2423317	.2125992	.88	.6961580	.6741049	.6535898
.39	.2836944	.2489033	.2189750	.89	.7093040	.6880913	.6683375
.40	.2904274	.2555121	.2254033	.90	.7230307	.7027128	.6837722
.41	.2971830	.2621593	.2318854	.91	.7374305	.7180686	.7000000
.42	.3039628	.2688471	.2384227	.92	.7526220	.7342862	.7171573
.43	.3107689	.2755765	.2450166	.93	.7687612	.7515343	.7354249
.44	.3176032	.2823475	.2516685	.94	.7860636	.7700454	.7550510
.45	.3244675	.2891679	.2583802	.95	.8048385	.7901537	.7763932
.46	.3313637	.2960333	.2651531	.96	.8255625	.8123723	.8000000
.47	.3382940	.3029472	.2719890	.97	.8490350	.8375651	.8267749
.48	.3452601	.3099120	.2788897	.98	.8768218	.8674173	.8585786
.49	.3522643	.3169293	.2858572	.99	.9129584	.9062814	.9000000
.50	.3593085	.3240014	.2928932	1.00	1.0000000	1.0000000	1.0000000

The $I_x(p, q)$ Function
 $q = 0.5$ $p = 1.20$ to 1.30

42-384
 NATIONAL
 BUREAU OF STANDARDS

x	$p = 1.20$	1.25	1.30	x	$p = 1.20$	1.25	1.30
$B(p, q)$	1.791044		1.7079163		1.791044		1.7079163
.01	.0018574		.0011345	.51	.2477474		.2257117
.02	.0042789		.0028016	.52	.2547680		.2326015
.03	.0069799		.0047596	.53	.2618896		.2396036
.04	.0098854		.0069383	.54	.2691146		.2467207
.05	.0129573		.0093005	.55	.2764461		.2539558
.06	.0161721		.0118229	.56	.2838863		.2613124
.07	.0195142		.0144893	.57	.2914383		.2687922
.08	.0229719		.0172878	.58	.2991059		.2764006
.09	.0265367		.0202093	.59	.3068916		.2841390
.10	.0302018		.0232466	.60	.3147995		.2920125
.11	.0339619		.0263940	.61	.3228343		.3000247
.12	.0378126		.0296468	.62	.3309989		.3081808
.13	.0417505		.0330009	.63	.3392990		.3164848
.14	.0457724		.0364532	.64	.3477383		.3249422
.15	.0498761		.0400009	.65	.3563224		.3335584
.16	.0540595		.0436418	.66	.3650570		.3423386
.17	.0583208		.0473736	.67	.3739483		.3512894
.18	.0626586		.0511949	.68	.3830017		.3604180
.19	.0670718		.0551043	.69	.3922251		.3697312
.20	.0715591		.0591004	.70	.4016259		.3792372
.21	.0761202		.0631824	.71	.4112118		.3889443
.22	.0807540		.0673494	.72	.4209920		.3988621
.23	.0854602		.0716008	.73	.4309763		.4090011
.24	.0902387		.0759362	.74	.4411750		.4193720
.25	.0950887		.0803550	.75	.4515998		.4299871
.26	.1000105		.0848571	.76	.4622641		.4408605
.27	.1050039		.0894423	.77	.4731808		.4520065
.28	.1100691		.0941107	.78	.4843680		.4634424
.29	.1152061		.0988623	.79	.4958407		.4751861
.30	.1204153		.1036973	.80	.5076206		.4872589
.31	.1256968		.1086157	.81	.5197282		.4996834
.32	.1310512		.1136181	.82	.5321901		.5124872
.33	.1364789		.1187049	.83	.5450344		.5257003
.34	.1419806		.1238767	.84	.5582944		.5393571
.35	.1475568		.1291340	.85	.5720084		.5534986
.36	.1532084		.1344775	.86	.5862206		.5681710
.37	.1589358		.1399079	.87	.6009847		.5834306
.38	.1647403		.1454263	.88	.6163640		.5993444
.39	.1706225		.1510334	.89	.6324348		.6159931
.40	.1765838		.1567302	.90	.6492921		.6334761
.41	.1826252		.1625180	.91	.6670551		.6519184
.42	.1887478		.1683981	.92	.6858768		.6714818
.43	.1949528		.1743714	.93	.7059602		.6923794
.44	.2012416		.1804397	.94	.7275835		.7149035
.45	.2076158		.1866042	.95	.7511480		.7394756
.46	.2140768		.1928666	.96	.7772681		.7667414
.47	.2206264		.1992287	.97	.8069774		.7977864
.48	.2272663		.2056922	.98	.8422912		.8347259
.49	.2339981		.2122589	.99	.8884081		.8830156
.50	.2408242		.2189312	1.00	1.0000000		1.0000000

The $I_x(p, q)$ Function
 $q = 0.5$
 $p = 1.35$ to 1.45

42-384
 NATIONAL
 MADE IN U.S.A.

x	$p = 1.35$	1.40	1.45	x	$p = 1.35$	1.40	1.45
$B(p, q)$	1.6703740	1.6351522	1.6020257		1.6703740	1.6351522	1.6020257
.01	.0008874	.0006944	.0005436	.51	.2155571	.2059278	.1967939
.02	.0022686	.0018378	.0014895	.52	.2223721	.2126637	.2034436
.03	.0039333	.0032518	.0026896	.53	.2293046	.2195208	.2102207
.04	.0058170	.0048791	.0040942	.54	.2363578	.2265037	.2171289
.05	.0078854	.0066885	.0056757	.55	.2435353	.2336167	.2241715
.06	.0101162	.0086598	.0074161	.56	.2508384	.2408605	.2313510
.07	.0124942	.0107787	.0093026	.57	.2582709	.2482399	.2386709
.08	.0150080	.0130396	.0113255	.58	.2658378	.2557588	.2461355
.09	.0176488	.0154194	.0134773	.59	.2735407	.2634196	.2537482
.10	.0204095	.0179265	.0157521	.60	.2813853	.2712272	.2615131
.11	.0232847	.0205506	.0181450	.61	.2893745	.2791867	.2694353
.12	.0262696	.0232873	.0206520	.62	.2975137	.2873013	.2775186
.13	.0293605	.0261329	.0232696	.63	.3058068	.2955766	.2857687
.14	.0325541	.0290844	.0259952	.64	.3142600	.3040184	.2941917
.15	.0358477	.0321393	.0288262	.65	.3228786	.3126311	.3027925
.16	.0392390	.0352953	.0317608	.66	.3316685	.3214223	.3115772
.17	.0427260	.0385506	.0347971	.67	.3406358	.3303978	.3205535
.18	.0463070	.0419035	.0379337	.68	.3497876	.3395653	.3297280
.19	.0499808	.0453527	.0411695	.69	.3591318	.3489316	.3391093
.20	.0537461	.0488972	.0445033	.70	.3686761	.3585057	.3487052
.21	.0576019	.0525360	.0479344	.71	.3784298	.3682969	.3585258
.22	.0615473	.0562683	.0514621	.72	.3884017	.3783144	.3685805
.23	.0655819	.0600936	.0550858	.73	.3986034	.3885694	.3788804
.24	.0697051	.0640113	.0588053	.74	.4090455	.3990735	.3894376
.25	.0739163	.0680212	.0626202	.75	.4197404	.4098393	.4002657
.26	.0782157	.0721231	.0665303	.76	.4307030	.4208819	.4113792
.27	.0826028	.0763167	.0705357	.77	.4419478	.4322158	.4227939
.28	.0870777	.0806023	.0746365	.78	.4534726	.4438601	.4345284
.29	.0916405	.0849798	.0788327	.79	.4653559	.4558328	.4466020
.30	.0962913	.0894495	.0831247	.80	.4775573	.4681573	.4590377
.31	.1010304	.0940117	.0875128	.81	.4901259	.4808563	.4718592
.32	.1058583	.0986667	.0919974	.82	.5030844	.4939588	.4850969
.33	.1107752	.1034150	.0965791	.83	.5164652	.5049772	.4987826
.34	.1157818	.1082574	.1012586	.84	.5303036	.5215068	.5129538
.35	.1208788	.1131943	.1060364	.85	.5446417	.5360308	.5276538
.36	.1260666	.1182264	.1109135	.86	.5595267	.5511180	.5429326
.37	.1313462	.1233548	.1158905	.87	.5750166	.5668273	.5588512
.38	.1367186	.1285801	.1209688	.88	.5911802	.5832292	.5754806
.39	.1421845	.1339035	.1261490	.89	.6080991	.6004074	.5929075
.40	.1477450	.1393261	.1314325	.90	.6258759	.6184667	.6112385
.41	.1534014	.1448489	.1368206	.91	.6446393	.6375388	.6306080
.42	.1591549	.1504736	.1423145	.92	.6645540	.6577923	.6511885
.43	.1650068	.1562012	.1479156	.93	.6858379	.6794502	.6732085
.44	.1709585	.1620332	.1536256	.94	.7087910	.7028193	.6969807
.45	.1770114	.1679712	.1594459	.95	.7338446	.7283403	.7229561
.46	.1831674	.1740170	.1653785	.96	.7611591	.7566887	.7518244
.47	.1894282	.1801724	.1714252	.97	.7933456	.7890002	.7847745
.48	.1957954	.1864392	.1775877	.98	.8310675	.8274868	.8239784
.49	.2022713	.1928194	.1838686	.99	.8804061	.8778507	.8753457
.50	.2088571	.1993156	.1902697	1.00	1.0000000	1.0000000	1.0000000

The $I_x(p, q)$ Function
 $q = 0.5$
 $p = 1.50 \text{ to } 1.70$

42-384
 NATIONAL
 BUREAU OF STANDARDS

x	$p = 1.50$	1.60	1.70	x	$p = 1.50$	1.60	1.70
$B(p, q)$	1.5707965	1.51336534	1.46171357		1.5707965	1.51336534	1.46171357
.01	.0004257	.0002644	.0001607	.51	.1881204	.1720482	.1575098
.02	.0012077	.0007948	.0005238	.52	.1946807	.1784197	.1636832
.03	.0022255	.0015254	.0010470	.53	.2013737	.1849339	.1700056
.04	.0034369	.0024247	.0017130	.54	.2082024	.1915915	.1764812
.05	.0048182	.0034762	.0025114	.55	.2151699	.1983976	.1831132
.06	.0063536	.0046686	.0034352	.56	.2222797	.2053550	.1899049
.07	.0080318	.0059939	.0044792	.57	.2295352	.2124681	.1968598
.08	.0098443	.0074459	.0056395	.58	.2369403	.2197405	.2039849
.09	.0117844	.0090197	.0069129	.59	.2444990	.2271770	.2112819
.10	.0138468	.0107115	.0082971	.60	.2522155	.2347810	.2187560
.11	.0160272	.0125179	.0097900	.61	.2600945	.2425588	.2264131
.12	.0183220	.0144365	.0113901	.62	.2681408	.2505137	.2342590
.13	.0207281	.0164651	.0130960	.63	.2763598	.2586527	.2422987
.14	.0232430	.0186017	.0149068	.64	.2847570	.2669817	.2505393
.15	.0258646	.0208450	.0168215	.65	.2933384	.2755065	.2589867
.16	.0285911	.0231936	.0188396	.66	.3021105	.2842346	.2676487
.17	.0314210	.0256466	.0209605	.67	.3110804	.2931726	.2765328
.18	.0343530	.0282032	.0231840	.68	.3202554	.3023285	.2856469
.19	.0373861	.0308625	.0255098	.69	.3296437	.3117115	.2950010
.20	.0405193	.0336242	.0279379	.70	.3392541	.3213303	.3046035
.21	.0437521	.0364879	.0304683	.71	.3490960	.3311948	.3144661
.22	.0470837	.0394533	.0331010	.72	.3591800	.3413163	.3245995
.23	.0505139	.0425203	.0358364	.73	.3695172	.3517061	.3350160
.24	.0540424	.0456889	.0386747	.74	.3801201	.3623778	.3457297
.25	.0576689	.0489591	.0416163	.75	.3910022	.3733449	.3567551
.26	.0613934	.0523313	.0446617	.76	.4021785	.3846241	.3681090
.27	.0652160	.0558055	.0478114	.77	.4136655	.3962313	.3798093
.28	.0691369	.0593823	.0510660	.78	.4254815	.4081877	.3918763
.29	.0731562	.0630620	.0544264	.79	.4376470	.4205124	.4043317
.30	.0772743	.0668452	.0578931	.80	.4501849	.4332315	.4172013
.31	.0814916	.0707325	.0614671	.81	.4631209	.4463699	.4305125
.32	.0858087	.0747245	.0651494	.82	.4764843	.4599597	.4442974
.33	.0902262	.0788222	.0689408	.83	.4903085	.4740353	.4585929
.34	.0947447	.0830263	.0728427	.84	.5046316	.4886364	.4734396
.35	.0993650	.0873379	.0768559	.85	.5194980	.5038095	.4888867
.36	.1040880	.0917580	.0809819	.86	.5349594	.5196079	.5049890
.37	.1089147	.0962873	.0852219	.87	.5510771	.5360966	.5218147
.38	.1138459	.1009276	.0895772	.88	.5679242	.5533510	.5394423
.39	.1188830	.1056799	.0940495	.89	.5855812	.5714641	.5579674
.40	.1240271	.1105455	.0986403	.90	.6041813	.5905489	.5775089
.41	.1292794	.1155259	.1033511	.91	.6238377	.6107485	.5982147
.42	.1346415	.1206231	.1081840	.92	.6447345	.6322467	.6202759
.43	.1401147	.1258382	.1131406	.93	.6671049	.6552858	.6439439
.44	.1457008	.1311731	.1182229	.94	.6912688	.6801987	.6695614
.45	.1514014	.1366298	.1234331	.95	.7176856	.7074636	.6976339
.46	.1572183	.1422103	.1287733	.96	.7470601	.7378134	.7289120
.47	.1631535	.1479168	.1342458	.97	.7805761	.7724779	.7646744
.48	.1692091	.1537511	.1398531	.98	.8205388	.8138530	.8074039
.49	.1753872	.1597160	.1455976	.99	.8728886	.8681089	.8634939
.50	.1816901	.1658139	.1514823	1.00	1.0000000	1.0000000	1.0000000

the $I_x(p, q)$ Function
 $q = 0.5$ $P = 1.80$ to 2.00

42-384
 NATIONAL
 BUREAU OF STANDARDS

x	$P = 1.80$	1.90	2.00	x	$P = 1.80$	1.90	2.00
$B(p, q)$	1.41494618	1.3723461	1.3333333		1.41494618	1.3723461	1.3333333
.01	.0000989	.0000610	.0000376	.51	.1493350	.1323756	.1215000
.02	.0003457	.0002284	.0001510	.52	.1503035	.1381338	.1270464
.03	.0007195	.0004950	.0003409	.53	.1564277	.1440543	.1327597
.04	.0012116	.0008580	.0006082	.54	.1627111	.1501398	.1386441
.05	.0018166	.0013155	.0009536	.55	.1691588	.1563963	.1447040
.06	.0025307	.0018664	.0013779	.56	.1757737	.1628265	.1509441
.07	.0033513	.0025102	.0018821	.57	.1825609	.1694359	.1573691
.08	.0042765	.0032465	.0024670	.58	.1895243	.1762289	.1639844
.09	.0053046	.0040750	.0031335	.59	.1966690	.1832110	.1707954
.10	.0064347	.0049958	.0038825	.60	.2039998	.1903864	.1778078
.11	.0076658	.0060090	.0047150	.61	.2115223	.1977630	.1850278
.12	.0089973	.0071148	.0056319	.62	.2192430	.2053454	.1924618
.13	.0104287	.0083137	.0066341	.63	.2271671	.2131401	.2001167
.14	.0119599	.0096059	.0077228	.64	.2353022	.2211552	.2080000
.15	.0135906	.0109920	.0088990	.65	.2436540	.2293973	.2161194
.16	.0153208	.0124725	.0101636	.66	.2522317	.2378753	.2244834
.17	.0171506	.0140481	.0115180	.67	.2610427	.2465960	.2331009
.18	.0190802	.0157194	.0129630	.68	.2700951	.2555905	.2419815
.19	.0211098	.0174871	.0145000	.69	.2793995	.2648070	.2511357
.20	.0232399	.0193521	.0161301	.70	.2889651	.2743174	.2605745
.21	.0254708	.0213152	.0178545	.71	.2988036	.2841128	.2703102
.22	.0278031	.0233772	.0196745	.72	.3089266	.2942055	.2803556
.23	.0302373	.0255393	.0215915	.73	.3193473	.3046099	.2907252
.24	.0327742	.0278022	.0236066	.74	.3300791	.3153396	.3014343
.25	.0354142	.0301671	.0257214	.75	.3411386	.3264115	.3125000
.26	.0381585	.0326352	.0279372	.76	.3525430	.3378441	.3239408
.27	.0410077	.0352075	.0302556	.77	.3643104	.3496556	.3357773
.28	.0439627	.0378853	.0326779	.78	.3764628	.3618702	.3480322
.29	.0470246	.0406698	.0352059	.79	.3890221	.3745097	.3607307
.30	.0501944	.0435624	.0378410	.80	.4020156	.3876027	.3739010
.31	.0534732	.0455645	.0405847	.81	.4154717	.4011788	.3875757
.32	.0568622	.0476775	.0434395	.82	.4294238	.4152731	.4017877
.33	.0603626	.0529029	.0464064	.83	.4439105	.4299245	.4165806
.34	.0639758	.0562424	.0494875	.84	.4589740	.4451785	.4320000
.35	.0677031	.0596975	.0526847	.85	.4746650	.4610863	.4480999
.36	.0715462	.0632700	.0560000	.86	.4910412	.4777085	.4649430
.37	.0755063	.0669616	.0594354	.87	.5081726	.4951177	.4826034
.38	.0795853	.0707744	.0629931	.88	.5261414	.5133982	.5011694
.39	.0837942	.0747101	.0666752	.89	.5450468	.5326538	.5207477
.40	.0881067	.0787708	.0704840	.90	.5650113	.5530114	.5414697
.41	.0925527	.0825587	.0744219	.91	.5861892	.5746297	.5635000
.42	.0971250	.0872760	.0784915	.92	.6087779	.5977141	.5870496
.43	.1018255	.0917250	.0826951	.93	.6330386	.6225358	.6123974
.44	.1066565	.0963082	.0870356	.94	.6593288	.6494586	.6399250
.45	.1116202	.1010280	.0915157	.95	.6881628	.6790203	.6701800
.46	.1167193	.1058871	.0961383	.96	.7203268	.7120304	.7040000
.47	.1219559	.1108884	.1009064	.97	.7571400	.7498522	.7427905
.48	.1273328	.1160346	.1058233	.98	.8011712	.7951359	.7892822
.49	.1328531	.1213289	.1108922	.99	.8590292	.8547016	.8505000
.50	.1385196	.1267745	.1161165	1.00	1.0000000	1.0000000	1.0000000

5
NATIONAL
42-354
MAY 1955

The $I_x(p, q)$ Function

$q = 0.5$

Miscellaneous values of p corresponding to Profile Fits
used in this paper

x	$p = .347261$	$.7994692$	$.7994086$	x	$p = .347261$	$.7994692$	$.7994086$
$B(p, q)$	4.0803160	2.3002594	2.3101818		4.0803160	2.3002594	2.3101818
.01	.1427877	.0137231	.0141019	.51	.6071199	.3665951	.3686270
.02	.1818832	.0239382	.0245072	.52	.6125470	.3737252	.3757522
.03	.2096594	.0331782	.0338926	.53	.6179621	.3809040	.3829243
.04	.2319922	.0418532	.0426879	.54	.6233685	.3881313	.3901449
.05	.2510183	.0501422	.0510803	.55	.6287684	.3954104	.3974165
.06	.2677842	.0581450	.0591742	.56	.6341641	.4027442	.4047421
.07	.2828922	.0659247	.0670354	.57	.6395582	.4101353	.4121239
.08	.2967235	.0735247	.0747091	.58	.6449534	.4175867	.4195657
.09	.3095368	.0809764	.0822279	.59	.6503519	.4251009	.4270695
.10	.3215170	.0883037	.0896167	.60	.6557562	.4326820	.4346387
.11	.3328013	.0955252	.0968951	.61	.6611701	.4403325	.4422768
.12	.3434938	.1026560	.1040784	.62	.6665949	.4480568	.4499882
.13	.3536778	.1097087	.1111798	.63	.6720340	.4558575	.4577752
.14	.3634175	.1166930	.1182095	.64	.6774905	.4637395	.4656427
.15	.3727677	.1236182	.1251771	.65	.6829672	.4717069	.4735943
.16	.3817722	.1304916	.1320901	.66	.6884671	.4797636	.4816351
.17	.3904679	.1373200	.1389552	.67	.6939934	.4879155	.4897695
.18	.3988863	.1441089	.1457789	.68	.6995499	.4961667	.4980025
.19	.4070547	.1508637	.1525660	.69	.7051400	.5045234	.5063485
.20	.4149956	.1575889	.1593215	.70	.7107675	.5129913	.5147885
.21	.4227298	.1642887	.1660497	.71	.7164364	.5215768	.5233532
.22	.4302744	.1709670	.1727545	.72	.7221513	.5302869	.5320419
.23	.4376459	.1776272	.1794396	.73	.7279164	.5391296	.5408618
.24	.4448579	.1842726	.1861083	.74	.7337371	.5481128	.5498211
.25	.4519219	.1909061	.1927635	.75	.7396183	.5572455	.5589292
.26	.4588504	.1975307	.1994084	.76	.7455676	.5665383	.5681958
.27	.4656526	.2041490	.2060454	.77	.7515891	.5760014	.5776317
.28	.4723383	.2107636	.2126776	.78	.7576916	.5856477	.5872592
.29	.4789149	.2173766	.2193067	.79	.7638822	.5954907	.5970632
.30	.4853911	.2239904	.2259357	.80	.7701701	.6055452	.6070870
.31	.4917730	.2306071	.2325661	.81	.7765645	.6158283	.6173379
.32	.4980573	.2372290	.2392008	.82	.7830769	.6263596	.6278355
.33	.5042799	.2438582	.2458416	.83	.7897199	.6371616	.6386024
.34	.5104166	.2504966	.2524906	.84	.7965077	.6482593	.6496632
.35	.5164825	.2571461	.2591496	.85	.8034570	.6596825	.6610475
.36	.5224823	.2638087	.2658204	.86	.8105863	.6714650	.6727895
.37	.5284202	.2704863	.2725055	.87	.8179204	.6836480	.6849296
.38	.5343004	.2771806	.2792064	.88	.8254851	.6962802	.6975169
.39	.5401278	.2838936	.2859251	.89	.8333138	.7094208	.7106096
.40	.5459054	.2906271	.2926633	.90	.8414471	.7231429	.7242806
.41	.5516372	.2973831	.2994230	.91	.8499366	.7375373	.7386209
.42	.5573268	.3041633	.3062061	.92	.8588481	.7527231	.7537487
.43	.5629767	.3109696	.3130145	.93	.8682693	.7688563	.7698193
.44	.5685908	.3178040	.3198502	.94	.8783201	.7861518	.7870466
.45	.5741713	.3246684	.3267149	.95	.8891742	.8049197	.8057395
.46	.5797223	.3315645	.3336106	.96	.9010978	.8256348	.8263712
.47	.5852456	.3384948	.3405395	.97	.9145396	.8490936	.8497386
.48	.5907446	.3454606	.3475033	.98	.9303765	.8768733	.8773978
.49	.5962209	.3524644	.3545042	.99	.9508770	.9129950	.9133672
.50	.6016788	.3595085	.3615447	1.00	1.0000000	1.0000000	1.0000000

the $I_x(p, q)$ Function

$q=0.5$

Miscellaneous values of p corresponding to Profile fits used in this paper

x	$p=$ 1.347261	1.799469	1.794086	x	$p=$ 1.347261	1.799469	1.794086
$B(p, q)$	1.6723683	1.4151829	1.4175897		1.6723683	1.4151829	1.4175897
.01	.0008994	.0001015	.0001018	.51	.2160995	.1444021	.1450783
.02	.0022949	.0003534	.0003543	.52	.2229184	.1503714	.1510597
.03	.0039745	.0007340	.0007355	.53	.2298547	.1564965	.1571959
.04	.0058734	.0012341	.0012367	.54	.2369121	.1627814	.1634912
.05	.0079569	.0018480	.0018516	.55	.2440923	.1692297	.1699497
.06	.0102029	.0025720	.0025768	.56	.2513986	.1758463	.1765757
.07	.0125959	.0034032	.0034092	.57	.2588345	.1826336	.1833724
.08	.0151246	.0043396	.0043469	.58	.2664031	.1895930	.1903455
.09	.0177800	.0053796	.0053882	.59	.2741089	.1967434	.1974995
.10	.0205553	.0065219	.0065319	.60	.2819546	.2040750	.2048385
.11	.0234448	.0077658	.0077772	.61	.2899455	.2115980	.2123704
.12	.0264439	.0091104	.0091233	.62	.2980856	.2193196	.2200985
.13	.0295488	.0105553	.0105698	.63	.3063802	.2272443	.2280303
.14	.0327561	.0121002	.0121163	.64	.3148338	.2353803	.2361718
.15	.0360632	.0137450	.0137627	.65	.3234524	.2437328	.2445381
.16	.0394678	.0154895	.0155088	.66	.3322414	.2523106	.2531132
.17	.0429678	.0173338	.0173547	.67	.3412082	.2611220	.2619285
.18	.0465618	.0192781	.0193087	.68	.3503591	.2701749	.2709857
.19	.0502482	.0213226	.0213469	.69	.3597018	.2794799	.2802933
.20	.0540258	.0234676	.0234936	.70	.3692442	.2890458	.2898528
.21	.0578938	.0257136	.0257413	.71	.3789955	.2988842	.2997020
.22	.0618511	.0280610	.0280905	.72	.3889650	.3090073	.3098260
.23	.0658974	.0305105	.0305417	.73	.3991631	.3194278	.3202471
.24	.0700320	.0330626	.0330956	.74	.4096017	.3301599	.3309790
.25	.0742545	.0357181	.0357529	.75	.4202924	.3412192	.3420370
.26	.0785648	.0384777	.0385142	.76	.4312505	.3526232	.3534389
.27	.0829626	.0413422	.0413834	.77	.4424899	.3643901	.3652029
.28	.0874481	.0443126	.0443526	.78	.4540293	.3765423	.3773589
.29	.0920212	.0473898	.0474315	.79	.4658860	.3891014	.3899145
.30	.0966820	.0505749	.0506183	.80	.4780828	.4020941	.4028916
.31	.1014309	.0538689	.0539140	.81	.4906418	.4155500	.4163398
.32	.1062682	.0572730	.0573199	.82	.5035923	.4295012	.4302826
.33	.1111944	.0607885	.0608371	.83	.5169641	.4439865	.4447539
.34	.1162101	.0644166	.0644670	.84	.5307927	.4590490	.4598183
.35	.1213158	.0681589	.0682109	.85	.5451205	.4747386	.4754877
.36	.1265121	.0720166	.0720703	.86	.5599939	.4911138	.4918485
.37	.1318001	.0759913	.0760467	.87	.5754718	.5082437	.5089528
.38	.1371804	.0800848	.0801418	.88	.5916219	.5262103	.5269119
.39	.1426541	.0842985	.0843572	.89	.6085261	.5451139	.5457956
.40	.1482221	.0886344	.0886947	.90	.6262873	.5650762	.5657362
.41	.1538858	.0930944	.0931562	.91	.6450332	.5862517	.5868871
.42	.1596463	.0976804	.0977437	.92	.6649289	.6088380	.6094455
.43	.1655050	.1023944	.1024593	.93	.6861922	.6330959	.6336721
.44	.1714631	.1072387	.1073051	.94	.7091221	.6593826	.6599239
.45	.1775222	.1122154	.1122833	.95	.7341498	.6882126	.6887136
.46	.1836842	.1173272	.1173966	.96	.7619348	.7203716	.7208263
.47	.1899506	.1225765	.1226472	.97	.7935863	.7571795	.7575785
.48	.1963232	.1279657	.1280378	.98	.8312662	.8012036	.8015341
.49	.2028041	.1334980	.1335714	.99	.8805479	.8590524	.8592693
.50	.2093955	.1391760	.1392509	1.00	1.0000000	1.0000000	1.0000000

B. Fortran Programs for Calculating Tables of the Incomplete Beta-Function

Abramowitz and Stegun² give a series expansion for the I_x function which is equivalent to the following:

$$I_x(p, q) = \frac{x^p (1-x)^q}{p B(p, q)} \left[1 + \sum_{n=1}^{\infty} \frac{B(p+1, n)}{B(p+q, n)} x^n \right] \quad (11)$$

This converges well if $x < .5$.

For $.5 < x < 1$ the symmetry relation:

$$I_x(a, b) = 1 - I_{(1-x)}(b, a) \quad (12)$$

may be used to evaluate Equation 11 within its region of good convergence.

The following Fortran programs use equations 1, 11, and 12 and a polynomial approximation² to the gamma function to tabulate I_x .

The main program "TABLE" calls the other functions, accepts the input parameters, and prints the output. When TABLE is called it requests a logical unit number for the output with "ENTER IPRNT". When this is typed in, the line "ENTER A, B, N, X0, K" appears. Type in, in free field form, the quantities p, q, N, X0, K where p and q are the quantities we have been using (p and .5), N is the number of terms in the expansion (25 for 6 place accuracy), X0 is just smaller than the smallest desired value (X0=0 for a complete table column), and K equals the number of tabular entries desired in the column to be printed. When these are typed in, the line "ENTER DELX" appears. Type in the increment between successive values of x and the program will type out the table with

$$x = x_0 + I * DELX$$

with $I = 1$ to K

Finally, the line "NEW VALUES, YE or NO" appears. Type in NO to exit the program or YE to repeat.


```

FTN4, L
      PROGRAM TABLE
5005  FORMAT("ENTER IPRNT")
5000  FORMAT("ENTER A, B, N, XO, K")
5012  FORMAT(5X, I4, 10X, F14.7)
5010  FORMAT(5X, F9.6, 4X, F14.7)
5020  FORMAT(7X, "X", 20X, "I")
5030  FORMAT("NEW VALUES, YE OR NO")
5040  FORMAT(A2)
5045  FORMAT(7X, "A", 20X, "B")
5047  FORMAT(7X, "N", 20X, "K")
5060  FORMAT("ENTER DELX")
      IYES=2HYES
      IITTY=1
      IOTTY=1
      WRITE(IOTTY, 5005)
      READ(IITTY, *)IPRNT
100  CONTINUE
      WRITE(IOTTY, 5060)
      READ(IITTY, *)DELX
      WRITE(IOTTY, 5000)
      READ(IITTY, *)A, B, N, XO, K
      WRITE(IPRNT, 5045)
      WRITE(IPRNT, 5010)A, B
      WRITE(IPRNT, 5047)
      WRITE(IPRNT, 5048)N, K
5048  FORMAT(5X, I4, 10X, I14)
      WRITE(IPRNT, 5020)
      DO 200 I=1, K
      X=XO+DELX*I
      OUT=BI(X, A, B, N)
      WRITE(IPRNT, 5010)X, OUT
200  CONTINUE
      WRITE(IOTTY, 5030)
      READ(IITTY, 5040)IQUER
      IF(IQUER.EQ.IYES)GO TO 100
      END

```

```

      FUNCTION BI(X, A, B, N)
      XA=X**A
      XB=(1.0-X)**B
      FCIR=XA*XB/(A*B*ETA(A, B))
      IF(X.GT.0.50)FCIR=XA*XB/(B*B*ETA(A, B))
      BIN=1.0
      AP=A+1.0
      IF(X.GT.0.50)AP=B+1.0
      AB=A+B
      XQ=X
      IF(X.GT.0.50)XQ=1.0-X
      DO 10 I=1, N
      XI=XQ**I
      YI=I
      BE1=BETA(AP, YI)
      BE2=BETA(AB, YI)
      BIN=BIN+XI*BE1/BE2
10  CONTINUE
      BI=BIN*FCIR
      IF(X.GT.0.50)BI=1.0-BI
      RETURN
      END

```

```

FUNCTION BETA(P,Q)
PG=GAMMA(P)
QG=GAMMA(Q)
PQG=GAMMA(P+Q)
BETA=PG*QG/PQG
RETURN
END

```

```

FUNCTION GAMMA(A)
DOUBLE PRECISION AK(8),RG,AD,ALD,DJ,AI,AG
DATA AK/-.577191652D0,.988205891D0,
1 -.897056937D0,.918206857D0,-.756704078D0,
2 -.482199394D0,-.193527818D0,.035868343D0/
AD=A
AG=1.0
RG=1.0
IF(AD.GE.1.0D0)GO TO 19
DO 10 I=1,8
RG=RG+AD**I*AK(I)
10 CONTINUE
RG=RG/AD
CONTINUE
15 GAMMA=RG
RETURN
19 J=AD
IF(AD.EQ.1.0D0)J=0
IF(AD.EQ.2.0D0)J=1
DJ=J
ALD=AD-DJ
DO 25 I=1,8
RG=RG+ALD**I*AK(I)
25 CONTINUE
IF(AD.LE.2.0D0)GO TO 15
J=J-1
26 CONTINUE
AI=J
AG=AG*(ALD+AI)
J=J-1
IF(J.GT.0)GO TO 26
RG=RG*AG
GO TO 15
END
END$

```


C. Programs for Profile Fitting and Acoustic Ray Plotting in Long
Range Deep Ocean Sound Transmission Studies

The following programs written for the Texas Instruments SR56 programmable pocket calculator would be easily adaptable to other programmables using algebraic notation.

Tables C1A and C1B give operating instructions and a listing of a program that finds values of α and β for a Hirsch-Carter model fit to the three points: $C_0, D; C_1, Z_1$; and C_2, Z_2 . The program iterates from a trial value of β to find α and a more accurate value of β . The iteration can be continued to any desired precision. The operating steps 7 and 8 use the final values of α and β to calculate C as a function of Z for comparison with the empirical profile.

Tables C2A and C2B give operating instructions and a listing of a program that calculates slope (dc/dz) and sound speed as a function of Z in a Hirsch-Carter type profile with parameters α , β , and C_0 .

Tables C3A and C3B give operating instructions and a listing of a program that calculates the depth increment Z that corresponds to a given sound velocity in a Hirsch-Carter type profile with parameters α , β , and C_0 .

Tables C4A and C4B give operating instructions and a listing of a program that calculates the depth increment, Z , that corresponds to a given slope dc/dz , in a Hirsch-Carter type profile with parameters α , β , and C_0 . The program iterates from a trial value of Z to find a more accurate value. The iteration can be continued to any desired precision.

Tables C5A and C5B give operating instructions and a listing of a program that computes range and travel time of a ray segment or multiple ray segments in a Hirsch-Carter type profile. See Section V of the basic paper for more detail of the equations programmed. The parameters

TABLE C-1A HIRSCH-CARTER PROFILE CURVE FITTING

42-334

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS				Display
1	KEY IN PROGRAM		2nd CP	LRN			00 00
			ALL KEY	ENTRIES	IN TABLE	IB	98 00
			LRN	RST			0
2	PRELOAD REGISTERS ($P_t < \beta$ is a test value. Final value will replace it)	C_0	X^2	STO	1		C_0
		C_1	X^2	STO	2		C_1
		Z_1	STO	3			Z_1
		C_2	X^2	STO	7		C_2
		Z_2	STO	8			Z_2
		Δ	STO	9			Δ
		P_t	STO	4			
3	compute β (β will increment by Δ until value is just greater than ideal. Pauses show each test.)		R/S				β_1
SEE NOTE							β_2
							β_3
							β_4
							
							β
4	CYCLE FOR NEXT SIGNIFICANT DIGIT		—	RCL	9	=	$\beta - \Delta$
			STO	4	RCL	9	
			$\div$	1	0	=	$\Delta_n = \Delta/10$
			STO	9	RST	R/S	β
	The same as STEP 3						β_1
							β_2
							β_3
5	REPEAT 3,4 UNTIL β has ENOUGH DIGITS						
6	COMPUTE α		RCL	6	$2nd \sqrt{y}$		
			RCL	4	=		α
7	CALCULATE SVP FOR VARIOUS Z_a , Z_b , etc.	Z_a	GTO	7	7	R/S	C_a
		Z_b	R/S				C_b
		Z_c	R/S				C_c
		Z_d	R/S				C_d
8	REPEAT FITTING PROCESS FOR OTHER BRANCH OF SVP (LOOPS BACK TO STEP 2)		RST				

Note: if β does not change from β_t , either β_t is already larger than β , or trial points 1+2 are in the wrong order in memory. To reverse trial points in memory, key in GTO 57 R/S.

TABLE C-1B HIRSCH-CARTER PROFILE CURVE FITTING

Register Contents									
0	1	2	3	4	5	6	7	8	9
C_0^2	C_1^2	Z_1	β	α^P	α^P_{prev}	C_2^2	Z_2	Δ	
Preloaded					Program				
Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments		
			MAIN LOOP						
00	57	2nd subr		50	01	1	C_0^2		
01	03	3	to	51	94	=	$C_2^2 - C_0^2$		
02	04	4	calculate	52	30	2nd PROD			
03	33	STO		53	05	5	α^P		
04	06	6	α^P from C_1, Z_1	54	34	RCL			
05	57	2nd subr		55	05	5	α^P		
06	05	5	to rotate	56	58	2nd rtn			
07	07	7	registers	57	34	RCL	This subr		
08	57	2nd subr	2-7, 3-8	58	02	2	rotates		
09	03	3	α^P from C_2, Z_2	59	39	2nd EXC	registers		
10	04	4		60	07	7	2-7,		
11	54	$\div$		61	33	STO	3-8		
12	34	RCL		62	02	2			
13	06	6		63	34	RCL			
14	74	-		64	03	3			
15	01	1		65	39	2nd EXC			
16	94	=	$\alpha_2^P / \alpha_1^P - 1$	66	08	8			
17	56	2nd CP	clear test reg.	67	33	STO			
18	47	2nd $X \geq t$	if $\alpha_2^P > \alpha_1^P$	68	03	3			
19	07	7	Go to 70	69	58	2nd rtn			
20	00	0		70	32	$X \geq t$			
21	34	RCL		71	57	2nd subr			
22	04	4	β	72	05	5	restore orig.		
23	84	+		73	07	7	pos of register		
24	34	RCL		74	34	RCL			
25	09	9	Δ	75	04	4	β to x reg.		
26	94	=	new $\beta = \beta + \Delta$	76	41	R/S			
27	59	2nd Pause	display β	77	45	2^x	2nd LOOP		
28	33	STO	store new β	78	34	RCL	with Z in X		
29	04	4		79	04	4	reg. calculat		
30	57	2nd subr	to rotate registers	80	64	X	C		
31	05	5	to original	81	34	RCL			
32	07	7	position	82	06	6			
33	42	RST	TO NEXT MAIN LOOP	83	94	=			
34	34	RCL	THIS SUBR	84	93	+/-			
35	03	3	CALCULATES α^P	85	84	+			
36	45	4^x		86	01	1			
37	34	RCL		87	94	=			
38	04	4		88	54	$\div$			
39	94	=	Z_1^P	89	34	RCL			
40	20	2nd $1/x$		90	01	1			
41	33	STO	$1/Z_1^P$ in 5	91	94	=			
42	05	5		92	48	2nd Vx			
43	34	RCL		93	20	2nd $1/x$			
44	02	2	C_1^2	94	41	R/S	C		
45	12	INV		95	22	GTO	2nd LOOP To		
46	30	2nd PROD	$1/C_1^2 Z_1^P$ in 5	96	07	7	calculate next		
47	05	5		97	07	7	C		
48	74	-		98					
49	34	RCL		99					

TABLE C-2 A dc/dz & c OF HIRSCH-CARTER PROFILE

PI
NATIONAL
42-394
REVISED

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS				Display
1	KEY IN PROGRAM		2nd CP	LRN			00 00
			ALL KEY	ENTRIES	IN		
			TABLE	2B			75 00
			LRN	RST			0
2	LOAD REGISTERS	α	R/S				α
		β	R/S				β
		c_0	R/S				c_0
3	CALCULATE dc/dz	Z	R/S				dc/dz
4	CALCULATE c		R/S				c
	REPEAT STEPS 3, 4 WITH NEW Z AS DESIRED						
5	TO CHANGE ALL PARAMETERS CONTINUE AT STEP 2		RST				

TABLE C-2 B dc/dz & c OF HIRSCH-CARTER PROFILE

Register Contents

0	1	2	3	4	5	6	7	8	9
	α	β	C_0	Z	αZ				

Program

Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments
			LOAD REGISTERS				
00	33	STO	$\leftarrow \alpha$	39	34	RCL	
01	01	1		40	05	5	
02	41	R/S	$\leftarrow \beta$	41	45	y^x	
03	33	STO		42	34	RCL	
04	02	2		43	02	2	
05	41	R/S	$\leftarrow C_0$	44	53	)	
06	33	STO		45	45	y^x	
07	03	3		46	01	1	
08	41	R/S	$\leftarrow Z$	47	92	.	
09	33	STO	MAIN	48	05	5	
10	04	4	LOOP	49	94	=	
11	64	X	THIS SECTION	50	41	R/S	$dc/dz-$
12	34	RCL	CALCULATES	51	34	RCL	THIS
13	01	1	dc/dz	52	04	4	SECTION
14	94	=		53	64	X	CALCULATE
15	33	STO		54	34	RCL	C
16	05	5		55	01	1	
17	45	y^x		56	94	=	
18	52	(		57	45	y^x	
19	34	RCL		58	34	RCL	
20	02	2		59	02	2	
21	74	-		60	74	-	
22	01	1		61	01	1	
23	53	)		62	94	=	
24	64	X		63	93	+/-	
25	34	RCL		64	20	2nd $1/x$	
26	03	3		65	64	X	
27	64	X		66	34	RCL	
28	34	RCL		67	03	3	
29	02	2		68	43	x^2	
30	64	X		69	94	=	
31	34	RCL		70	48	2nd $\sqrt{x}$	
32	01	1		71	41	R/S	$C \rightarrow$
33	54	$\div$		72	22	GTO	$\leftarrow Z$
34	02	2		73	00	0	RETURN TO
35	54	$\div$		74	09	9	MAIN LOOP
36	52	(		75			
37	01	1		76			
38	74	-		77			

TABLE C-3A Depth for a given sound velocity. Hirsch-Carter Profile

NATIONAL
42-384
Rev. 1-10-66

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS	Display
1	KEY IN PROGRAM		2nd CP LRN	00 00
			ALL ENTRIES IN	
			TABLE C-3B	32 00
			LRN RST	0
2	LOAD REGISTERS	α	R/S	α
		β	R/S	β
		C_0	R/S	C_0
3	CALCULATE	C	R/S	Z
4	REPEAT STEP 3 AS DESIRED			
5	TO LOAD NEW CONSTANTS AND REPEAT STEPS 2, 3		RST	

TABLE C-3B Depth for a given sound velocity. Hirsch-Carter Profile

NATIONAL
42-384
Rev. 1-10-66

Register Contents

0	1	2	3	4	5	6	7	8	9
	α	β	C_0						

Program

Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments
			LOAD REGISTERS	17	94	=	
00	33	STO	$\leftarrow \alpha$	18	93	+/-	$1 - C_0^2 / C^2$
01	01	1		19	45	$\times$	
02	41	R/S	$\leftarrow \beta$	20	34	RCL	
03	33	STO		21	02	2	
04	02	2		22	20	2nd 1/x	$(1 - C_0^2 / C^2)^{1/2}$
05	41	R/S	$\leftarrow C_0$	23	94	=	
06	33	STO		24	54	$\div$	
07	03	3		25	34	RCL	
08	41	R/S	$\leftarrow C$	26	01	1	
09	20	2nd 1/x	MAIN LOOP	27	94	=	
10	64	X		28	41	R/S	Z $\rightarrow$
11	34	RCL		29	22	GTO	$\leftarrow C$
12	03	3		30	00	0	RETURN TO MAIN LOOP
13	94	=		31	09	9	
14	43	$\times^2$	C_0^2 / C^2				
15	74	-					
16	01	1					

TABLE C-4A Depth for a given slope. Hirsch-Carter Profile.

NATIONAL
42-334
Made in U.S.A.

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS				Display
1	KEY IN PROGRAM		2nd CP	LRN			00 00
			ALL KEY	ENTRIES	IN		
			Table	B-4B			70 00
			LRN	RST			0
2	LOAD REGISTERS	α	R/S				α
		β	R/S				β
		C_0	R/S				C_0
3	Put dc/dz in t register	dc/dz	$\times \rightarrow t$				0
4	Initialize ΔZ , Z_t & START	ΔZ	STO	6			ΔZ
		Z_t	R/S				
5	PROGRAM PAUSES AT EACH NEW TRIAL Z HALTS AT FIRST Z with dc/dz G.E. t						Z_1 Z_2 Z_3 Z
6	if desired restart for more accurate Z		—	RCL	6	=	$dc/dz *$
			2nd EXC	4			
			RCL	6	$\div$	1	$\Delta Z / 10$
			0	=			
			STO	6	RCL	4	
			R/S				
	RECYCLES TO STEP 5						
	* dc/dz here is trial value from last run						

TABLE C-4B Depth for a given slope. Hirsch-Carter Profile

Register Contents									
0	1	2	3	4	5	6	7	8	9
α	β	C_0	Z	αZ	ΔZ				
Program									
Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments		
			LOAD REGISTERS	36	52	(			
00	33	STO	$\leftarrow \alpha$	37	01	1			
01	01	1		38	74	-			
02	41	R/S	$\leftarrow \beta$	39	34	RCL			
03	33	STO		40	05	5			
04	02	2		41	45	y^x			
05	41	R/S	$\leftarrow C_0$	42	34	RCL			
06	33	STO		43	02	2			
07	03	3		44	53	)		$(1-(\alpha Z)^p)$	
08	41	R/S	put dc/dz in t register	45	45	y^x			
			ΔZ in 6	46	01	1			
			Initiate MAIN LOOP with trial Z	47	92	.			
			$\leftarrow Z_t$	48	05	5		1.5	
				49	94	=		dc/dz	
09	33	STO		50	12	INV			
10	04	4		51	47	2nd $X \geq t$		test whether to iterate Z	
11	64	X		52	06	6			
12	34	RCL		53	00	0		on final iteration	
13	01	1		54	39	2nd EXC			
14	94	=	αZ_t	55	04	4			
15	33	STO		56	41	R/S		$Z \rightarrow$	
16	05	5						{ NEW ΔZ	
17	45	y^x						{ trial Z	
18	52	(						REPEAT FOR MORE ACCURATE Z	
19	34	RCL		57	22	GTO			
20	02	2		58	00	0			
21	74	-		59	09	9			
22	01	1							
23	53	)		60	34	RCL			
24	64	X	$(\alpha Z)^{p-1}$	61	04	4		Z_t	
25	34	RCL		62	84	+			
26	03	3		63	34	RCL		ΔZ	
27	64	X		64	06	6			
28	34	RCL		65	94	=			
29	02	2		66	59	2nd Pause		$Z + \Delta Z$	
30	64	X		67	22	GTO		to MAIN LOOP for next iteration	
31	34	RCL		68	00	0			
32	01	1		69	09	9			
33	54	$\div$							
34	02	2							
35	54	$\div$	$\alpha p C_0 (\alpha Z)^{p-1} / 2$						

α , β , c_0 and values B_1 and B_2 of the beta-function (see Section 1 of this supplement are needed for initialization. If the segment is not an integral multiple of the path from reference level to vertex, values of I_1 and I_2 must be entered from a table of the incomplete beta-function (section 1 of this supplement). Otherwise $I_1 = I_2 = 1$. The program recycles for each new value of the reference angle θ_0 , which in a case 1 profile, fitted at the axis, is the axial angle, θ_A .

Tables C6A and C6B give operating instructions and a listing of a program that calculates angles θ_0 and θ_j at points c_0 , O and C_j , Z_j of a Hirsch-Carter type profile with parameters α , β . The angles are calculated for given values of the X parameter of the incomplete beta-function to avoid interpolation in the tables when using the previous program (C5A,C5B) in this section.

Tables C7A and C7B give operating instructions and a listing of a program that calculates the characteristic time J (see Milder³ and section IX of the basic paper) of a sound ray path when the axial sound speed c_0 , the axial angle θ_A , the full cycle range X and the full cycle travel time T are known. The characteristic time is needed to convert axial angles in one profile to those in the next when a ray propagates through a horizontal gradient.

Tables C8A and C8B give operating instructions and a listing of a program that calculates ray angle with the horizontal, and depth increment, of a ray in a Hirsch-Carter type profile. The independent variable is range, which may be specified from an axis crossing or a vertex. The program is initialized with values of α , β , c_0 , and the beta-function, $B(\frac{1}{\beta}, \frac{1}{2})$. The ray is designated by its reference angle θ_0 ,

TABLE C-5A Ray plot by Hirsch-Carter profile

NOTE: IF LOC 86 is 2nd rtn Computes Equations 8-10
 IF. LOC 86 is (computes equations 9-11
 of the main report

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS	Display
1	KEY IN PROGRAM		2nd CP LRN ALL KEY EUTRIES IN TABLE C-5B LRN RST	00 00 93 00
2	PRELOAD 2N = 1 single segment 2N = 2 axis to axis or even number for higher order	2N	STO 6	2N
3	LOAD REGISTERS	α β B_1 B_2 C_0	R/S R/S R/S R/S R/S	α 2/ β B_1 B_2 C_0
4	RUN MAIN LOOP	θ I_1 I_2	R/S R/S R/S R/S	Z_v R T
5	REPEAT STEP 4 AS DESIRED WITH ANY θ			
6	TO CHANGE CONSTANTS AND REPEAT STEPS 2, 5		RST	

TABLE C-5B Ray Plot by Hirsch-Carter Profile
Register Contents

0	1	2	3	4	5	6	7	8	9
θ	$B_1(1-I_1)$	α	β	c_0	$2 \times N$	$2/\beta$	B_1	B_2	
or $B_1 I_1$ Program									
Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments		
			LOAD REGISTERS	49	54	$\div$			
00	33	STO	$\leftarrow \alpha$	50	34	RCL			
01	03	3		51	01	1			
02	41	R/S	$\leftarrow \beta$	52	25	tan			
03	33	STO		53	94	=			
04	04	4		54	41	R/S	R $\rightarrow$		
05	20	2nd 1/x		55	54	$\div$			
06	64	X		56	34	RCL			
07	02	2		57	05	5			
08	94	=	$2/\beta$	58	54	$\div$			
09	33	STO		59	34	RCL			
10	07	7		60	01	1			
11	41	R/S	$\leftarrow B_1$	61	24	cos			
12	33	STO		62	64	X			
13	08	8		63	52	(			
14	41	R/S	$\leftarrow B_2$	64	01	1			
15	33	STO		65	74	-			
16	09	9		66	34	RCL			
17	41	R/S	$\leftarrow c_0$	67	01	1			
18	33	STO		68	23	sin			
19	05	5		69	43	X^2			
20	41	R/S		70	54	$\div$			
			MAIN LOOP	71	34	RCL			
21	33	STO	$\leftarrow \theta$	72	02	2			
22	01	1		73	64	X			
23	23	sin		74	41	R/S	$\leftarrow I_2$		
24	45	y^x		75	57	2nd subr	for choice		
25	34	RCL		76	08	8	of ray segmen		
26	07	7		77	06	6			
27	54	$\div$		78	64	X			
28	34	RCL		79	34	RCL			
29	03	3		80	09	9			
30	94	=		81	94	=			
31	64	X		82	41	R/S	T $\rightarrow$		
32	52	(					$\leftarrow \theta$		
33	41	R/S	$Z_v \rightarrow$	83	22	GTO	RETURN TO		
			$\leftarrow I_1$	84	02	2	BEGIN MAIN		
34	57	2nd subr	for choice of	85	01	1	LOOP AGAIN		
35	08	8	ray segment						
36	06	6		86	58	2nd rtn	for segment		
37	64	X					ref. level to		
38	34	RCL					intermediate		
39	08	8					depth or		
40	53	)	$B_1(1-I_1)$	86	52	(	For segment		
41	33	STO		87	51	CE	intermediate		
42	02	2		88	74	-	level to		
43	64	X		89	01	1	vertex		
44	34	RCL		90	53	)			
45	06	6		91	93	+/-	(1-I)		
46	54	$\div$		92	58	2nd rtn			
47	34	RCL							
48	04	4							

TABLE C-6B Given X and C_j , Find θ_0 and θ_j . Hirsch-Carter profile
Register Contents :

NATIONAL
42-394

0	1	2	3	4	5	6	7	8	9
	α	β	C_0	Z_i	θ_0	C_j			

Program

Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments
			LOAD REGISTERS	29	52	(	
00	33	STO	$\leftarrow X$	30	34	RCL	
01	01	1		31	02	2	
02	41	R/S	$\leftarrow \beta$	32	54	$\div$	
03	33	STO		33	02	2	
04	02	2		34	53	)	
05	41	R/S	$\leftarrow C_0$	35	94	=	$(\alpha Z_v)^{p/2}$
06	33	STO		36	12	INV	
07	03	3		37	23	sin	
08	41	R/S	$\leftarrow Z_j$	38	33	STO	
09	33	STO		39	05	5	
10	04	4		40	41	R/S	$\theta_0 \rightarrow$
11	41	R/S	$\leftarrow C_j$	41	34	RCL	
12	33	STO		42	05	5	
13	06	6		43	24	cos	
14	41	R/S		44	64	X	
			MAIN LOOP	45	34	RCL	
			$\leftarrow X$	46	06	6	
15	48	2nd $\sqrt{y}$		47	54	$\div$	
16	34	RCL		48	34	RCL	
17	02	2		49	03	3	
18	94	=		50	94	=	
19	20	2nd $1/x$		51	12	INV	
20	64	X		52	24	cos	
21	34	RCL		53	41	R/S	$\theta_j \rightarrow$
22	04	4					$\leftarrow X$
23	94	=	Z_v	54	22	GTO	LOOP
24	64	X		55	01	1	AGAIN
25	34	RCL		56	05	5	WITH
26	01	1					NEW X
27	94	=					
28	45	y^x					

which for a case 1 profile (see Section IV of the basic paper) fitted at the axis is the axial angle Θ_A . Range may be specified from an axis crossing or a vertex. Tables of the incomplete beta-function are used to convert $I_1 = I_\chi(\frac{1}{\rho}, \frac{1}{2})$ to χ . This program can be used to generate data for a range annotated ray angle diagram as described by Flatte⁴ and Cox⁵.

TABLE C-8A RANGE ANNOTATED RAY ANGLE DIAGRAM
COMPUTED BY HIRSCH-CARTER MODEL

42-384

* If I_1 comes out negative or G.T. 1 range is LARGER than that of reference level to vertex path.

OPERATING INSTRUCTIONS

STEP	PROCEDURE	ENTER	PRESS	Display
1	KEY IN PROGRAM		2nd CP ALL TABLE LRN LRN R/S	00 00 86 00
2	LOAD REGISTERS	α β C_0 B_1	R/S R/S R/S R/S	α β C_0 B_1
3	SELECT RAY BY REF. ANGLE	θ_0	R/S	Z_v
4	CONTINUE		R/S	
5	SUPPLY RANGE READ I_1	R	R/S	I_1 *
6	LOOK UP X IN TABLE OF I_x FUNCTION AND ENTER	X	R/S	Z
7	COMPUTE θ AT RANGE R AND DEPTH Z		R/S	θ
8	REPEAT STEPS 4-7 AS NECESSARY TO GET DATA FOR PLOT			
9	FOR NEW RAY AND REPEAT STEPS 4-7	θ_0	GTO 1 2 R/S	Z_v

TABLE C-8B RANGE ANNOTATED RAY ANGLE DIAGRAM

Register Contents

 NATIONAL
 42-394
 PRINTED IN U.S.A.

0	1	2	3	4	5	6	7	8	9
α	β	C_0	B_1	θ_0	Z_v	Z			
Program									
Loc.	Code	Key Entry	Comments	Loc.	Code	Key Entry	Comments		
			LOAD REGISTERS	44	94	=	$R \beta \tan \theta_0 / B_1 Z$		
00	33	STO	$\leftarrow \alpha$	45	46	NOP	for R		
01	01	1		46	46	NOP	measured		
02	41	R/S	$\leftarrow \beta$	47	46	NOP	from ref. level		
03	33	STO		48	46	NOP	or		
04	02	2		45	93	+/-	for R		
05	41	R/S	$\leftarrow C_0$	46	84	+	measured		
06	33	STO		47	01	1	from vertex		
07	03	3		48	94	=			
08	41	R/S	$\leftarrow B_1$						
09	33	STO		49	41	R/S	$I_1 \rightarrow$ FIND		
10	04	4					$\leftarrow \alpha$		
11	41	R/S	SELECT RAY	50	40	2nd $\sqrt{y}$	IN TABLE		
			BY REF. ANGLE	51	34	RCL			
			$\leftarrow \theta_0$	52	02	2			
12	33	STO		53	64	X	$X^{1/p}$		
13	05	5		54	34	RCL			
14	23	sin		55	06	6			
15	45	y^x		56	94	=			
16	52	(		57	33	STO			
17	02	2		58	07	7			
18	54	$\div$		59	41	R/S	$Z \rightarrow$		
19	34	RCL		60	34	RCL			
20	02	2		61	07	7			
21	53	)		62	64	X			
22	54	$\div$	$(\sin \theta_0)^{2/p}$	63	34	RCL			
23	34	RCL		64	01	1			
24	01	1		65	94	=	αZ		
25	94	=		66	45	y^x			
26	33	STO		67	34	RCL			
27	06	6		68	02	2			
28	41	R/S	$Z_v \rightarrow$	69	74	-			
			MAIN LOOP	70	01	1			
			STARTS HERE	71	94	=			
29	34	RCL		72	93	+/-	$(1 - (\alpha Z)^\beta)$		
30	06	6		73	20	2nd $1/x$			
31	20	2nd $1/x$		74	48	2nd $\sqrt{x}$			
32	54	$\div$		75	64	X			
33	34	RCL		76	34	RCL			
34	04	4		77	05	5			
35	64	X	$1/B_1 Z_v$	78	24	cos			
				79	94	=	$\cos \theta$		
36	41	R/S	$\leftarrow R$	80	12	INV			
37	64	X		81	24	cos			
38	34	RCL		82	41	R/S	$\theta \rightarrow$		
39	02	2		83	22	GTO	RETURN		
40	64	X	$R \beta / B_1 Z_v$	84	02	2	TO START		
41	34	RCL		85	09	9	OF MAIN		
42	05	5					LOOP		
43	25	tan							

REFERENCES FOR SUPPLEMENT

1. E. S. Pearson and N. L. Johnson, "Tables of the Incomplete Beta-function 2nd Edition". Cambridge University Press (1968).
2. M. Abramowitz and I. A. Stegun, "Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables". U. S. Department of Commerce, National Bureau of Standards Applied Mathematics Series 55. Ninth printing. U. S. Government Printing Office, Washington, D.C. (1970).
3. D. Michael Milder, "Ray and Wave Invariants for SOFAR Channel Propagation". J.Acoust.Soc.Am. 46, pp 1259-1263. (1969).
4. Stanley M. Flatte, "Angle-depth diagram for use in underwater acoustics". J.Acoust.Soc.Am. 60, pp 1020-1023. (1976).
5. Henry Cox, "Approximate ray angle diagram". J.Acoust.Soc.Am. 61, pp 353-359. (1977).

DISTRIBUTION LIST

Director
Strategic Systems Project Office (PM-1)
Department of the Navy
Washington, DC 20360

OASN (RE&S)
Room 4D, 745 Pentagon
Washington, DC 20301
Attn: G. A. Cann

Assistant Director
Ocean Control DDR&E
Room 3D, 1048 Pentagon
Washington, DC 20301

Chief of Naval Operations
Department of the Navy
Washington, DC 20350
Attn: Code OP 02
OP 03
OP 023
OP 032
OP 092U
OP 095
OP 095E
OP 098
OP 951
OP 952
OP 955
OP 955F
OP 966
OP 967
OP 981

Chief of Naval Material
Department of the Navy
Washington, DC 20360
Attn: Code NMAT 08T
NMAT 98T1
NMAT 08T2
NMAT 08T24
NMAT 08T4

Commander
Pacific Missile Test Center
Point Mugu, CA 93047
Attn: Library

Commanding Officer
Fleet Numerical Weather Central
Monterey, CA 93940

President
Naval War College
Newport, RI 02840
Attn: Library

Commanding Officer
Fleet ASW Training Center, Pacific
San Diego, CA 92147

Commanding Officer
Fleet ASW Training Center, Atlantic
Naval Station
Norfolk, VA 23511

Commanding Officer and Director
Defense Documentation Center
Cameron Station, Bldg. 5
5010 Duke St.
Alexandria, VA 22314

Commander
Naval Oceanography Command
NSTL Station, MS 39529

Anti-Submarine Warfare Systems
Project Office
Department of the Navy
Washington, DC 20360
Attn: Code PM-4 (A. Bernard)

Naval Ocean Systems Center
San Diego, CA 92152

U. S. Coast Guard HQ
400 Seventh St., SW
Washington, DC 20591
Attn: R & D Director

National Oceanic & Atmos. Admin.
6001 Executive Blvd.
Rockville, MD 20852

Applied Physics Laboratory
University of Washington
1013 NE Fortieth St.
Seattle, WA 98195
Attn: Library

Applied Research Laboratories
University of Texas
P. O. Box 8029
Austin, TX 78712
Attn: L. D. Hampton
G. E. Ellis
S. K. Mitchell
K. C. Focke
K. E. Hawker
S. R. Rutherford
T. G. Muir
D. J. Shirley
Library

Applied Research Laboratory
Pennsylvania State Univ.
P. O. Box 30
State College, PA 16801
Attn: Library

Applied Physics Laboratory
Johns Hopkins University
Laural, MD 20810
Attn: G. L. Smith
J. R. Austin
Library

Marine Physical Laboratory
University of California, San Diego
San Diego, CA 92152
Attn: F. N. Spiess
V. C. Anderson
F. H. Fisher
Library

Lamont-Doherty Geol. Obs.
Palisades, NY 10964
Attn: G. M. Bryan
W. J. Ludwig
B. Tucholke
R. E. Houtz
J. E. Nafe
W. B. Kutschera

Woods Hole Oceanographic Inst.
Woods Hole, MA 02543
Attn: J. Ewing
E. E. Hayes

Library

TRACOR, Inc.
6500 Tracor Lane
Austin, TX 78721
Attn: A. F. Wittenborn
Library

Mechanics Research, Inc.
7929 Westpark Dr.
McLean, VA 22101
Attn: N. Peck

Western Electric Co.
Gelford Center
P. O. Box 20046
Greensboro, NC 27420
Attn: R. Scudder

Applied Hydroacoustics, Inc.
Montgomery Plaza Bldg.
656 Quince Orchard Rd.
Gaithersburg, MD 20760
Attn: F. Ryder

TRW
7600 Colshire Dr.
McLean, VA 22101
Attn: R. Murawski
I. B. Gereben

TRACOR Inc.
1601 Research Blvd.
Rockville, MD 20850
Attn: R. J. Urlick

Planning Systems Inc.
7900 Westpark Dr.
McLean, VA 22101
Attn: L. Solomon

Arthur D. Little, Inc.
15 Acorn Park
Cambridge, MA 02140
Attn: W. G. Sykes
G. Raisbeck

Commander
Naval Electronic Systems Command
Department of the Navy
Washington, DC 20360
Attn: Code 035
320
320 (J. Cybulski)
320 (J. A. Sinsky)
PME 124
124-20
124-30
124-34 (J. Reeves)
124-40
124-60
124TA

Commander
Naval Sea Systems Command
Department of the Navy
Washington, DC 20362
Attn: Code 06H1
06H1-4

Commander
Naval Air Systems Command
Department of the Navy
Washington, DC 20360
Attn: Code 350
360
370
604
PMA 264

Office of Naval Research
Department of the Navy
Arlington, VA 22217
Attn: Code 102-OS
480
483
485
486

U. S. Naval Oceanographic Office
NSTL Station, MS 39522
Attn: Code 3400
3721 (R. Anderson)

Commander
New London Laboratory
Naval Underwater Systems Center
New London, CT 06320
Attn: P. Herstein
F. R. DiNapoli
S. R. Santaniello
R. L. Deavenport
R. W. Hasse
J. J. Hanrahan
R. K. Dullea
B. F. Cole
T. G. Bell

Commanding Officer
Newport Laboratory
Naval Underwater Systems Center
Newport, RI 02840
Attn: Library

Commander
Naval Air Development Center
Warminster, PA 18974
Attn: J. Howard
C. L. Bartberger
P. S. Haas
J. H. Rubisch
Library

Director
Naval Research Laboratory
Department of the Navy
Washington, DC 20375
Attn: Code 2627
8000
8100
8106 (R. K. Perry)
8106 (H. S. Fleming)
8108
8120 (R. L. Dicus)
8120 (R. H. Ferris)
8160 (B. B. Adams)
8403 (R. T. Swim)
Library
B. G. Hurdle
O. I. Diachok

Chief Scientist
Naval Underwater Sound Reference Division
Naval Research Laboratory
P. O. Box 8337
Orlando, FL 32806
Attn: Library

Commanding Officer
Naval Coastal Systems Laboratory
Panama City, FL 32401
Attn: E. G. Mcleroy, Jr.
W. Tolbert
Library

Commander
David W. Taylor Naval Ship
Research and Development Center
Bethesda, MD 20084
Attn: Library

Officer in Charge
Naval Ship R & D Center
Annapolis, MD 21402
Attn: Library

Superintendent
Naval Postgraduate School
Monterey, CA 93940
Attn: H. Medwin
O. B. Wilson, Jr.
R. S. Andrews
Library

Commander
Naval Surface Weapons Center
White Oak Laboratory
Silver Spring, MD 20910

Commanding Officer
Civil Engineering Laboratory
Naval Construction Bn. Center
Port Hueneme, CA 93043
Attn: Library

Defense Advanced Research Projects Agency
1400 Wilson Blvd.
Arlington, VA 22209
Attn: Dr. Gustafson

ARPA Research Center
Unit 1, Bldg. 301A
NAS Moffet Field, CA 94035
Attn: E. L. Smith

Science Applications, Inc.
8400 Westpark Dr.
McLean, VA 22101
Attn: J. S. Hanna
C. W. Spofford

Bell Telephone Laboratories
Whippany, NJ 07981
Attn: R. Lauver
Mr. Pennotti

Commander-in-Chief, Atlantic Fleet
Naval Base
Norfolk, VA 23511

Commander-in-Chief, U. S. Navy Europe
FPO
San Francisco, CA 96610

Commander-in-Chief, Pacific Fleet
Box 11, FPO
San Francisco, CA 96610
Attn: Code 352

Commander
Operational Test & Eval. Force
Naval Base
Norfolk, VA 23511

Commander
Submarine Development Group 1
FPO
San Diego, CA 92132

Commander
Submarine Development Group 2
Naval Submarine Base-New London
Groton, CT 06340

Office of Naval Research
Department of the Navy
NORDA Liaison Office
Arlington, VA 22217
Attn: K. W. Lackie

Office of Naval Research
Resident Representative
University of California
La Jolla, CA 92093

Commander
Naval Intelligence Support Center
4301 Suitland Road
Washington, DC 20390
Attn: Code 222

Director of Naval Warfare Analysis
Institute of Naval Studies
1401 Wilson Blvd.
Arlington, VA 22209

Director for Naval Matters
Center for Naval Analysis
Arlington, VA 22209
Attn: C. E. Woods

Commander
Naval Oceanographic Office
Washington, DC 20374
Attn: Code 3410 (J. L. Carrol)

Commanding Officer
Naval Ocean Research and Development Activity
NSTL Station, MS 39529
Attn: Code 110

200 (K. W. Mackensie)
200 (CDR T. McClosky)
230 (H. E. Morris)
300 (H. Eppert)
320
330
340
341 (D. F. Fenner)
350
360
360 (G. L. Maynard)
360 (T. L. Holcomb)
361 (A. Lowrie)
362 (J. E. Matthews)

Code 410
420
440
460
500
500 (E. M. Stanley)
500 (M. G. Lewis)
600
Library

ADDITIONAL DISTRIBUTION LIST-----

Dr. D. C. Stickler
Courant Inst. Math. Sciences
New York, NY 10012

Commander, Third Fleet
FPO
San Francisco, CA 96610
Attn: N-32

Dr. G. H. Sutton
Hawaii Inst. of Geophysics
The University of Hawaii
2525 Correa Rd.
Honolulu, HI 96822

Commander, Seventh Fleet
FPO
San Francisco, CA 96610
Attn: N-34

Dr. Morris Schulkin
9325 Orchard Brook Dr.
Potomac, MD 20854

Commander, Submarine Force
U. S. Pacific Fleet
FPO
San Francisco, CA 96601

Dr. P. A. Rona
National Oceanic and Atmos. Admin.
15 Rickenbacker Causeway
Miami, FL 33149

Marvin S. Weinstein
Underwater Systems, Inc.
3121 Georgia Ave.
Silver Spring, MD 20910

Director
Geophysics Laboratory
University of Texas
700 The Strand
Galveston, TX 77550

Dr. C. S. Clay
Dept. of Geology and Geophysics
University of Wisconsin
Madison, WI 53700

Dr. A. O. Williams, Jr.
Dept. of Physics
Brown University
Providence, RI 02912

Dr. H. M. Uberall
The Catholic University of America
6220 Michigan Ave., NE
Washington, DC 20017

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER WHOI-78-65	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) RAY CALCULATIONS OF OCEAN SOUND CHANNELS USING A POCKET PROGRAMMABLE CALCULATOR AND EXTENDED FORMS OF THE HIRSCH-CARTER MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE BETA FUNCTION		5. TYPE OF REPORT & PERIOD COVERED Technical
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Lincoln Baxter, II		8. CONTRACT OR GRANT NUMBER(s) N00014-77-C-0196
9. PERFORMING ORGANIZATION NAME AND ADDRESS Woods Hole Oceanographic Institution Woods Hole, MA 02543		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS NORDA National Space Technology Laboratory Bay St. Louis, MS 39529		12. REPORT DATE October 1978
		13. NUMBER OF PAGES 36 & 34-S
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release, distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) 1. Acoustic Ray Computations 2. Ocean Sound Channels 3. Programmable Pocket Calculators		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) Formulas for curve fitting and ray computation using compound models made up of several different layers of form $C^2 = C_0^2 (1 - aZ ^P)^{-1}$ are presented. Examples of computation by pocket programmable calculator on two Sargasso Sea profiles, one from the center of a cold ring eddy are given. Necessary tables of the incomplete beta-function and calculator programs are included in a supplement.		

Woods Hole Oceanographic Institution WHOI-78-65	RAY CALCULATIONS OF OCEAN SOUND CHANNELS USING A POCKET PROGRAMMABLE CALCULATOR AND EXTENDED FORMS OF THE HIRSCH-CARTER MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE BETA FUNCTION by Lincoln Baxter II. Prepared for the Office of Naval Research under Contract N00014-77-C-0196. (70 pages)	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED	Woods Hole Oceanographic Institution WHOI-78-65 RAY CALCULATIONS OF OCEAN SOUND CHANNELS USING A POCKET PROGRAMMABLE CALCULATOR AND EXTENDED FORMS OF THE HIRSCH-CARTER MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE BETA FUNCTION by Lincoln Baxter II. Prepared for the Office of Naval Research under Contract N00014-77-C-0196. (70 pages)	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED
Woods Hole Oceanographic Institution WHOI-78-65	RAY CALCULATIONS OF OCEAN SOUND CHANNELS USING A POCKET PROGRAMMABLE CALCULATOR AND EXTENDED FORMS OF THE HIRSCH-CARTER MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE BETA FUNCTION by Lincoln Baxter II. Prepared for the Office of Naval Research under Contract N00014-77-C-0196. (70 pages)	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED	Woods Hole Oceanographic Institution WHOI-78-65 RAY CALCULATIONS OF OCEAN SOUND CHANNELS USING A POCKET PROGRAMMABLE CALCULATOR AND EXTENDED FORMS OF THE HIRSCH-CARTER MATHEMATICAL MODEL WITH TABLES OF THE INCOMPLETE BETA FUNCTION by Lincoln Baxter II. Prepared for the Office of Naval Research under Contract N00014-77-C-0196. (70 pages)	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED	1. Acoustic Ray Computations Sound Channels 2. 3. Programmable Pocket Calculators I. Baxter, Lincoln, II II. N00014-77-C-0196 This card is UNCLASSIFIED