

RESEARCH ARTICLE

Searching for the Oldest Baobab of Madagascar: Radiocarbon Investigation of Large *Adansonia rubrostipa* Trees

Adrian Patrut^{1*}, Karl F. von Reden², Pascal Danthu^{3,4}, Jean-Michel Leong Pock-Tsy⁴, Roxana T. Patrut⁵, Daniel A. Lowy⁶

1 Babeş-Bolyai University, Faculty of Chemistry, Cluj-Napoca, Romania, **2** NOSAMS Facility, Department of Geology & Geophysics, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, United States of America, **3** Cirad, UPR BSEF, Montpellier, France, **4** DP Forêts et Biodiversité, Antananarivo, Madagascar, **5** Babeş-Bolyai University, Faculty of Biology and Geology, Cluj-Napoca, Romania, **6** Northern Virginia Community College (Nova University), Alexandria Campus, Alexandria, Virginia, United States of America

* apatrut@gmail.com



OPEN ACCESS

Citation: Patrut A, von Reden KF, Danthu P, Leong Pock-Tsy J-M, Patrut RT, Lowy DA (2015) Searching for the Oldest Baobab of Madagascar: Radiocarbon Investigation of Large *Adansonia rubrostipa* Trees. PLoS ONE 10(3): e0121170. doi:10.1371/journal.pone.0121170

Academic Editor: John P. Hart, New York State Museum, UNITED STATES

Received: October 12, 2014

Accepted: January 30, 2015

Published: March 25, 2015

Copyright: © 2015 Patrut et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper.

Funding: The research was fully funded by the Romanian Ministry of National Education CNCS-UEFISCDI under grant PN-II-ID-PCE-2013-76 (URL: <http://uefiscdi.gov.ro/>). The funder had no role in study design, data collection and analysis, decision to publish or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

We extended our research on the architecture, growth and age of trees belonging to the genus *Adansonia*, by starting to investigate large individuals of the most widespread Malagasy species. Our research also intends to identify the oldest baobabs of Madagascar. Here we present results of the radiocarbon investigation of the two most representative *Adansonia rubrostipa* (fony baobab) specimens, which are located in south-western Madagascar, in the Tsimanampetsotse National Park. We found that the fony baobab called “Grandmother” consists of 3 perfectly fused stems of different ages. The radiocarbon date of the oldest sample was found to be 1136 ± 16 BP. We estimated that the oldest part of this tree, which is mainly hollow, has an age close to 1,600 yr. This value is comparable to the age of the oldest *Adansonia digitata* (African baobab) specimens. By its age, the Grandmother is a major candidate for the oldest baobab of Madagascar. The second investigated specimen, called the “polygamous baobab”, consists of 6 partially fused stems of different ages. According to dating results, this fony baobab is 1,000 yr old. This research is the first investigation of the structure and age of Malagasy baobabs.

Introduction

The genus *Adansonia* belonging to the Bombacoideae, a subfamily of Malvaceae, consists of nine species out of which six are endemic to Madagascar [1–3].

Up to present, the research dedicated to the architecture, growth and age of *Adansonia* trees was performed exclusively on the African baobab (*Adansonia digitata* L.). We found that all large African baobabs are multi-stemmed and exhibit preferentially ring-shaped structures; the oldest dated specimens were found to be between 1,000 and 2,000 years old [4–10]. We extended our research on the *Adansonia* genus by starting to investigate large individuals of the most

representative Malagasy species, *i.e.*, *Adansonia rubrostipa* Jum. & H. Perrier (fony baobab), *Adansonia za* Baill. (za baobab) and *Adansonia grandidieri* Baill. (Grandidier's baobab or reniala), which grow in west and south; each of these species is represented by over 1 million individuals [2,11,12]. The research is based on our approach which consists of AMS (accelerator mass spectrometry) radiocarbon dating of small wood samples collected from different areas of the trunk/stems of large baobabs [6,8]. Our research also intends to identify the oldest baobabs of Madagascar and to answer two main questions: (i) are there Malagasy baobabs over 1,000 years old? (ii) if yes, which species do these individuals belong to?

A. rubrostipa is represented by small to medium and also by some relatively large, rather bizarre-looking trees. The fony baobabs have bottle-shaped or quasi-cylindrical trunks and short horizontal branches that turn upwards at the ends [2]. According to recent estimates, there are up to 1.5 million *A. rubrostipa* individuals which are widespread along the west coast of Madagascar. The largest known and probably the oldest fony baobabs are both located in the Tsimanampetsotse National Park, Toliara Province (Fig. 1). Here we present the radiocarbon investigation results of these two most representative *A. rubrostipa* individuals.

Materials and Methods

Ethics statement

The investigation and sampling of the baobabs was approved and authorised by the Forestry Direction of the Ministry of Environment, Ecology and Forestry of Madagascar and by the Madagascar National Parks. The Madagascar National Parks, Tsimanampetsotse National Park and PARRUR project (ECOBABO) provided scientific assistance and support in the investigation. The baobabs were not endangered by the sampling. After each coring, the increment borer was disinfected with methyl alcohol. The small coring holes were sealed with Steriseal (Efekto), a special polymer sealing product, for preventing any possible infection of the trees.

The two baobabs and their area

The two fony baobabs are located in the Tsimanampetsotse National Park, at a distance of only 1.5 km apart from each other. The mean annual rainfall in this very arid area is around 350 mm.

The "Grandmother" (La grand-mère) is generally considered the oldest *A. rubrostipa*. Its GPS coordinates are 24°02.707' S, 043°45.266' E and the altitude is 36 m. It has a height of 7.47 m and a circumference at breast height (cbh) of 9.67 m. The overall wood volume is 25 m³. The tree, which has a pot shape, consists of 3 perfectly fused stems (Figs. 2–4). However, some fusion lines of stems can be identified at the upper part of the trunk.

The "polygamous baobab" (Le baobab polygame), the largest known *A. rubrostipa*, is located at 24°03.060' S, 043°45.418' E and the altitude is 27 m. It has a maximum height of 14.2 m and a restored circumference (cbh) of 13.50 m. The overall wood volume is 60 m³. It exhibits a cluster structure with 6 partially fused stems. The stem from the northern extremity collapsed some time ago, but it is still alive (Figs. 5 and 6).

Sample collection

Several wood samples were collected from the outer part of the 3 stems of Grandmother, at convenient heights between 1.27–1.55 m. The longest samples from each stem (labelled GM-1, GM-2, GM-3) were investigated. The stem numbering (I, II, III) and the sampling positions of the longest samples are shown in Fig. 4.




Fig 1. Map of Madagascar, showing the Toliara province (in yellow) and the position of the Tsimanampetsotse National Park (marked by the red arrow).

doi:10.1371/journal.pone.0121170.g001

Additional 5 wood samples (labelled PB-1 to PB-5) were collected from 5 stems of the polygamous baobab, at heights between 1.30–1.53 m. The stem numbering (from I to VI) is displayed in Figs. 5 and 6.

The samples were collected by using Haglöf increment borers (0.80–1.00 m long, 0.010–0.012 m inner diameter). Several small pieces/segments of the length of 0.001 m were extracted from determined positions of the 10 samples and investigated by AMS radiocarbon dating.

Sample preparation

The standard acid-base-acid pretreatment method [13] was used for removing soluble and mobile organic components. The resulting samples were combusted to CO₂ *via* the closed tube



Fig 2. General view of the old Grandmother.

doi:10.1371/journal.pone.0121170.g002

combustion method [14]. Next, CO₂ was reduced to graphite on iron catalyst, under hydrogen atmosphere [15]. The graphite samples were analysed by AMS.

AMS measurements

AMS radiocarbon measurements were performed at the NOSAMS Facility of the Woods Hole Oceanographic Institution by using the Pelletron Tandem 500 kV AMS system [16,17]. The obtained fraction modern values were ultimately converted to a radiocarbon date.

Calibration

Fraction modern values were calibrated and converted into calendar ages with the OxCal v4.2 for Windows [18], by using the SHCal13 atmospheric data set [19].

Results

AMS results and calibrated ages

Fraction modern values and radiocarbon dates *BP* (before present, referring to AD 1950) of 10 wood samples/segments which originate from the Grandmother (GM) and the polygamous baobab (PB) are displayed in Tables 1 and 2. Radiocarbon dates and errors were rounded to



Fig 3. The strange looking tri-stemmed trunk of the Grandmother showing its various sized bulbous formations.

doi:10.1371/journal.pone.0121170.g003

the nearest year. The accession numbers in the NOSAMS database of the 10 radiocarbon analyses are also listed in Tables 1 and 2.

The 1σ probability distribution (68.2%) was typically selected to derive calibrated age ranges. For 4 sample segments, the 1σ distribution is consistent with one range of calendar years, while for other 4 segments, the 1σ distribution corresponds to two or three ranges of calendar years. For these segments, the confidence interval of one range is considerably greater than that of the others; therefore, it was selected as the cal AD range of the sample for the purpose of this discussion. For 2 sample segments, there are two 1σ ranges with close confidence intervals. In these special cases *i.e.*, GM-1(a) and GM-1(c), we used the higher 2σ probability distribution (95.4%) for calibration.

For obtaining single age values, we derived a mean calibrated (cal) age of each segment from the selected range (marked in bold). Calibrated ages of samples/segments represent the difference between AD 2014 and the mean AD value of the selected range, with the corresponding error. All calibrated sample/segment ages, as well as stem ages and tree ages are expressed in (calendar) yr in AD 2014. Calibrated ages and errors were rounded to the nearest 5 yr.




Fig 4. Cross-section of Grandmother (at 1.30 m above ground) showing the 3 fused stems (I, II and III) and the projection of sampling points' positions (1, 2 and 3).

doi:10.1371/journal.pone.0121170.g004

Results of samples collected from the Grandmother

Even if the borer penetration in the soft wood was quasi-complete in all cases, the collected samples were much shorter. This shows that more than half of the tree is hollow, due to decay. We extracted and dated 3 segments (labelled a, b, c) from sample GM-1, which originate from the largest stem I. The innermost and oldest segment GM-1(c) corresponds to the sample end located at 0.46 m from the sampling point. Its radiocarbon date was found to be 1136 ± 16 BP, which corresponds to a calibrated age of 1070 ± 50 yr. The second oldest dated segment GM-2 (b), which corresponds to a depth of 0.20 m in the wood, has an age of 650 yr (neglecting the error). The ages and distances of segments GM-1(b) and GM-1(c) indicate that stem I grew 0.26 m in 420 yr in the sampling direction. This value indicates a mean radial increase of $0.62 \times 10^{-3} \text{ m yr}^{-1}$ over the time frame AD 945–1365.

The outermost segment GM-1(a), which was adjacent to the bark, is not as young as it would be expected. Its calibrated age of 250 yr shows that the large stem I stopped growing around AD 1765. One should, however, mention that the calibration of such low radiocarbon dates is very difficult and also somewhat uncertain, particularly with the southern data sets. According to ages and distances of segments GM-1(a) and GM-1(b), stem I grew 0.20 m in 400 yr in the sampling direction. The value indicates a slightly lower mean radial increase, of $0.50 \times 10^{-3} \text{ m yr}^{-1}$ over the time frame AD 1365–1765.

We dated the deepest segments from samples GM-2 and GM-3, which were collected from the second stem II and from the smallest stem III. These segments correspond to a depth in the wood of 0.52 and 0.55 m and have ages of 735 ± 10 and 535 ± 20 yr.



Fig 5. Western view of the polygamous baobab with stem numbering (from I to VI).

doi:10.1371/journal.pone.0121170.g005

Results of samples collected from the polygamous baobab

The 5 samples PB-1, PB-2, PB-3, PB-4 and PB-5 originate from the five standing stems, *i.e.*, I, II, III, IV and V. The 5 samples were also shorter than the penetration distance of the increment borer, showing the presence of significant hollow parts in each stem. In all cases, we extracted and dated the innermost/ deepest segment of each sample.

The oldest dated segment PB-1 originates from stem I, at a distance of 0.52 m from the sampling point. Its radiocarbon date of 852 ± 28 BP corresponds to a calibrated age of 785 ± 25 yr. The other 4 sample segments exhibit a continuous age decrease from PB-2 to PB-5 having ages between 565 and 375 yr.

Discussion

Ages of Grandmother and its stems

For determining the true age of Grandmother, which is identical to the age of stem I, we used the age of the oldest segment GM-1(a), the estimated distance to the calculated pith of stem I



Fig 6. Eastern view of the polygamous baobab with stem numbering.

doi:10.1371/journal.pone.0121170.g006

from the sampling point (1.15 m) and from the oldest segment (0.69 m) and also the mean growth rates for 650 and 420 yr (see [Table 1](#) and [Fig. 4](#)). In a conservative estimate, the age of stem I and, consequently, of Grandmother is 1,500–1,700 yr, *i.e.*, $1,600 \pm 100$ yr. According to this value, the Grandmother started growing around AD 400.

The ages of stems II and III were determined from the ages of samples GM-2 and GM-3 and the distances from these sample ends to the presumptive piths of stems II and III. According to these values, the estimated ages of stems II and III are of ca. 1,100 and 750 yr.

Ages of polygamous baobab and its stems

The age of the polygamous baobab, which is identical to the age of its oldest stem I, was determined by extrapolating the calibrated age of sample PB-1 to the corresponding pith (see [Table 2](#)). Therefore, we consider that the polygamous baobab and its stem I are $1,000 \pm 100$ yr old. The ages of the other 4 sampled stems were estimated in a similar manner. Thus, stem II and stem III, which are fused completely up to a height of 2.60 m, are both around 750 yr old.

Table 1. AMS Radiocarbon dating results and calibrated calendar ages of samples collected from the Grandmother.

Sample (Segment)	Depth in the wood ¹ [Distance to stem pith ¹] (10 ⁻² m)	Fraction modern [error]	Radiocarbon date [error] (¹⁴ C yr BP)	Cal AD range (1σ or 2σ) ² [confidence interval]	Sample age [error] (yr in AD 2014)	NOSAMS Accession #
GM-1(a)	0.5 [115]	0.9765 [±0.0026]	191 [±21]	1666–1712 [25.8%] 1718–1814 [49.7%] 1836–1884 [12.7%] 1924–... [7.3%]	250 [±50]	OS-110648
GM-1(b)	20 [115]	0.9189 [±0.0032]	679 [±26]	1300–1320 [25.6%] 1344–1388 [42.6%]	650 [±20]	OS-108121
GM-1(c)	46 [115]	0.8681 [±0.0020]	1136 [±16]	892–995 [93.7%] 1005–1013 [1.7%]	1070 [±50]	OS-112137
GM-2	51 [80]	0.9074 [±0.0020]	781 [±16]	1268–1288 [68.2%]	735 [±10]	OS-111763
GM-3	47 [65]	0.9492 [±0.0026]	419 [±21]	1457–1498 [58.2%] 1599–1608 [10.0%]	535 [±20]	OS-110647

¹from the sampling point

²with the SHCal13 calibration

doi:10.1371/journal.pone.0121170.t001

Stems IV and V have estimated ages of 600 and 500 yr. The fallen stem VI, which broke partially in the collapse, was not sampled. According to its size, we estimate that it is around 300 yr old.

Architecture and growth of the two fony baobabs

The two large and old *A. robustipa* specimens started growing as single-stemmed and became over time multi-stemmed, thus developing different shapes. This evolution is due to the general ability of baobabs, which we identified for the first time for *A. digitata* [5,9,10], to produce stems during their life cycle, such as other trees produce branches.

The Grandmother started growing its main stem I around AD 400. The tree produced new stems in AD 900 and 1250, developing over time its well-known pot shape with 3 perfectly fused stems. The main stem I stopped growing in AD 1750, because of old age.

Table 2. AMS Radiocarbon dating results and calibrated calendar ages of samples collected from the polygamous baobab.

Sample (Segment)	Depth in the wood ¹ [Distance to stem pith ¹] (10 ⁻² m)	Fraction modern [error]	Radiocarbon date [error] (¹⁴ C yr BP)	Cal AD range (1σ) ² [confidence interval]	Sample age [error] (yr in AD 2014)	NOSAMS Accession #
PB-1	55 [82]	0.8994 [±0.0034]	852 [±28]	1214–1266 [68.2%]	785 [±25]	OS-110646
PB-2	47 [75]	0.9420 [±0.0025]	480 [±20]	1440–1456 [68.2%]	565 [±10]	OS-108826
PB-3	39 [60]	0.9463 [±0.0029]	443 [±23]	1448–1487 [68.2%]	545 [±20]	OS-110644
PB-4	41 [65]	0.9602 [±0.0030]	326 [±24]	1512–1549 [34.9%] 1560–1572 [8.3%] 1622–1646 [25.1%]	485 [±20]	OS-108824
PB-5	40 [65]	0.9621 [±0.0029]	310 [±23]	1514–1543 [25.3%] 1624–1654 [42.9%]	375 [±15]	OS-109280

¹from the sampling point

²with the SHCal13 calibration

doi:10.1371/journal.pone.0121170.t002

The main stem I of the polygamous baobab started growing around AD 1000. Stems II and III emerged probably together in AD 1250 and the fony baobab gained a triangular structure. Next, 3 new stems emerged successively toward north in AD 1400, 1500 and 1700. Thus, the polygamous baobab developed its cluster structure, with 6 very distinct bottle-shaped stems, which are fused only in the proximity of their base and have separate canopies.

Conclusions

We investigated by AMS radiocarbon dating two representative *Adansonia rubrostipa* specimens, which are located in a very arid area of the Tsimanampetsotse National Park, in southwestern Madagascar. According to our research, the fony baobab called Grandmother, which has a typical pot shape, consists of 3 fused stems of different ages. The radiocarbon date of the oldest collected wood sample was of 1136 ± 16 BP, which corresponds to a calibrated age of 1070 ± 50 yr. We determined that the oldest stem of this baobab is $1,600 \pm 100$ yr old. This value is close to the age of the oldest dated *Adansonia digitata* specimens. The other two stems of Grandmother exhibit ages of 1,100 and 750 yr. The second investigated specimen, the polygamous baobab, is composed of 6 partially fused stems. Its 6 stems have ages between 300 and 1,000 yr.

According to accurate dating results, the fony baobab becomes the second *Adansonia* species with individuals that can live over 1,000 yr. On the other hand, by its age, the Grandmother is a major candidate for being the oldest baobab of Madagascar.

This research is the first investigation of the architecture, growth and age of Malagasy baobabs.

Acknowledgments

The authors thank Laszlo Rakosy, Emilson Rakotoarisoa, Wilfried Ramahafaly and Lucien Rasoanaivoson for helpful assistance and discussions.

Author Contributions

Conceived and designed the experiments: AP. Performed the experiments: AP KFVR PD JMLPT RTP. Analyzed the data: AP KFVR PD JMLPT RTP DAL. Contributed reagents/materials/analysis tools: AP KFVR PD JMLPT RTP. Wrote the paper: AP.

References

1. Wickens GE, Lowe P. The Baobabs: Pachycauls of Africa, Madagascar and Australia. Dordrecht: Springer; 2008.
2. Baum DA. A systematic revision of *Adansonia* (Bombacaceae). *Ann Mo Bot Gard*. 1995; 82: 440–471.
3. Pettigrew JD, Bell LK, Bhagwandin A, Grinan E, Jillani N, Meyer J, et al. Morphology, ploidy and molecular phylogenetics reveal a new diploid species from Africa in the baobab genus *Adansonia* (Malvaceae; Bombacoideae). *Taxon*. 2013; 61: 1240–1250.
4. Patrut A, von Reden KF, Lowy DA, Alberts AH, Pohlman JW, Wittmann R, et al. Radiocarbon dating of a very large African baobab. *Tree Physiol*. 2007; 27: 1569–1574. PMID: [17669746](#)
5. Patrut A, von Reden KF, Lowy DA, Mayne DH, Elder KE, Roberts ML, et al. Comparative AMS radiocarbon dating of pretreated versus non-pretreated wood samples. *Nucl Instr and Meth B*. 2010; 268: 910–913.
6. Patrut A, Mayne DH, von Reden KF, Lowy DA, Van Pelt R, McNichol AP, et al. Fire History of a Giant African Baobab evinced by Radiocarbon Dating. *Radiocarbon*. 2010; 52(2–3): 717–726. PMID: [22228915](#)
7. Patrut A, Mayne DH, von Reden KF, Lowy DA, Venter S, McNichol AP, et al. Age and Growth Rate Dynamics of an Old African Baobab determined by Radiocarbon Dating. *Radiocarbon*. 2010; 52(2–3): 727–734. PMID: [22228915](#)

8. Patrut A, von Reden KF, Van Pelt R, Mayne DH, Lowy DA, Margineanu D. Age determination of large live trees with inner cavities: Radiocarbon dating of Platland tree, a large African baobab. *Ann Forest Sci.* 2011; 68: 993–1003.
9. Patrut A, von Reden KF, Mayne DH, Lowy DA, Patrut RT. AMS radiocarbon investigation of the African baobab: Searching for the oldest tree. *Nucl Instr Meth B.* 2013; 294: 622–626.
10. Patrut A, Woodborne S, von Reden KF, Hall G, Hofmeyr M, Lowy DA, et al. African Baobabs with False Inner Cavities: The Radiocarbon Investigation of the Lebombo Eco Trail Baobab. *PLoS ONE.* 2015 Jan 26. 10(1): e0117193. doi: [10.1371/journal.pone.0117193](https://doi.org/10.1371/journal.pone.0117193) PMID: [25621989](https://pubmed.ncbi.nlm.nih.gov/25621989/)
11. Vielledent G, Cornu C, Cuni Sanchez A, Leong Pock-Tsy J- M, Danthu P. Vulnerability of baobab species to climate change and effectiveness of the protected area network in Madagascar: Towards new conservation priorities. *Biol Conserv.* 2013; 166: 11–22.
12. Pakenham T. *The Remarkable Baobab.* London: Weidenfield & Nicholson; 2004.
13. Olsson IU. Radiometric Methods. In: Berglung B, editor. *Handbook of Holocene palaeoecology and palaeohydrology.* Chichester: Wiley; 1986. pp. 273–312. PMID: [17794573](https://pubmed.ncbi.nlm.nih.gov/17794573/)
14. Sofer Z. Preparation of carbon dioxide for stable carbon isotope analysis of petroleum fractions. *Anal Chem.* 1980; 52: 1389–1391.
15. Vogel JS, Southon JR, Nelson DE, Brown TA. Performance of catalytically condensed carbon for use in accelerator mass-spectrometry. *Nucl Instr Meth B.* 1984; 5: 289–293.
16. Povinec PP, Litherland AE, von Reden KF. Developments in Radiocarbon Technologies: From the Libby Counter to Compound-Specific AMS Analyses. *Radiocarbon.* 2009; 51: 45–78.
17. Roberts ML, Burton JR, Elder KL, Longworth BE, McIntyre CP, von Reden KF, et al. A high-performance ¹⁴C Accelerator Mass Spectrometry system. *Radiocarbon.* 2010; 52: 228–235.
18. Bronk Ramsey C. Bayesian analysis of radiocarbon dates. *Radiocarbon.* 2009; 51: 337–360.
19. Hogg AG, Hua Q, Blackwell PG, Niu M, Buck CE, Guilderson TP, et al. SHCal13 Southern Hemisphere calibration, 0–50,000 years cal BP. *Radiocarbon.* 2013; 55: 1889–1903.