WHOI-87-12

**Options for Radioactive and Other Hazardous Waste Siting within
the U.S. Exclusive Economic Zone**

by

Frank T. Manheim

U. S. Geological Survey
Woods Hole, MA 02543

and

Allyn Vine

Woods Hole Oceanographic Institution
Woods Hole, MA 02543

March 1987

Technical Report

Reproduction in whole or in part is permitted for any purpose of the
United States Government. This report should be cited as:
Woods Hole Oceanog. Inst. Tech. Rept. WHOI-87-12.

Approved for publication; distribution unlimited.

Approved for Distribution:

David A. Ross, Chairman
Department of Geology and Geophysics

OPTIONS FOR RADIOACTIVE AND OTHER HAZARDOUS WASTE SITING WITHIN
THE U.S. EXCLUSIVE ECONOMIC ZONE

by

Frank T. Manheim* and Allyn Vine**
October, 1986

ABSTRACT

Some areas of the E.E.Z. (Exclusive Economic Zone) offer technical, political and economic options that may complement existing approaches to hazardous waste storage and disposal.

NOTES AND ADDENDA

A shortened version of this report, edited for a wide audience, but without references was published in Geotimes, v. 31, December, 1986, (p. 8-10) under the title "E.E.Z. may have waste-disposal options".

As stated in Geotimes, this earlier draft with references is hereby made available to interested individuals.

Appendix I is an additional list of references that may be of interest.

Appendix II is a copy of the two pages of references from the May 1986 Office of Technology Assessment Staff Paper on the Sub-seabed Disposal of High-Level Radioactive Waste.

Again the authors would like to thank the many people who have contributed ideas, criticism, and references.

* U.S. Geological Survey, Woods Hole, MA 02543

** Woods Hole Oceanographic Institution, Woods Hole, MA 02543

**OPTIONS FOR RADIOACTIVE AND OTHER HAZARDOUS WASTE SITING WITHIN
THE U.S. EXCLUSIVE ECONOMIC ZONE**

Introduction

The long-term storage and disposal of concentrated, high-level radioactive and other highly toxic wastes remains a difficult and controversial issue. Over 300,000 and 19,000 m³ (over 1,300 and 35,000 million curies) of defense and commercial high-level radioactive waste, respectively, are projected to require disposal by the end of 2000 (Office of Technology Assessment, 1985). In the United States most of the funded effort applied to disposal siting has been on land sites, with only a small scientific effort devoted to research and development on disposal beneath the deep ocean bottom. As a part of a continuing effort to find reliable and politically and economically satisfactory solutions for the disposal problem, we suggest that some geographic segments of the new U.S. Exclusive Economic Zone (EEZ) offer technical and political options that may complement constructively the existing approaches to waste storage final disposal.

Present status for land storage and disposal

In the United States, most research and development on high-level radioactive-waste disposal in deep geological repositories has been focused on nine potential sites, distributed in six States, for the first repository, Washington, Nevada, Utah (two), Texas (two), Louisiana, and Mississippi (two) (Fig. 1a). Of these sites, seven are in salt and one each in tuff and basalt. Very recently Nevada, Washington, and Texas have been given priority by the Department of Energy (DOE, 1986). Following provisions of the Nuclear Waste Policy Act of 1982, four additional regions in the north-central and

eastern part of the United States were considered for the second repository (Fig. 1b). However, the apparent growing opposition of State governments to any nuclear waste sites within their borders poses at least temporary political barriers to permanent emplacement of wastes in geological repositories within the continental United States.

Present status of subseabed storage and disposal

Another seriously considered disposal option is permanent disposal within unconsolidated sediments (pelagic clays) under the central deep ocean bed far from land. Research on this option by the United States and several other nuclear nations has focused on international waters for technical and cooperative reasons (Office of Technology Assessment, 1986), and a rewarding international cooperation among principal user nations has been achieved (Sandia Laboratories, 1983; International Atomic Energy Agency, 1983). The main study areas for the Deep Ocean Nuclear Waste Disposal Program are shown in Fig. 2. Efforts were made to find standards and methodologies acceptable to all users of nuclear power and to nonusers and other coastal nations. However, this international program is threatened by existing and potential new restrictions on nuclear waste disposal in international areas of the ocean. Blanket restrictions against nuclear waste disposal in international waters, and even against research involving radionuclides, are being proposed for the London Dumping Convention, to which the United States is a signatory.

Implications of the EEZ for disposal options

The EEZ may offer a third geographic option. The offshore EEZ areas constitute about 15,800,000 km², compared with 9,370,000 km² of U.S. land territory (McGregor and Offield, 1984). A part of this area is shown in

Figs. 3 and 4. The EEZ covers a wide choice of geological substrates, including salt domes, igneous rock bodies, relatively flat-lying extensions of coastal plain strata, and such deep-water pelagic sediments as those considered for deep-ocean waste emplacement. As with site selections on land or in the deep sea, choices would trend toward areas of minimal commercial, resource, and tectonic activities.

Some segments of the EEZ, beyond the State territorial seas but within the 200-nm limit, offer logical areas for research and development, training, storage and perhaps permanent disposal systems. Initially, these may be considered backup or complementary sites to principal on-land sites in the context of the Nuclear Waste Policy Act of 1982. We suggest the following opportunities for these purposes: (1) sediment areas that display the main pelagic clay characteristics of deep ocean disposal sites but occurring within 200 nm of U.S. coasts, (2) low permeability rocks which may include granites, basalts, and, especially, rock salt, (3) bedded strata into which caverns or lined shafts might be placed for temporary (decades to centuries) monitoring and storage by drilled emplacement beneath the sea floor (Bury, 1985), and (4) development and training sites.

Deep-water hemipelagic and pelagic clay sites are especially well developed within the EEZ off the coast of California and Oregon (Fig. 3) and off the Pacific Islands Trust Territory. Although igneous rocks like granite and basalt are found off the Atlantic and the Pacific coasts, the rock type least permeable through fracturing is rock salt. Among the most extensive offshore salt deposits in the world are those in the northern Gulf of Mexico in the form of salt domes and ridges (Fig. 4); other deposits exist off the Atlantic margin. Many domes are well studied because some serve as traps for petroleum hydrocarbons. Training sites might be located reasonably close to

logistic ports. They would simulate more remote sites for the purpose of testing investigatory, emplacement, monitoring, and retrieval systems. Of particular importance, they could serve to better train crew and engineers to conduct operational and engineering functions at sea more easily, reliably, and cheaply than at more remote (including midocean) final repositories.

Discussion

Development of engineering expertise for EEZ disposal systems would draw upon both land and deep-ocean (including offshore petroleum) technology, as well as new scientific and technological approaches. Normally, it is assumed that offshore operations are more difficult than those on land. However, the constraints of work with hazardous materials in inhabited areas complicate and add to the expense of land activity. In addition, some aspects of the EEZ geological and physical environments will be more favorable than those of land environments for waste disposal in comparable substrates; for example, the artesian systems that drive meteoric water circulation through land aquifers are believed to be largely absent offshore (Manheim and Sayles, 1974). With ion movement largely limited to diffusive gradients, salt domes under the ocean should be far less erodable and leachable than comparable land forms. In addition, submarine salt domes do not present a hazard to ground-water systems in areas of human habitation. Uncommon systems, such as possible upward hydrothermal circulation in rift zones (Sleep, 1985) or aquifers that are continuous between land and sea (Kohout et al., 1986) would be excluded from consideration.

Criteria for disposal/storage areas: first, we have limited consideration to storage or disposal and training sites within the U.S. EEZ and outside State waters*. Second, we assume that sites and methods will

allow for monitoring, retrieval, and, if need be, relocation of waste. The first criterion corresponds to the Federal responsibility and legal authority for nuclear waste disposal projected for 1998 by the Nuclear Policy Act of 1982. Temporary placement of waste within sediment or rocks beneath the U.S. territorial waters will reduce, though perhaps not wholly eliminate, potential foreign opposition to waste siting in the ocean. States retain input into decision-making regarding areas off their coasts through the Coastal Zone Management Act and other statutes.

The monitoring-retrieval criterion was mandated already by the Nuclear Regulatory Commission in the final rules for geologic repositories issued in 1983. We believe that with the evolution and practice of specific techniques retrievability is as feasible under the seabed as on land. Monitorability and retrievability can allow for early detection of potential problems and transfer of waste products should on-site problems be identified or should better sites be developed. These safeguards are likely to increase public acceptability of waste sites. Variable storage time frames (decades to centuries) provide a variety of options, such as special EEZ site or sites developed for extended cooling periods, to supplement special land cooling or storage areas, to provide more time to study "permanent" sites that have optimum storage properties. Finally, in addition to high-level nuclear waste, we suggest that with modified or other less costly techniques, the EEZ could serve as a safe repository or storage area for nonradioactive hazardous wastes having lesser hazard.

*For international claim purposes the EEZ is taken from the coast to 200 nm seaward, according to the proclamation of March 10, 1983, by President Reagan. State waters usually extend to 3 nm from shore.

Conclusion

In the past, the geographical equivalents of the EEZ were considered less attractive than land or the central deep ocean for high level radioactive waste disposal. However, selected areas of the huge U.S. EEZ now offer more attractive options for radioactive waste disposal and storage.

The storage and disposal concepts discussed above should be applicable to hazardous waste of either nuclear or non-radioactive type. For example, successful storage/disposal in the EEZ might aid in solution of land disposal programs, especially toxic or corrosive chemical wastes that might cause ground water contamination. The information and experience gained in this effort might be helpful in assessing applicability of EEZ areas to the placement of high-level radioactive waste. However, the non-nuclear hazardous waste program might be a separate and distinct program, not necessarily involving commitment to nuclear waste programs.

Appropriate investigation of these options requires a multidisciplinary research program to which Federal agencies and private and commercial organizations could contribute. We assume that the next few decades will continue to bring forward new physical data and changing political-economic emphases. The authors hope that this short report will stimulate discussion and broaden the field of possible alternatives to solve our waste disposal problems.

References

- Addicott, W.O., 1985, Coordinator, Manganese nodules, seafloor sediment, and sedimentation rates of the circum-Pacific regions: American Association of Petroleum Geologists, Tulsa, OK (map).
- Bury, M.R.C., 1985, The offshore disposal of radioactive waste by drilled emplacement: a feasibility study: Graham and Trotman, Ltd., London, 13 chapters, chapter-paginated.
- Department of Energy, 1986, DOE announces decisions on high-level waste repository program, Press Release, May 28, 1986. DOE, Washington D.C.
- Dept. of Energy, Publications Office, Chicago, Ill.
- Department of Energy, 1986a, Area recommendation report for the crystalline repository project, Volumes I and II, DOE-CH-15(1) prepared for the crystalline repository project by the Office of Crystalline Repository Development, Battelle Project Management Division, Argonne Ill.
- Kohout, F.A., Meisler, H., Meyer, F.W., Johnston, R.H., Leve, G.W., and Wait, R.L., 1986, Hydrogeology of the Atlantic continental margin. Decade of North American Geology, Ch. 6b, in press.
- International Atomic Energy Agency, 1983, Waste Management Research Abstracts, NBo. 14, International Atomic Energy Agency, Vienna, 287 p.
- Manheim, F.T., and Sayles, F.L., 1974, Composition and origin of interstitial waters of marine sediments, based on deep sea drill cores, in Goldberg, E.D. (ed). The Sea: Marine Chemistry, v. 5, p. 527-568.
- Martin, R.F., 1980, Distribution of salt structures in the Gulf of Mexico, U.S. Geological Survey M.F. 1213, Map and descriptive text.
- McGregor, B.A. and Offield, T.W., 1984, The exclusive economic zone: an exciting new frontier. Washington, D.C., U.S. Government Printing Office, 20 p.

Office of Technology Assessment, 1985, Managing the Nation's high-level commercial radioactive waste. U.S. Congress, Office of Technology Assessment, Pub. OTA-0171, 348 p.

Office of Technology Assessment, 1986, Subseabed Disposal of High-Level Radioactive Waste. Staff Paper, Oceans and Environment Program. 42 p.

Sandia Laboratories, 1983, The Subseabed Disposal Program: 1983 Status Report. Albuquerque NM, Sandia National Laboratories, Sandia Report SA D83-1887, 179 p.

Sleep, N.H., 1985, Hydrothermal convection at ridge axes. In Rona, P.A., Bostrom, K., Laubier, L., and Smith, K.L., Jr. (Eds), Hydrothermal processes at seafloor spreading centers. Plenum Press, N.Y. and London, p. 71-82.

List of Figures

- Fig. 1. Location of candidate sites for the first deep geologic repository for high-level radioactive waste (from Office of Technology Assessment, 1985) "Sites" are named.
- Fig. 1b. Location of study areas for a second deep-ocean high-level radioactive waste disposal site (from U.S. Dept. Energy, 1986a).
- Fig. 2. Areas selected for the Deep Ocean Nuclear Waste Disposal Program (from Sandia Laboratories, 1983).
- Fig. 3. Hemipelagic and pelagic clay areas in the U.S. West Coast EEZ. Heavy stippling denotes gravel, sand, and silty deposits. Light stippling indicates other terrigenous sediments, including sediments in southern California offshore basins. Area shaded with irregular subparallel oblique lines refers to hemipelagic and pelagic sediments having accumulation rates of less than 150 mm/1000 years. Vertical shading refers to pelagic sediments having accumulation rates of less than 50 mm/1000 years. Heavy line denotes EEZ boundary. Map modified largely from Addicott (1985).
- Fig. 4. Distribution of salt domes in northern Gulf of Mexico area. Black areas represent salt domes and ridges. Hollow areas offshore represent possible deep salt "pillows." Diagram modified from Martin (1980), by K. Klitgord, U.S. Geological Survey, Woods Hole, MA 02543. Heavy line denotes EEZ boundary.

F.T. Manheim, U.S. Geological Survey, Woods Hole Ma 02543

A. Vine, Woods Hole Oceanographic Institution, Woods Hole Ma 02543

2.

4.

Appendix I

Selected Supplementary References for Hazardous Waste Disposal in U.S. EEZ

Anonymous, 1986. Staff paper on the subseabed disposal of high-level radioactive waste: a summary of the Oceans and Environment Program, prepared by the Office of Technology Assessment, Washington, D.C. 20510. 42 pages.

Carter, Luther V., 1987, Nuclear Imperatives and Public Trust Dealing with Radioactive Waste ,pages 46-61, Issues in Science and Technology, winter 1987, .

Frosch, R.A., Hollister, C.D., Heath, G.R., Silva, A.J., Hessler, R.R., Jumars, P.A., Deese, D.A., High-level Nuclear Wastes in the Seabed. In Oceanus, Winter, 1977, Volume 20, Number 1.

Jenyon, M.K., 1986, Salt with a pinch of water, Nature, vol. 324, pages 515-516.

Milnes, A. G., 1985. Geology and radwaste. Academic Press, London, Orlando, 328 p.

Neiheisel, J. 1983, Prediction parameters of radionuclide retention at low-level radioactive waste sites. U.S. Environmental Protection Agency, Office of Radiation Programs, Washington, D.C., 27 p.

Noshkin, V.E., Wong, Kai M., Jokela, T.A., Eagle, R.J., and Brunk, J.L., 1978. Radionuclides in the marine environment near the Farallon Islands. Lawrence Livermore Laboratory, UCRL 52381, 17 pp. Authors determined total plutonium and 137 cesium inventory in the Farallon water columns. Remobilization of waste products from the disposal site is discussed.

Office of Technology Assessment, 1985, Managing the Nations's high-level commercial radioactive waste. U.S. Congress, Office of Technology Assessment, Pub. OTA-0171, 348 p.

Office of Technology Assessment, 1986, Subseabed Disposal of High-Level Radioactive Waste. Staff Paper, Oceans and Environment Program. 42 p.

Sandia Laboratories, 1983, The Subseabed Disposal Program: 1983 Status Report. Albuquerque NM, Sandia National Laboratories, Sandia Report SA D83-1887, 179 p.

Schell, W. R., and Sugai, S., 1980. Radionuclides at the U.S. radioactive waste disposal site near the Farallon Islands. Health Physics, vol. 39, p. 475-496. Investigation of canisters of low-level radioactive waste at water depths of 914-1829 m W. of San Francisco Bay. 137 Cs, 239, 240 Pu, 90 Sr and 238 Pu were analyzed in sediments and fish.

Sleep, N.H., 1985, Hydrothermal convection at ridge axes. In Rona, P.A., Bostrom, K., Laubier, L., and Smith, K.L., Jr. (Eds), Hydrothermal processes at seafloor spreading centers. Plenum Press, N.Y. and London, p. 71-82.

Urai, J.L., Spiers, C.J., Zwart, H.J., Lister, G.S., 1986, Weakening of rock salt by water during long term creep, Nature, vol. 324, p.554-557.

Appendix II

Staff Paper on the Subseabed Disposal of High-Level Radioactive Waste

Office of Technology Assessment, May 1986, Washington, D.C. 20510

REFERENCES

- Anderson, D.R., "The Concept of Disposing of High-Level Radioactive Wastes in the Stable Geologic Formations Underlying the Ocean Floor," SAND84-0031, Sandia National Laboratories, February 1984.
- Anderson, D.R. (ed.), "Engineering Studies Task Group," *Tenth Annual Meeting of the NEA Coordinated Program on the Assessment of the Subseabed Disposal of Radioactive Waste*, SAND85-1365, Halifax, Nova Scotia, Apr. 30 - May 3, 1985, in press.
- Brush, L.H., "Subseabed Radionuclide Migration Studies and Preliminary Repository Design Concepts," *The Technology of High-Level Nuclear Waste Disposal, Volume 2*, Hofmann, P.L. (ed.), U.S. Department of Energy Technical Information Center, 1984, pp. 287-294.
- Burdett, J.R.F., and Karnes, C.H. (eds.), "Report on the Sixth Interim Meeting of the Seabed Working Group's Engineering Studies Task Group," ISMES SpA., Rome, Italy, Oct. 7-11, 1985.
- Bury, M.R.C., "The Offshore Disposal of Radioactive Waste by Drilled Emplacement: A Feasibility Study" (Commission of European Communities, London: Graham and Trotman Ltd., 1985).
- Ensminger, D.A., Koplik, C.M., and Klett, R.D., "Preliminary Pre-Emplacement Safety Analysis of the Subseabed Disposal of High-Level Nuclear Waste," SAND83-7105, Sandia National Laboratories, January 1985.
- Hartwig, E., Office of Naval Research, personal communication, Feb. 21, 1986.
- Heath, G.R., C.D. Hollister, D.R. Anderson, and M. Leinen, "Why Consider Subseabed Disposal of High-Level Nuclear Wastes?" *Wastes in the Ocean: Volume 3 Radioactive Wastes and the Ocean*, Park, P.K., Kester, D.R., Duedall, I.W., and Ketchum, B. (eds.) (New York: John Wiley & Sons, 1983).
- Hickerson, J., unpublished design and test information, to be included in the 1986 Annual Report on Engineering Studies and the 1986 Status Report of the Subseabed Disposal Project, Sandia National Laboratories.
- Interagency Review Group on Nuclear Waste Management, "Report to the President," T10-29442, Washington, DC, March 1979.
- International Atomic Energy Agency (IAEA), "The Radiological Basis of the IAEA Revised Definition and Recommendations Concerning High-Level Radioactive Waste Unsuitable for Dumping at Sea," IAEA-211, 1978.
- JK Associates; "The Subseabed Disposal Project: Briefing Book 1985," Austin, Texas, 1985.
- Jumars, P., Sandia National Laboratories, personal communication, Sept. 3, 1985.
- Kaplan, M.F. and Klett, R.D., "Biological and Physical Oceanographic Sensitivity Analysis for Subseabed Disposal of High-Level Waste," SAND83-7107, Sandia National Laboratories, November 1984.
- Kaplan, M.F., Klett, R.D., Koplik, C.M., and Ensminger, D.A., "Radiological Protection Options for Subseabed High-Level Waste Disposal," SAND84-0548, Sandia National Laboratories, March 1985.
- Karnes, C.H., Sandia National Laboratories, personal communication, Apr. 9, 1986.
- League of Women Voters Education Fund, "The Nuclear Waste Primer: A Handbook for Citizens" (New York: Nick Lyons Books, 1985).
- Livingston, H.D., Gowen, V.T., Kasso, S.A., Volchok, H.L., Noshkin, V.E., Wong, K.M., and Beasley, T.M., "Fallout Nuclides in Atlantic and Pacific Water Columns: GEOSECS Data," Woods Hole Oceanographic Institution 85-19, May 1985.

- Marietta, M.G. and Robinson, A.R., "Status and Outlook of Ocean Modeling Research Dispersion and Related Applications," SAND85-2806, Sandia National Laboratories, December 1985.
- National Advisory Committee on Oceans and Atmosphere (NACOA), "Nuclear Waste Management and the Use of the Sea," 1984.
- Nuclear Regulatory Commission, "An Evaluation of Radionuclide Concentrations in High-Level Radioactive Wastes," NUREG-0946, October 1985.
- Organization for Economic Co-operation and Development (OECD), "Seabed Disposal of HLRW: A Status Report of the NEA Coordinated Research Programme," Nuclear Energy Agency, 1984.
- Ove Arup and Partners, "Studies of Penetrator Emplacement," Commission of European Communities, in press 1986.
- Richter, K., Bennett, J.C., and Smith, K.L., Jr., "Bottom-Moored Acoustic Array To Monitor Density and Vertical Movement of Deep-Sea Benthos-Pelagic Animals," *IEEE Journal of Ocean Engineering* OE-10:32-37, January 1985.
- Robinson, A.R., and Marietta, M.G., "Research, Progress and the Mark A Model for Physical, Biological and Chemical Transports," SAND84-0646, Sandia National Laboratories, March 1985.
- Rowe, G., Brookhaven National Laboratory, personal communication at OTA Workshop on Subseabed Disposal of HLRW, Sept. 4, 1985.
- Sandia National Laboratories, "The Subseabed Disposal Program: 1983 Status Report," SAND83-1387, October 1983.
- Schreiner, F.S., Fried, S., and Friedman, A.M., "Measurement of Radionuclide Diffusion in Ocean Floor Sediment and Clay," *Nuclear Technology* 59:429-438, 1982.
- Taylor, P.J., Political Ecology Research Group, Ltd., personal communication, Feb. 10, 1986.
- Thornton, E.C. and Seyfried, W.E., Jr., "Sediment-Seawater Interaction at 200° and 300° C, 500 Bars Pressure: The Role of Sediment Composition in Diagenesis and Low-Grade Metamorphism of Marine Clay," *Geological Society of America Bulletin* 96:1287-1295, October 1985.
- U.S. Congress, Office of Technology Assessment, "Managing the Nation's Commercial High-Level Radioactive Waste" OTA-O-171, (Washington, DC: U.S. Government Printing Office, 1985a).
- U.S. Congress, Office of Technology Assessment, "Workshop on Subseabed Disposal of HLRW," Washington, DC, Sept. 4, 1985b.
- U.S. Congress, Office of Technology Assessment, "Oil and Gas Technologies for the Arctic and Deepwater" OTA-O-270, (Washington, DC: U.S. Government Printing Office, 1985c).
- U.S. Department of Energy, "Spent Fuel and Radioactive Waste Inventories, Projections, and Characteristics," DOE/RW-0006, Rev. 1, December 1985.
- U.S. Department of Energy, "Report to OMB on the Subseabed Disposal Concept," Sept. 26, 1983.
- U.S. House of Representatives, "Achieving Performance Objectives for the Engineered Barrier System," staff report prepared by the Subcommittee on Energy Conservation and Power of the Committee on Energy and Commerce (Washington, DC: Government Printing Office, 1984).
- Yayanos, A.A. and Nevenzel, J.C., "Rising-Particle Hypothesis: Rapid Ascent of Matter From the Deep Ocean," *Naturwissenschaften* 65:255, 1978.

REPORT DOCUMENTATION PAGE		1. REPORT NO. <i>WHOI-87-12</i>	2.	3. Recipient's Accession No.
4. Title and Subtitle <i>Options for Radioactive and Other Hazardous Waste Siting within the U.S. Exclusive Economic Zone</i>		5. Report Date <i>March 1987</i>		6.
7. Author(s) <i>Frank T. Manheim and Allyn Vine</i>		8. Performing Organization Rept. No. <i>WHOI-87-12</i>		9. Performing Organization Name and Address <i>Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543</i>
10. Project/Task/Work Unit No.		11. Contract(C) or Grant(G) No. (C) (G)		12. Sponsoring Organization Name and Address
13. Type of Report & Period Covered <i>Technical</i>		14.		
15. Supplementary Notes <i>This report should be cited as: Woods Hole Oceanog. Inst. Tech. Rept., WHOI-87-12.</i>				
16. Abstract (Limit: 200 words) <i>Some areas of the E.E.Z. (Exclusive Economic Zone) offer technical, political and economic options that may complement existing approaches to hazardous waste storage and disposal.</i>				
17. Document Analysis a. Descriptors b. Identifiers/Open-Ended Terms c. COSATI Field/Group				
18. Availability Statement: <i>Approved for publication; distribution unlimited.</i>		19. Security Class (This Report) <i>UNCLASSIFIED</i>	21. No. of Pages <i>16</i>	
		20. Security Class (This Page)	22. Price	

DOCUMENT LIBRARY

November 21, 1986

Distribution List for Technical Report Exchange

Institute of Marine Sciences Library
University of Alaska
O'Neill Building
905 Koyukuk Ave., North
Fairbanks, AK

Attn: Stella Sanchez-Wade
Documents Section
Scripps Institution of Oceanography
Library, Mail Code C-075C
La Jolla, CA 92093

Hancock Library of Biology & Oceanography
Alan Hancock Laboratory
University of Southern California
University Park
Los Angeles, CA 90089-0371

Gifts & Exchanges
Library
Bedford Institute of Oceanography
P.O. Box 1006
Dartmouth, NS, B2Y 4A2, CANADA

Office of the International
Ice Patrol
c/o Coast Guard R & D Center
Avery Point
Groton, CT 06340

Library
Physical Oceanographic Laboratory
Nova University
8000 N. Ocean Drive
Dania, FL 33304

NOAA/EDIS Miami Library Center
4301 Rickenbacker Causeway
Miami, FL 33149

Library
Skidaway Institute of Oceanography
P.O. Box 13687
Savannah, GA 31416

Institute of Geophysics
University of Hawaii
Library Room 252
2525 Correa Road
Honolulu, HI 96822

Library
Chesapeake Bay Institute
4800 Atwell Road
Shady Side, MD 20876

MIT Libraries
Serial Journal Room 14E-210
Cambridge, MA 02139

Director, Ralph M. Parsons Laboratory
Room 48-311
MIT
Cambridge, MA 02139

Marine Resources Information Center
Bldg. E38-320
MIT
Cambridge, MA 02139

Library
Lamont-Doherty Geological Observatory
Columbia University
Palisades, NY 10964

Library
Serials Department
Oregon State University
Corvallis, OR 97331

Pell Marine Science Library
University of Rhode Island
Narragansett Bay Campus
Narragansett, RI 02882

Working Collection
Texas A&M University
Dept. of Oceanography
College Station, TX 77843

Library
Virginia Institute of Marine Science
Gloucester Point, VA 23062

Fisheries-Oceanography Library
151 Oceanography Teaching Bldg.
University of Washington
Seattle, WA 98195

Library
R.S.M.A.S.
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149

Maury Oceanographic Library
Naval Oceanographic Office
Bay St. Louis
NSTL, MS 39522-5001
ATTN: Code 4601