Reference No. 84-11

ABSTRACTS OF PAPERS
SUBMITTED IN 1983 FOR PUBLICATION

Woods Hole Oceanographic Institution
Woods Hole, Massachusetts

Editor: Anne I. Rabushka

APPROVED FOR DISTRIBUTION

When citing this report, it should be referenced
as: Woods Hole Oceanographic Institution
Technical Report WHOI-84-11.

MBL/WHOI

0 0301 0096524 0

PREFACE

This volume contains all abstracts submitted for publication during calendar year 1983 by the staff and students of the Woods Hole Oceanographic Institution. Because some of the abstracts may not be published in the journal to which they have been submitted initially, we have purposely omitted identifying the journals. The volume is intended to be informative, but not a bibliography.

The abstracts are listed by title in the Table of Contents and are grouped into one of our five departments, marine policy, or the student category. An author index is presented in the back to facilitate locating specific papers.

A handwritten signature in dark ink, appearing to read 'Derek W. Spencer', is written over a horizontal line.

Derek W. Spencer
Associate Director for Research

DEPARTMENT OF BIOLOGY

B-1

Animal Physiology and Biochemistry

MICROSOMAL ELECTRON TRANSPORT AND XENOBIOTIC MONOOXYGENASE ACTIVITIES DURING THE EMBRYONIC PERIOD OF DEVELOPMENT IN THE KILLIFISH, <u>Fundulus Heteroclitus</u> . Robert L. Binder and John J. Stegeman	B-3
BLUEFIN TUNA WARM THEIR VISCERA DURING DIGESTION. Francis G. Carey, John W. Kanwisher and E. Don Stevens	B-3
PECTORAL FIN LOCOMOTION: SWIMMING ENERGETICS OF CUNNERS AND TAUTOGS (LABRIDAE). Marvin A. Freadman	B-3
SWIMMING/LOCOMOTOR CAPACITY OF HUDSON RIVER STRIPED BASS. Marvin A. Freadman, Frederick P. Thurberg and Anthony C. Calabrese	B-4
INTRODUCTION TO THE BIOLOGY OF HYDROTHERMAL VENTS. J. Frederick Grassle	B-4
AN ALTERNATIVE 7-ETHOXYRESORUFIN O-DEETHYLASE ACTIVITY ASSAY: A CONTINUOUS VISIBLE SPECTROPHOTOMETRIC METHOD FOR MEASUREMENT OF CYTOCHROME P-450 MONOOXYGENASE ACTIVITY Alan V. Klotz, John J. Stegeman and Chris Walsh	B-4
AN ARYL HYDROCARBON HYDROXYLATING HEPATIC CYTOCHROME P-450 FROM THE MARINE FISH <u>Stenotomus Chrysops</u> . Alan V. Klotz, John J. Stegeman and Christopher Walsh	B-5
PHYSIOLOGY OF THE WOOD BORING MOLLUSC <u>Martesia cuneiformis</u> SAY. Roger Mann and Scott M. Gallager	B-5
SWIMMING BEHAVIOUR OF LARVAE OF THE OCEAN QUAHOG <u>Arctica islandica</u> IN RESPONSE TO PRESSURE AND TEMPERATURE. Roger Mann and Charlotte C. Wolf	B-6
HYPOSMOREGULATION IN AN ANADROMOUS TELEOST: INFLUENCE OF SEX AND MATURATION. Stephen D. McCormick and Robert J. Naiman	B-6
SOME DETERMINANTS OF MATURATION IN BROOK TROUT, <u>Salvelinus fontinalis</u> . Stephen D. McCormick and Robert J. Naiman	B-6
DIFFERENTIAL REGULATION OF HEPATIC XENOBIOTIC AND STEROID METABOLISM IN MARINE TELEOST SPECIES. John J. Stegeman and Bruce R. Woodin	B-7
FOOD SELECTION BY LARVAL GULF MENHADEN (<u>Brevoortia patronus</u>). Diane Stoecker and John J. Govoni	B-7

Aquaculture

BIVALVE MOLLUSC HATCHERIES: A CRITICAL APPRAISAL OF THEIR DEVELOPMENT AND A REVIEW OF THEIR POTENTIAL VALUE IN ENHANCING THE FISHERIES OF DEVELOPING NATIONS. Roger Mann	B-7
THE ROLE OF INTRODUCED BIVALVE MOLLUSC SPECIES IN MARICULTURE. Roger Mann	B-8
GROWTH-STASIS AND LIMITED SHELL CALCIFICATION IN LARVAE OF THE GASTROPOD <u>Cymatium parthenopeum</u> DURING TRANS-ATLANTIC TRANSPORT. Jan A. Pechenik, Rudolf S. Scheltema and Linda S. Eyster	B-8
SEASONAL VARIATION IN THE NITRATE CONTENT OF WATER HYACINTH (<u>Eichhornia crassipes</u> [MART.] SOLMS). C. S. Tucker and Thomas A. Debusk	B-9

Ecosystem StudiesPredation

PREDATION ON FISH LARVAE BY <u>Physalia</u> , THE PORTUGUESE MAN OF WAR. Jennifer E. Purcell	B-9
---	-----

Riverine Systems

NITROGEN FIXATION IN BEAVER (<u>Castor canadensis</u>) INFLUENCED SUBARCTIC STREAMS. Margaret M. Francis, Robert J. Naiman and Jerry M. Melillo	B-9
--	-----

RESOURCE PARTITIONING AND AGGRESSIVE INTERACTIONS BETWEEN BROOK TROUT (<i>Salvelinus fontinalis</i>) FRY AND NINESPINE STICKLEBACKS (<i>Pungitius pungitius</i>).	B-10
Alaine Gaudreault, Tandy Miller, Gerry J. FitzGerald and W. Linn Montgomery	
DYNAMICS OF PLANT LITTER DECAY IN NORTHERN STREAMS.	B-10
Jerry M. Melillo, Robert J. Naiman, John D. Aber and Alan E. Linkins	
NITROGEN BUDGET OF A SUBARCTIC STREAM ALTERED BY BEAVER (<i>Castor canadensis</i>).	B-10
Robert J. Naiman and Jerry M. Melillo	
 Salt Marsh Systems	
WATER UPTAKE BY ROOTS IN SHORT <i>Spartina</i> MARSH: MASS FLOW OF AIR INTO SEDIMENT AND INTO PLANTS	B-11
John W. H. Dacey and Brian L. Howes	
CARBON FLOW THROUGH OXYGEN AND SULFATE REDUCTION PATHWAYS IN SALT MARSH SEDIMENTS.	B-11
Brian L. Howes, John W. H. Dacey and Gary M. King	
ANNUAL CARBON REMINERALIZATION AND BELOW-GROUND PRODUCTION OF <i>Spartina alterniflora</i> IN A NEW ENGLAND SALT MARSH.	B-11
Brian L. Howes, John W. H. Dacey and John M. Teal	
EFFECTS OF SAMPLING TECHNIQUE ON MEASUREMENTS OF POREWATER CONSTITUENTS IN SALT MARSH SEDIMENTS	B-12
Brian L. Howes, John W. H. Dacey and Stuart Wakeham	
EFFECTS OF O ₂ ON AMMONIUM UPTAKE AND ROOT RESPIRATION BY <i>Spartina alterniflora</i> .	B-12
James T. Morris and John W. H. Dacey	
THE ROLE OF ONE SALT MARSH IN COASTAL PRODUCTIVITY.	B-12
John M. Teal	
THE IMPORTANCE OF CHEMICAL COMPOSITION OF SALT MARSH LITTER ON DECAY RATES AND FEEDING BY DETRITIVORES.	B-12
Ivan Valiela, John Wilson, Robert Buchsbaum, Carol Rietsma, Donald Bryant, Kenneth Foreman and John Teal	
 Marine Mammals	
ANATOMY AND VASCULATURE OF A MINKE WHALE HEART.	B-13
Christina Ochrymowych and Richard Lambertsen	
STATUS OF <i>Eubalaena glacialis</i> OFF CAPE COD.	B-13
William E. Schevill, William A. Watkins and Karen E. Moore	
SONAR OBSERVATIONS EXPLAIN BEHAVIORS NOTED DURING BOAT MANEUVERS FOR RADIO TAGGING OF HUMPBACK WHALES (<i>Megaptera novaeangliae</i>) IN THE GLACIER BAY AREA.	B-14
William A. Watkins and Camille A. Goebel	
SOUNDS OF THE ROSS SEAL, <i>Ommatophoca rossi</i> .	B-14
William A. Watkins and G. Carleton Ray	
 Marine Pollution	
THE EFFECTS OF POLLUTANTS ON MARINE ZOOPLANKTON POPULATIONS: SUBLETHAL PHYSIOLOGICAL INDICES OF POPULATION STRESS.	B-14
Judith M. Capuzzo	
THE IMPACT OF WASTE DISPOSAL IN NEAR SHORE ENVIRONMENTS.	B-15
Judith M. Capuzzo, Wayne V. Burt, Iver W. Duedall, Dana R. Kester and P. Kilho Park	
THE EFFECTS OF PETROLEUM HYDROCARBONS ON LIPID METABOLISM AND ENERGETICS OF LARVAL DEVELOPMENT AND METAMORPHOSIS IN THE AMERICAN LOBSTER (<i>HOMARUS AMERICANUS</i> MILNE EDWARDS).	B-15
Judith M. Capuzzo, Bruce A. Lancaster and Glenn C. Sasaki	
EFFECTS OF EXPOSURE TO SUBLETHAL CONCENTRATIONS OF CRUDE OIL ON THE COPEPOD <i>Centropages hamatus</i> . II. ACTIVITY PATTERNS.	B-16
Timothy J. Cowles	
EFFECTS OF EXPOSURE TO SUBLETHAL CONCENTRATIONS OF CRUDE OIL ON THE COPEPOD <i>Centropages hamatus</i> . I. FEEDING AND EGG PRODUCTION.	B-16
Timothy J. Cowles and James F. Remillard	
PHYSIOLOGICAL AND BIOCHEMICAL EFFECTS OF THE POLYCYCLIC AROMATIC HYDROCARBON BENZ(α)ANTHRACENE ON THE DEPOSIT FEEDING POLYCHAETE <i>Nereis virens</i> .	B-16
Anne E. McElroy	

EFFECTS OF DRILLING FLUIDS ON LARVAE OF THE AMERICAN LOBSTER, <u>Homarus americanus</u> . Jennifer G. Smith Derby and Judith M. Capuzzo	B-16
OIL SPILL STUDIES: A REVIEW OF ECOLOGICAL EFFECTS. John M. Teal and Robert W. Howarth	B-17
UNDERSTANDING SUBLETHAL POLLUTANT EFFECTS IN THE MARINE ENVIRONMENT. John Vandermeulen and Judith M. Capuzzo	B-17

Microbiology

<u>Chloroherpeton thalassion</u> GEN. NOV. OF SPEC. NOV., A NON-FILAMENTOUS, FLEXING AND GLIDING GREEN SULFUR BACTERIUM. Jane Gibson, Norbert Pfennig and John B. Waterbury	B-18
MICROBES IN THE OCEANIC ENVIRONMENT. Holger W. Jannasch	B-18
RECENT PROGRESS IN THE MICROBIOLOGY OF HYDROTHERMAL VENTS. Holger W. Jannasch and D. C. Nelson	B-18
DEEP SEA MICROBIOLOGY. Holger W. Jannasch and Craig D. Taylor	B-18
CHEMOAUTOTROPHIC GROWTH OF A MARINE <u>Beggiatoa</u> IN SULFIDE-GRADIENT CULTURES. Douglas C. Nelson and Holger W. Jannasch	B-18
CELLULOLYTIC NITROGEN-FIXING BACTERIUM CULTURED FROM THE GLAND OF DESHAYES IN SHIPWORMS (BIVALVIA: TEREDINIDAE). John B. Waterbury, C. Bradford Calloway and Ruth D. Turner	B-19
IN SITU STUDIES ON DEEP-SEA AMPHIPODS AND THEIR INTESTINAL MICROFLORA. Carl O. Wirsen and Holger W. Jannasch	B-19

Modeling

AN INJUNCTIVE FORM OF THE AXIOMS OF TRIALECTICS. Hal Caswell	B-19
TRIALECTICS, CYBERNETICS, AND ZADEH'S THEORY OF STATE. Hal Caswell	B-19
ZADEH'S THEORY OF STATE AND ICHAZO'S TRIALECTIC ANALYSIS: CONVERGENT APPROACHES TO UNDERSTANDING DYNAMICS. Hal Caswell	B-20

Phytoplankton

COPPER COMPLEXATION DURING SPRING PHYTOPLANKTON BLOOMS IN COASTAL WATERS. Donald M. Anderson, John S. Lively and Ralph F. Vaccaro	B-20
PRODUCTION AND ROLE OF HYALINE HAIRS IN <u>Ceramium rubrum</u> . J. A. DeBoer and F. G. Whoriskey	B-21
UPTAKE AND ASSIMILATION OF AMMONIUM AND NITRATE BY PHYTOPLANKTON: INDICES OF NUTRITIONAL STATUS FOR NATURAL ASSEMBLAGES. Patricia M. Glibert and James J. McCarthy	B-21
CONCEPTUAL ROLE FOR MICROAGGREGATES IN PELAGIC WATERS. Joel C. Goldman	B-21
INFLUENCE OF BOTTLE PHOTOINHIBITION ON THE MEASUREMENT OF SEASONAL PRIMARY PRODUCTION IN A TEMPORATE COASTAL WATER. Joel C. Goldman and Mark R. Dennett	B-21

Dinoflagellates

THE IMPORTANCE OF DORMANT CYSTS IN TOXIC DINOFLAGELLATE BLOOMS AND SHELLFISH TOXICITY. Donald M. Anderson	B-22
SEXUALITY AND CYST FORMATION IN THE DINOFLAGELLATE <u>Gonyaulax tamarensis</u> : I. CYST YIELD IN BATCH CULTURES. Donald M. Anderson, David M. Kulis and Brian J. Binder	B-22
SEXUALITY AND CYST FORMATION IN THE DINOFLAGELLATE <u>Gonyaulax tamarensis</u> : II. TIME-COURSE MEASUREMENTS OF PHOSPHORUS LIMITATION. Donald M. Anderson and Niels L. Lindquist	B-23

SEXUAL PROCESSES IN THE LIFE CYCLE OF THE BLOOM FORMING DINOFLAGELLATE	B-23
<u>Gyrodinium uncatenum</u> : A MORPHOGENETIC OVERVIEW.	
D. Wayne Coats, Mary A. Tyler and Donald M. Anderson	

Population Ecology

OPTIMAL LIFE HISTORIES AND AGE-SPECIFIC COSTS OF REPRODUCTION: TWO EXTENSIONS.	B-23
Hal Caswell	
EVALUATING THE CONSEQUENCES OF REPRODUCTION IN COMPLEX SALMONID LIFE CYCLES. . .	B-24
Hal Caswell, Robert J. Naiman and Rod Morin	
PLANKTONIC LARVAE ACT LIKE PASSIVE PARTICLES IN TURBULENT NEAR-BOTTOM FLOWS. . .	B-24
Cheryl Ann Hannan	
SEASONAL CHANGES IN THE DEPTH DISTRIBUTION OF BIVALVE LARVAE ON	B-24
THE SOUTHERN NEW ENGLAND SHELF.	
Roger Mann	
SEASONAL CHANGES IN THE DEPTH DISTRIBUTION OF LARVAE <u>Arctica islandica</u> L: . . .	B-25
A COMPARISON OF FIELD DATA FROM THE SOUTHERN NEW ENGLAND SHELF AND A LABORATORY GENERATED MODEL.	
Roger Mann	
RECRUITMENT OF COPEPOD NAUPLII INTO THE PLANKTON: THE IMPORTANCE OF	B-25
DIAPAUSE EGGS AND BENTHIC PROCESSES.	
Nancy Marcus	
VARIATION IN THE DIAPAUSE RESPONSE OF <u>Labidocera aestiva</u> (COPEPODA:CALANOIDA) . .	B-25
FROM DIFFERENT LATITUDES AND ITS IMPORTANCE IN THE EVOLUTIONARY PROCESS.	
Nancy H. Marcus	
THE INFLUENCE OF BEAVER (<u>Castor canadensis</u>) ON THE PRODUCTION DYNAMICS	B-26
OF AQUATIC INSECTS.	
Robert J. Naiman, Donald M. McDowell and Barbara S. Farr	
ENERGY BUDGET OF AN OMNIVOROUS ROCKY SHORE AMPHIPOD,	B-26
<u>Calliopius laevisculus</u> (KROYER).	
Judith B. Pederson and Judith M. Capuzzo	

Zooplankton

A TECHNIQUE FOR THE ENUMERATION OF HETEROTROPHIC AND PHOTOTROPHIC NANOPLANKTON . .	B-26
USING EPIFLUORESCENCE MICROSCOPY, AND A COMPARISON WITH OTHER PROCEDURES.	
David A. Caron	
PREDICTED AND OBSERVED FEEDING RATES OF THE SPINOSE PLANKTONIC FORAMINIFER . . .	B-27
<u>Globigerinoides sacculifer</u> .	
David A. Caron and Allan W. H. Bé	
GEOGRAPHIC, VERTICAL, SEASONAL AND DIEL DISTRIBUTIONS OF PHOTOTROPHIC AND . . .	B-27
HETEROTROPHIC COMPONENTS OF THE PICOPLANKTON AND NANOPLANKTON IN THE NORTH ATLANTIC.	
Paul G. Davis, David A. Caron, Paul W. Johnson and John McN. Sieburth	
COMPARISON OF BEHAVIORAL PATTERNS IN THE COPEPODS <u>Acartia tonsa</u> AND	B-27
<u>Centropages typicus</u> : A QUANTITATIVE VIDEO ANALYSIS.	
Timothy J. Cowles	
THE DEVELOPMENTAL STAGES OF <u>Longipedia americana</u> (COPEPODA:HARPACTICOIDA) . . .	B-28
REARED IN THE LABORATORY.	
Takashi Onbé	
THE FUNCTIONS OF NEMATOCYSTS IN PREY CAPTURE BY EPIPELAGIC SIPHONOPHORES . . .	B-28
(COELENTERATE, HYDROZOA).	
Jennifer E. Purcell	
PARTICLE PRODUCTION BY PLANKTONIC CILIATES.	B-28
Diane K. Stoecker	
FINE SCALE SPATIAL CORRELATIONS BETWEEN PLANKTONIC CILIATES AND DINOFLAGELLATES.	B-29
Diane K. Stoecker, Linda H. Davis and Donald M. Anderson	

Organic and Biological Chemistry

THE CONCH OF <u>Limacina</u> AND <u>Peraclis</u> (PTEROPODA) AND A MODEL FOR THE EVOLUTION OF PLANKTONIC GASTROPODS.	C-3
K. Bandel, A. Almogi-Labin, C. Hemleben and W. G. Deuser	
MEMBRANE LIPID FROM A DEEP-SEA HYDROTHERMAL VENT METHANOGEN: IDENTIFICATION OF A NEW MACROCYCLIC GLYCEROL DIETHER.	C-3
Paul B. Comita and Robert B. Gagosian	
STRUCTURAL ELUCIDATION OF A UNIQUE MACROCYCLIC MEMBRANE LIPID FROM A NEW, EXTREMELY THERMOPHILIC, DEEP-SEA HYDROTHERMAL VENT ARCHAEABACTERIUM, <u>Methanococcus jannaschii</u> .	C-3
P. B. Comita, R. B. Gagosian, H. Pang and C. E. Costello	
SUSPENDED PARTICULATE ORGANIC MATERIAL FROM HYDROTHERMAL VENT WATERS AT 21°N.	C-3
Paul B. Comita, Robert B. Gagosian and Peter M. Williams	
FLY-ASH PARTICLES INTERCEPTED IN THE DEEP SARGASSO SEA.	C-4
W. G. Deuser, K. Emeis, V. Ittekkot and E. T. Degens	
PROCESSES CONTROLLING THE DISTRIBUTION OF BIOGENIC COMPOUNDS IN RECENT SEDIMENTS.	C-4
Robert B. Gagosian	
REVIEW OF MARINE ORGANIC GEOCHEMISTRY.	C-4
Robert B. Gagosian	
SEASONALITY IN THE FLUXES OF SUGARS, AMINO ACIDS AND AMINO SUGARS TO THE DEEP SARGASSO SEA.	C-5
V. Ittekkot, W. G. Deuser and E. T. Degens	
PARTICULATE AMINO ACIDS IN THE SEA: EFFECTS OF PRIMARY PRODUCTIVITY AND BIOLOGICAL DECOMPOSITION.	C-5
Cindy Lee and Carolyn Cronin	
DISSOLVED, EXCHANGEABLE, AND BOUND ALIPHATIC AMINES IN MARINE SEDIMENTS: INITIAL RESULTS.	C-5
Cindy Lee and Brenda L. Olson	
TRANSFORMATION REACTIONS AND RECYCLING OF CAROTENOIDS AND CHLORINS IN THE PERU UPWELLING REGION (15°S, 75°W).	C-5
Daniel J. Repeta and Robert B. Gagosian	
PARTICLE SIZE DISTRIBUTIONS OF N-ALKANES AND ²¹⁰ Pb IN AEROSOLS OFF THE COAST OF PERU.	C-6
J. K. Schneider, R. B. Gagosian, J. K. Cochran and T. W. Trull	
FREE AND ESTERIFIED STEROLS OF THE MARINE DINOFLAGELLATE <u>Gonyaulax polygramma</u> .	C-6
J. K. Volkman, R. B. Gagosian and S. G. Wakeham	
BIOGEOCHEMISTRY OF PARTICULATE ORGANIC MATTER IN THE OCEANS: RESULTS FROM SEDIMENT TRAP EXPERIMENTS.	C-6
S. G. Wakeham, C. Lee, J. W. Farrington and R. B. Gagosian	
STERENES IN SUSPENDED PARTICULATE MATTER IN THE EASTERN TROPICAL NORTH PACIFIC.	C-7
S. G. Wakeham, R. B. Gagosian, J. W. Farrington and E. A. Canuel	

Seawater and Geochemistry

DSDP HOLE 504B, THE FIRST REFERENCE SECTION OVER 1 KM THROUGH LAYER 2 OF THE OCEANIC CRUST.	C-7
R. N. Anderson, J. Honnorez, K. Becker, A. C. Adamson, J. C. Alt, R. Emmermann, P. D. Kempton, H. Kinoshita, C. Laverne, M. J. Mottl and R. L. Newmark	
IN SITU ELECTRICAL RESISTIVITY AND BULK POROSITY OF THE OCEANIC CRUST COSTA RICA RIFT.	C-7
K. Becker, R. P. Von Herzen, T. J. G. Francis, R. N. Anderson, J. Honnorez, A. C. Adamson, J. C. Alt, R. Emmermann, P. D. Kempton, H. Kinoshita, C. Laverne, M. J. Mottl and R. L. Newmark	
RESIDUAL GEOID ANOMALIES IN ATLANTIC OCEAN BASIN: RELATIONSHIP TO MANTLE PLUMES.	C-7
Carl Bowin and Geoffrey Thompson	
CHEMICAL OCEANOGRAPHY OF THE PERSIAN GULF.	C-8
Peter G. Brewer.	
A CLIMATIC FRESHENING OF THE DEEP NORTH ATLANTIC (NORTH OF 50°N) OVER THE PAST 20 YEARS.	C-8
P. G. Brewer, W. S. Broecker, W. J. Jenkins, P. B. Rhines, C. G. Rooth, J. H. Swift, T. Takahashi and R. T. Williams	

THE MIXING PROFILE OF ORGANIC CARBON AND SEVERAL RADIOISOTOPES IN ABYSSAL RED CLAY SEDIMENTS.	C-9
E. R. M. Druffel, P. M. Williams, H. D. Livingston and M. Koide	
ALKALI AND Sr ISOTOPE GEOCHEMISTRY OF WATERS COLLECTED FROM BASALTIC BASEMENT, DEEP SEA DRILLING PROJECT HOLE 504B, COSTA RICA RIFT.	C-9
S. R. Hart and M. J. Mottl	
GEOCHEMISTRY OF RARE-EARTH ELEMENTS IN BASALTS FROM THE WALVIS RIDGE: IMPLICATIONS FOR ITS ORIGIN AND EVOLUTION.	C-9
Susan E. Humphris and Geoffrey Thompson	
TRACING OCEAN CIRCULATION BY MEANS OF CHEMICAL AND RADIOCHEMICAL DATA.	C-10
William J. Jenkins	
THE SEDIMENTATION RATES OF TRACE ELEMENTS IN THE SARGASSO SEA MEASURED BY SEDIMENT TRAP.	C-10
T. D. Jickells, W. G. Deuser and A. H. Knap	
TROPOSPHERIC NO _x AND O ₃ BUDGETS IN THE EQUATORIAL PACIFIC.	C-11
S. C. Liu, M. McFarland, D. Kley, O. Zafiriou and B. Huebert	
PROGRAMMED PYROLYSIS OF ORGANIC MATTER FROM THERMALLY ALTERED CRETACEOUS BLACK SHALES.	C-11
K. E. Peters, J. K. Whelan, J. M. Hunt and M. E. Tarafa	
HYDROTHERMAL ACTIVITY IN THE TAG HYDROTHERMAL FIELD, MID-ATLANTIC RIDGE CREST AT 26°N.	C-11
P. A. Rona, G. Thompson, M. J. Mottl, J. A. Karson, W. J. Jenkins, D. Graham, M. Mallette, K. Von Damm and J. M. Edmond	
ALTERATION OF THE OCEANIC CRUST: IMPLICATIONS FOR GEOCHEMICAL CYCLES OF LITHIUM AND BORON.	C-12
W. E. Seyfried, D. R. Janecky and M. J. Mottl	
BASALT - SEAWATER INTERACTION.	C-12
Geoffrey Thompson	
MORPHOLOGY, MINERALOGY AND CHEMISTRY OF HYDROTHERMAL DEPOSITS FROM THE TAG AREA, 26°N MID-ATLANTIC RIDGE.	C-13
Geoffrey Thompson, Michael J. Mottl and Peter A. Rona	
PHOTOCHEMISTRY AND THE SEA-SURFACE MICROLAYER: NATURAL PROCESSES AND POTENTIAL AS A TECHNIQUE.	C-13
Oliver C. Zafiriou	

Radiochemistry

SEASONALITY IN THE FLUX OF NATURAL RADIONUCLIDES AND PLUTONIUM IN THE DEEP SARGASSO SEA.	C-14
Michael P. Bacon, Chih-An Huh, Alan P. Fleer and Werner G. Deuser	
²¹⁰ Po AND ²¹⁰ Pb DISTRIBUTIONS IN THE CENTRAL AND EASTERN INDIAN OCEAN.	C-14
J. K. Cochran, M. P. Bacon, S. Krishnaswami and K. K. Turekian	
NEUTRON ACTIVATION ANALYSIS OF RARE EARTH ELEMENTS IN SEAWATER.	C-14
Hein J. W. DeBaar	
DETERMINATION OF LEAD-210 AND POLONIUM-210 IN SEAWATER AND MARINE PARTICULATE MATTER.	C-15
Alan P. Fleer and Michael P. Bacon	
HELIUM ISOTOPIC VARIATIONS IN VOLCANIC ROCKS FROM LOIHI SEAMOUNT AND THE ISLAND OF HAWAII.	C-15
M. D. Kurz, W. J. Jenkins, S. R. Hart and D. Clague	
ANTHROPOGENIC TRITIUM IN SOUTH ATLANTIC BOTTOM WATER.	C-15
W. J. Jenkins, D. E. Lott, M. W. Pratt and R. D. Boëdureau	
IN SITU CHEMISORPTION OF TRANSURANICS FROM SEAWATER.	C-16
Don R. Mann, Lolita D. Surprenant and Susan Casso	
PLUTONIUM OXIDATION STATE DISTRIBUTIONS IN THE PACIFIC OCEAN DURING 1980-1981.	C-16
D. M. Nelson, A. E. Carey and V. T. Bowen	
THE DISTRIBUTION OF ^{239,240} Pu, ¹³⁷ Cs, AND ⁵⁵ Fe IN CONTINENTAL MARGIN SEDIMENTS: RELATION TO SEDIMENTARY REDOX ENVIRONMENT.	C-16
F. L. Sayles and H. D. Livingston	
MEASUREMENT OF CURIUM IN MARINE SAMPLES.	C-17
David L. Schneider and Hugh D. Livingston	
THE COAGULATION OF DISSOLVED ^{239,240} Pu IN ESTUARIES AS DETERMINED FROM A MIXING EXPERIMENT.	C-17
Glen T. Shen, Edward R. Sholkovitz and Don R. Mann	

REDOX-RELATED GEOCHEMISTRY IN LAKES - ALKALI METALS, ALKALINE EARTH ELEMENTS AND ^{137}Cs .		C-17
Edward R. Sholkovitz		
THE PORE WATER CHEMISTRY OF $^{239,240}\text{Pu}$ AND ^{137}Cs IN SEDIMENTS OF BUZZARDS BAY, MASSACHUSETTS.		C-18
E. R. Sholkovitz and D. R. Mann		
TRIPOD MODIFICATION OF A SPHINCTER CORER: CONSTRUCTION, OPERATION, CORE EXTRUSION AND SAMPLING EFFICIENCY.		C-18
J. C. Burke, R. E. Hamblin and S. A. Casso		

Geology

NEOGENE GEOCHRONOLOGY AND CHRONOSTRATIGRAPHY.	GG-3
William A. Berggren, Dennis V. Kent and John A. Van Couvering	
MORPHOLOGY, SEDIMENTS AND LATE QUATERNARY HISTORY OF THE EAST CHINA SEA.	GG-3
Jorge Butenko, Ye Yincin and John D. Milliman	
MAGNETIC CHRON 12R AT FLAGSTAFF RIM, WYOMING.	GG-3
Charles R. Denham	
TECTONICS AT THE INTERSECTION OF THE EAST PACIFIC RISE WITH THE	GG-4
TAMAYO TRANSFORM FAULT.	
D.G. Gallo, W.S.F. Kidd, P.J. Fox, J.A. Karson, K. Macdonald,	
K. Crane, P. Choukroune, M. Seguret, R. Moody and K. Kastens	
SEDIMENTATION UNDER DEEP SEA-STORMS.	GG-4
Charles D. Hollister and Ian N. McCave	
ABYSSAL TELECONNECTIONS II. INITIATION OF ANTARCTIC BOTTOM WATER FLOW	GG-4
IN THE SOUTHWESTERN ATLANTIC.	
David A. Johnson	
THE VEMA CHANNEL: PHYSIOGRAPHY, STRUCTURE AND SEDIMENT-CURRENT INTERACTIONS. . .	GG-5
David A. Johnson	
DISPLACED ANTARCTIC DIATOMS IN VEMA CHANNEL SEDIMENTS: LATE PLEISTOCENE/HOLOCENE	
FLUCTUATIONS IN AABW FLOW.	GG-5
Glenn A. Jones and David A. Johnson	
GEOLOGIC AND SEISMIC VELOCITY STRUCTURE OF THE CRUST/MANTLE TRANSITION IN THE . .	GG-6
BAY OF ISLANDS OPHIOLITE COMPLEX.	
Jeffrey A. Karson, John A. Collins and J.F. Casey	
GEOLOGY OF THE OCEANOGRAPHER TRANSFORM: THE RIDGE-TRANSFORM INTERSECTION. . . .	GG-6
J.A. Karson, P.J. Fox, K. Crane, H. Sloane, E. Bonatti, W. Kidd, J. Stroup,	
D. Fornari, D. Elthon, P. Hamlyn, J. Casey, D. Gallo, D. Needham and R. Sartori	
TECTONICS OF RIDGE-TRANSFORM INTERSECTIONS AT THE KANE FRACTURE ZONE.	GG-7
J.A. Karson and H.J.B. Dick	
HIGH RESOLUTION ISOTOPIC STUDY OF THE LATEST DEGLACIATION FROM BERMUDA RISE CORES.	GG-7
L. D. Keigwin, Bruce H. Corliss, Ellen M. Druffel and E. P. Laine	
A CLOSE LOOK AT THE LATE MIOCENE CARBON ISOTOPIC SHIFT AT DSDP SITE 158.	GG-8
L. D. Keigwin, S.V. Margolis and B.U. Haq	
OLIGOCENE CALIBRATION OF THE MAGNETIC POLARITY TIMESCALE.	GG-8
Donald R. Prothero, Charles R. Denham and Harlow G. Farmer	
THE RED SEA.	GG-8
David A. Ross	
CENOZOIC REGIONAL EROSION OF THE ABYSSAL SEAFLOOR OFF SOUTH AFRICA.	GG-9
Brian E. Tucholke and Robert W. Embley	
ABYSSAL CURRENT CHARACTER DETERMINED FROM SEDIMENT BEDFORMS ON THE	GG-9
NOVA SCOTIAN CONTINENTAL RISE.	
B.E. Tucholke and C.D. Hollister	

Geophysics

DEPTH OF PRINCIPAL MASS ANOMALIES CONTRIBUTING TO THE EARTH'S GEOIDAL	GG-9
UNDULATIONS AND GRAVITY ANOMALIES.	
Carl Bowin	
GLOBAL GRAVITY MAPS AND THE STRUCTURE OF THE EARTH.	GG-10
Carl Bowin	
GRAVITY, TOPOGRAPHY AND CRUSTAL EVOLUTION OF VENUS.	GG-11
Carl Bowin	
STRUCTURE AND VARIABILITY OF OCEANIC LITHOSPHERE ON THE FLANKS OF THE	GG-11
EAST PACIFIC RISE BETWEEN 11° and 13°W.	
Steven R. Bratt and G.M. Purdy	
SEISMIC REFLECTION PROFILING WITH A DEEP-TOWED HYDROPHONE ARRAY.	GG-11
John A. Collins, Donald E. Koelsch, G. M. Purdy	
MODELING PALEOMAGNETIC REMANENCE.	GG-12
Charles R. Denham and Loren Shure	
THE AMERICA-ANTARCTIC RIDGE.	GG-12
Lawrence A. Lawver and Henry J.B. Dick	

HYPERBOLIC STACKING OF WIDE-ANGLE REFLECTIONS FROM DEEP-SEA SEDIMENTS.	GG-12
Kristin Rohr	
THE APPLICATION OF TAU-P MAPPING TO LOW SIGNAL TO NOISE RATIO WIDE-ANGLE REFLECTION DATA.	GG-13
K. Rohr	
VARIATION OF THE AMPLITUDES OF REFLECTORS FROM OCEANIC BASEMENT.	GG-13
Kristin Rohr	
DECONVOLUTION OF SEAFLOOR TOPOGRAPHY.	GG-13
Hans Schouten and Charles R. Denham	
ESTIMATING THE ROUGHNESS OF SEAFLOOR.	GG-13
Hans Schouten and Charles R. Denham	
FINITE DIFFERENCE SEISMOGRAMS FOR Laterally Varying Marine Models.	GG-14
Ralph A. Stephen	
SYNTHETIC VERTICAL SEISMIC PROFILES BY THE METHOD OF FINITE DIFFERENCES.	GG-14
Ralph A. Stephen	
BOREHOLE SEISMIC EXPERIMENTS AND THE STRUCTURE OF UPPER OCEANIC CRUST.	GG-14
R.A. Stephen	
TRAVEL-TIME ANALYSIS OF BOREHOLE SEISMIC DATA.	GG-15
R.A. Stephen and A.J. Harding	
CRUSTAL STRUCTURE IN THE OROZCO TRANSFORM ZONE.	GG-15
A.M. Trehu and G.M. Purdy	
CRUSTAL VELOCITY STRUCTURE ON THE FLANKS OF THE MID-ATLANTIC RIDGE AT 24°N.	GG-16
Robert S. White and G. M. Purdy	

Marine Chemistry

REDUCED ADVECTION INTO ATLANTIC OCEAN DEEP EASTERN BASINS DURING LAST GLACIATION MAXIMUM.	GG-16
Bill Curry and G. P. Lohmann	

Plate Tectonics

OVERLAPPING SPREADING CENTERS ON EAST PACIFIC RISE.	GG-16
Hans Schouten and Kim Klitgord	

Paleontology

CENOZOIC GEOCHRONOLOGY.	GG-17
William A. Berggren, Dennis V. Kent, John J. Flynn and John A. Van Courvering	
LATE TERTIARY/QUATERNARY MAGNETOSTRATIGRAPHY AND BIOSTRATIGRAPHY OF THE VEMA CHANNEL SEDIMENTS.	GG-17
David A. Johnson, Michael T. Ledbetter, Eric Tappa and Robert Thunell	
HIGH-RESOLUTION STRATIGRAPHY IN LATE PLEISTOCENE-HOLOCENE SEDIMENTS OF THE VEMA CHANNEL.	GG-17
Glenn A. Jones, David A. Johnson and William B. Curry	
EVIDENCE OF PUNCTUATED GRADUALISM IN THE <i>Globorotalia tumida</i> LINEAGE (PLANKTONIC FORAMINIFERA).	GG-18
Bjorn A. Malmgren, W.A. Berggren and G.P. Lohmann	
RADIOLARIA.	GG-18
Kozo Takahashi	

Petrology

SYSTEMATICS OF MODAL PHENOCRUST ASSEMBLAGES IN SUBMARINE BASALTS: PETROLOGIC IMPLICATIONS.	GG-19
W.B. Bryan	
RECONSTRUCTION OF THE GEOMETRY OF ACCRETION DURING FORMATION OF THE BAY OF ISLANDS OPHIOLITE COMPLEX.	GG-20
J.F. Casey, J.A. Karson, D. Elthon, Elric Rosencrantz and Marchall Titus	
MINERALOGIC STUDIES OF THE RESIDUES OF MANTLE MELTING: ABYSSAL AND ALPINE-TYPE PERIDOTITES.	GG-20
H.J.B. Dick and R.L. Fisher	

FRACTIONATION OF PYROXENE-PHYRIC MORB AT LOW PRESSURE: AN EXPERIMENTAL STUDY. . .	GG-21
T.L. Grove and W.B. Bryan	
VARIATIONS IN STRUCTURE AND PETROLOGY IN THE COASTAL COMPLEX, NEWFOUNDLAND: . .	GG-21
THE ANATOMY OF AN OCEANIC FRACTURE ZONE.	
J.A. Karson	
GEOCHEMISTRY, MINERALOGY AND PETROGENESIS OF LAVAS ERUPTED ALONG THE SOUTHWEST	GG-21
INDIAN RIDGE BETWEEN THE BOUVET TRIPLE JUNCTION AND 11 DEGREES EAST.	
A.P. leRoex, H.J.B. Dick, A.J. Erlank, F.A. Frey and S.R. Hart	

Sedimentology QE

BEACH CHANGES ON COASTS WITH DIFFERENT WAVE CLIMATES.	GG-22
David G. Aubrey	
UPDRIFT MIGRATION OF TIDAL INLETS: A NATURAL RESPONSE TO WAVE, TIDE	GG-23
AND STORM FORCING.	
David G. Aubrey, Paul E. Speer and Eric Ruder	
SEDIMENTATION OF PARTICULATE MANGANESE IN THE DEEP OCEAN.	GG-23
Susumu Honjo, Jonathan J. Cole, Alan P. Fleer and Cynthia H. Pilska	
SEASONALITY IN THE FLUXES OF SUGARS, AMINO ACIDS AND AMINO SUGARS	GG-23
TO DEEP OCEAN. PART II. PANAMA BASIN.	
Venugopalan Ittekkot, Egon T. Degens and Susumu Honjo	
LATE CENOZOIC TURBIDITE AND CONTOURITE DEPOSITION IN THE SOUTHERN BRAZIL BASIN.	GG-24
David A. Johnson and Kenneth A. Rasmussen	
ANALYSIS OF A LONGITUDINAL RIPPLE FROM THE NOVA SCOTIAN CONTINENTAL RISE. . . .	GG-24
I.N. McCave, C.D. Hollister, D.J. DeMaster, C.A. Nittrouer,	
A.J. Silva and J.Y. Yingst	
SEDIMENT DISCHARGE FROM THE INDUS RIVER TO THE OCEAN: PAST, PRESENT AND FUTURE.	GG-25
John D. Milliman, G.S. Quraishee and M.A.A. Beg	
TIDAL PHASE CONTROL OF SEDIMENT DISCHARGE FROM THE YANGTZE RIVER.	GG-25
J.D. Milliman, Ya Hsueh, Dun-Xin Hu, David J. Pashinski,	
Huang-ting Shen, Zuo-sheng Yang and Peter Hacker	
TRANSFER OF PARTICULATE ORGANIC CARBON AND NITROGEN FROM THE YANGTZE RIVER . . .	GG-25
TO THE OCEAN.	
John D. Milliman, Xie Qinchon and Yang Zuosheng	
FLUX OF SUSPENDED SEDIMENT IN THE CHANG JIANG (YANGTZE) ESTUARY.	GG-25
J.D. Milliman, Z.-s. Yang and R.H. Meade	
PHAEODARIAN SKELETONS: ROLE OF SILICA TRANSPORT TO THE DEEP SEA AND DISSOLUTION.	GG-26
Kozo Takahashi, David C. Hurd and Susumu Honjo	
LATE QUATERNARY SEDIMENTATION ON THE SOUTHERN SOHM ABYSSAL PLAIN.	GG-26
Gustavs Vilks, D.E. Buckley and L.D. Keigwin, Jr.	

Technical Reports

COASTAL SEDIMENT TRANSPORT POPPONSETT BEACH, MA.	GG-26
David G. Aubrey and Margaret R. Goud	
SEDIMENT TRANSPORT IN A TIDAL INLET.	GG-27
David G. Aubrey and Paul E. Speer	

Wave Mechanics QC

FIELD INTERCOMPARISON OF NEARSHORE DIRECTIONAL WAVE SENSORS.	GG-27
W.G. Grosskopf, D.G. Aubrey and M.G. Mattie	

Acoustic Signal Processing GC

CONVERGENCE ANALYSIS OF LMS FILTERS WITH UNCORRELATED GAUSSIAN DATA.	OE-3
A. Feuer and E. Weinstein	
ANGLE-INDEPENDENT DOPPLER MEASUREMENT.	OE-3
Nadav Levanon and Ehud Weinstein	
DELAY AND DOPPLER ESTIMATION BY TIME-SPACE PARTITION OF THE ARRAY DATA.	OE-3
Ehud Weinstein and Doron Kletter	
FUNDAMENTAL LIMITATIONS IN PASSIVE TIME-DELAY ESTIMATION--PART I:	OE-3
NARROW-BAND SYSTEMS.	
Anthony J. Weiss and Ehud Weinstein	
FUNDAMENTAL LIMITATIONS IN PASSIVE TIME DELAY ESTIMATION--PART II:	OE-4
WIDEBAND SYSTEMS.	
Ehud Weinstein and Anthony J. Weiss	

Coastal Dynamics

RIPPLED SCOUR DEPRESSIONS ON THE CONTINENTAL SHELF OFF CENTRAL CALIFORNIA.	OE-4
D.A. Cacchione, D.E. Drake, W.D. Grant and G.B. Tate	
VARIABILITY OF SEA-FLOOR ROUGHNESS WITHIN THE COASTAL OCEAN DYNAMICS	OE-5
EXPERIMENT (CODE) REGION.	
David A. Cacchione, David E. Drake, William D. Grant,	
Albert J. Williams III and George B. Tate	
BOTTOM STRESS ESTIMATES AND THEIR PREDICTION ON THE NORTHERN CALIFORNIA	OE-5
CONTINENTAL SHELF DURING CODE-1: THE IMPORTANCE OF WAVE-CURRENT INTERACTION.	
W.D. Grant, A.J. Williams and S.M. Glenn	
CONTINENTAL SHELF BOTTOM BOUNDARY LAYER MODEL VOLUME I:	OE-5
THEORETICAL MODEL DEVELOPMENT.	
William D. Grant and Scott M. Glenn	
CONTINENTAL SHELF BOTTOM BOUNDARY LAYER MODEL VOLUME II:	OE-6
MODEL/DATA COMPARISON.	
W.D. Grant and S.M. Glenn	
HIGH FREQUENCY BOTTOM STRESS VARIABILITY AND ITS PREDICTION IN THE CODE REGION.	OE-6
W.D. Grant, A.J. Williams, 3rd, S.M. Glenn, D.A. Cacchione and D.E. Drake	

Computer Science QA

FINDIF: A SOFTWARE PACKAGE TO CREATE SYNTHETIC SEISMOGRAMS BY FINITE DIFFERENCES.	OE-7
Mary M. Hunt, Lee Gove and Ralph A. Stephen	

Geology GE

THE EAST PACIFIC RISE NEAR 21°N, 13°N AND 20°S: INFERENCES FOR	OE-7
ALONG-STRIKE VARIABILITY OF AXIAL PROCESSES OF THE MID-OCEAN RIDGE.	
Jean Francheteau and Robert D. Ballard	
GEOLOGIC PROCESSES OF THE MID-OCEAN RIDGE AND THEIR RELATIONSHIP TO	OE-7
SULFIDE DEPOSITION.	
Robert D. Ballard and Jean Francheteau	
INTENSE HYDROTHERMAL ACTIVITY AT THE AXIS OF THE EAST PACIFIC RISE NEAR 13°N:	OE-8
SUBMERSIBLE WITNESSES THE GROWTH OF SULFIDE CHIMNEY.	
R. Hekinian, J. Francheteau, V. Renard, R.D. Ballard,	
P. Choukroune, J.L. Cheminee, F. Albarede, J.F. Minster,	
J.C. Marty, J. Boulegue and J.L. Charlou	
PROJECT CYATHERM: DETAILED INVESTIGATION OF HYDROTHERMAL ACTIVITY AT	OE-8
12°48'N ON THE EAST PACIFIC RISE.	
R.D. Ballard	

Instrumentation

A CCD CHIRP-Z FFT DOPPLER SIGNAL PROCESSOR FOR LASER VELOCIMETRY.	OE-8
Y.C. Agrawal	

DIRECTIONAL LASER VELOCIMETRY WITHOUT FREQUENCY BIASING - PART II.	OE-9
Y.C. Agrawal and J.B. Riley	
CTD LOWERING MECHANICS.	OE-9
H.O. Berteaux and R.G. Walden	
EQUIPMENT LOWERING MECHANICS.	OE-9
H.O. Berteaux	
EXPERIMENTAL EVALUATION OF CTD PACKAGE HYDRODYNAMIC BEHAVIOR AND RECOMMENDATIONS FOR IMPROVED LOWERING TECHNIQUES.	OE-10
H.O. Berteaux, R.G. Walden, P.R. Clay and R.E. McDevitt	
AIR DEPLOYED OCEANOGRAPHIC MOORING - AB1034.	OE-10
L.W. Bonde, D.B. Dillon, E.J. Softley, R. Walden, H. Berteaux and T. Popp	
A COHERENT APPROACH TO INSTRUMENT INTERCOMMUNICATION AND TESTING USING SAIL.	OE-10
Albert M. Bradley and William E. Terry, Jr.	
INTEGRATED COMMUNICATIONS IN BUOY SYSTEMS.	OE-11
Ed Mellinger and Al Bradley	
THE LOTUS DISCUS MOORING.	OE-11
Peter R. Clay	
SEISMIC REFLECTION PROFILING WITH DEEP TOWED HYDROPHONE ARRAY.	OE-11
J. Collins, D. Koelsch, and G. Purdy	
DIRECT OPTICAL ASSESSMENT OF MACROSCOPIC AGGREGATES IN THE DEEP OCEAN.	OE-11
Susumu Honjo, Kenneth W. Doherty, Yogesh C. Agrawal and Vernon L. Asper	
AN EVENT RECORDING OCEAN BOTTOM SEISMOGRAPH.	OE-12
K.R. Peal and R.E. Kirk	
A DEEP OCEAN AMBIENT NOISE SPECTRA RECORDING INSTRUMENT.	OE-12
K. Prada, D. Koelsch, K. Von der Heydt, E. Mellinger, S. Liberatore and F. Schuler	
FREE DRIFTING RELAYS BUOY SYSTEM.	OE-12
R.G. Walden and H.O. Berteaux	
HEBBLE SEA DUCT ANNUAL PROJECT REPORT #3 YEAR TWO, 1982-1983.	OE-12
Clifford L. Winget	

Ocean Acoustics

FLUCTUATIONS OF SOUND PROPAGATING VERTICALLY THROUGH THE OCEAN.	OE-13
Yves Desaubies	
SHALLOW WATER WAVEGUIDE CHARACTERIZATION USING THE HANKEL TRANSFORM.	OE-13
George V. Frisk and James F. Lynch	
ON THE USE OF FOCUSED HORIZONTAL ARRAYS AS MODE SEPARATION AND SOURCE LOCATION DEVICES IN OCEAN ACOUSTICS. PART I: THEORY.	OE-13
James F. Lynch	
SOME SIMPLE EXPRESSIONS FOR THE BEAMFORMING PROPERTIES OF FOCUSED HIGH RESOLUTION CIRCULAR ARRAYS, WITH APPLICATIONS TO REFOCUSING SYSTEMS.	OE-13
James F. Lynch	
A HYBRID NUMERICAL/ANALYTIC TECHNIQUE FOR THE COMPUTATION OF WAVE FIELDS IN STRATIFIED MEDIA BASED ON THE HANKEL TRANSFORM.	OE-14
Douglas R. Mook, George V. Frisk and Alan V. Oppenheim	
MEASUREMENTS OF GULF STREAM MEANDERING AND EVIDENCE OF SEASONAL THERMOCLINE DEVELOPMENT USING LONG RANGE ACOUSTIC TRANSMISSION.	OE-14
J.L. Spiesberger, T.G. Birdsall, K. Metzger, R.R. Knox, C.W. Spofford and R.C. Spindel	
PERTURBATIONS IN TRAVEL TIME AND RAY GEOMETRY DUE TO MESOSCALE DISTURBANCES: A COMPARISON OF EXACT AND APPROXIMATE CALCULATIONS.	OE-14
John L. Spiesberger and Peter F. Worcester	
EDDIES AND ACOUSTICS.	OE-15
R.C. Spindel and Y.J.F. Desaubies	

Submersibles

DSRV ALVIN WEIGHT AND STABILITY.	OE-15
Arnold G. Sharp	

Coastal Dynamics GC

STRUCTURE OF THE CHANGJIANG RIVER PLUME IN THE EAST CHINA SEA DURING JUNE 1980. . .	PO-3
Robert C. Beardsley, Richard Limeburner, Dun Xin Hu, Ken Tang Le, Glenn A. Cannon and David J. Pashinski	
LOW-FREQUENCY CURRENT AND BOTTOM-PRESSURE VARIABILITY.	PO-3
Kenneth H. Brink, Bruce A. Magnell and Marlene A. Noble	
OBSERVATIONS OF THE COASTAL UPWELLING REGION NEAR 34°30'N OFF CALIFORNIA:	PO-3
SPRING 1981. Kenneth H. Brink, David W. Stuart and John C. Van Leer	
LONG-TERM OBSERVATIONS ON THE SOUTHERN FLANK OF GEORGES BANK: SEASONAL	PO-4
CYCLE OF CURRENT, TEMPERATURE, STRATIFICATION AND WIND STRESS. Bradford Butman and Robert C. Beardsley	
THE PHYSICAL OCEANOGRAPHY OF GEORGES BANK: INTRODUCTION AND SUMMARY.	PO-4
Bradford Butman and Robert C. Beardsley	
THE SEASONAL MEAN CIRCULATION ON GEORGES BANK: OBSERVATION AND THEORY.	PO-4
Bradford Butman, John W. Loder and Robert C. Beardsley	
AN UPPER BOUND FOR THE TIDALLY RECTIFIED CURRENT AT ONE LOCATION ON THE	PO-5
SOUTHERN FLANK OF GEORGES BANK. Bradford Butman, Marlene A. Noble, David C. Chapman and Robert C. Beardsley	
DISPERSAL BY RANDOMLY VARYING CURRENTS.	PO-5
Gabriel T. Csanady	
THE "INSULATING" EFFECT OF A STEEP CONTINENTAL SLOPE.	PO-6
Gabriel T. Csanady and Ping Tung Shaw	
THE HYDROGRAPHIC STRUCTURE AND VARIABILITY OF GEORGES BANK.	PO-6
Charles N. Flagg	
WATER MASSES AND CIRCULATION IN THE EAST CHINA SEA.	PO-7
Richard Limeburner and Robert C. Beardsley	

Instrumentation

POPUP: A PROTOTYPE BOTTOM-MOORED LONG-TERM CURRENT PROFILER.	PO-7
Arthur D. Voorhis and Albert M. Bradley	

Laboratory Models

LABORATORY MODELS OF BAY-TYPE CONTINENTAL SHELVES IN THE WINTER.	PO-7
Takashige Sugimoto and John A. Whitehead, Jr.	
DISLOCATIONS IN CONVECTION AND THE ONSET OF CHAOS.	PO-8
John A. Whitehead	
A LABORATORY STUDY OF GYRES AND UPLIFT NEAR THE STRAIT OF GIBRALTAR.	PO-8
John A. Whitehead, Jr.	

Large Scale Oceanic Heat and Water Properties

WARM WATER TO COLD WATER CONVERSION IN THE NORTHERN NORTH ATLANTIC OCEAN.	PO-8
Michael S. McCartney and Lynne D. Talley	
THE STRUCTURE, DYNAMICS AND ORIGIN OF A SMALL SCALE LENS OF WATER IN THE	PO-9
WESTERN NORTH ATLANTIC THERMOCLINE. Stephen C. Riser, W. Brechner Owens, H. Thomas Rossby and Curtis C. Ebbesmeyer	

Ocean Circulation GC

LIMITING PROCESSES THAT DETERMINE BASIC FEATURES OF THE CIRCULATION IN THE . . .	PO-9
MEDITERRANEAN SEA. Harry L. Bryden and Henry M. Stommel	
CURRENT VARIABILITY NEAR THE SOUTHEAST NEWFOUNDLAND RIDGE.	PO-9
Nick P. Fofonoff and Ross M. Hendry	

EVIDENCE FOR BAROCLINIC INSTABILITY IN THE GULF STREAM RECIRCULATION.	PO-10
Nelson G. Hogg	
THE DYNAMICS OF THE OCEANIC GENERAL CIRCULATION: THE POTENTIAL VORTICITY FIELD.	PO-10
William R. Holland, Thomas Keffer and Peter B. Rhines	
THE VENTILATION OF THE WORLD'S OCEANS: MAPS OF THE POTENTIAL VORTICITY FIELD.	PO-11
Thomas Keffer	
THE DENSITY JUMP ACROSS LITTLE BAHAMA BANK.	PO-11
James R. Luyten and Henry M. Stommel	
UPSTREAM EFFECTS OF THE GULF STREAM ON THE STRUCTURE OF THE MID-OCEAN THERMOCLINE.	PO-11
James R. Luyten and Henry M. Stommel	
A SYNOPTIC AND STATISTICAL DESCRIPTION OF THE GULF STREAM AND SUBTROPICAL GYRE	PO-11
USING SOFAR FLOATS.	
W. Brechner Owens	
DRIFTING DERELICTS IN THE NORTH ATLANTIC 1883-1902.	PO-12
Philip L. Richardson	
AVERAGE SEASONAL VARIATION OF THE ATLANTIC NORTH EQUATORIAL COUNTERCURRENT	PO-12
FROM SHIP DRIFT DATA.	
Philip L. Richardson and Theresa K. McKee	
OBSERVATIONS OF HORIZONTAL VELOCITIES AND VERTICAL DISPLACEMENTS IN THE EQUATORIAL	PO-12
PACIFIC OCEAN ASSOCIATED WITH THE EARLY STAGES OF THE 1982/83 EL NINO.	
John M. Toole and Mark D. Borges	
WIND FORCED VARIABILITY IN THE CENTRAL EQUATORIAL PACIFIC DURING PEQUOD.	PO-12
Arthur D. Voorhis, James R. Luyten, Gerald J. Needell and	
John H. Thomson	
SOME PRELIMINARY RESULTS CONCERNING DEEP NORTHERN-BOUNDARY CURRENTS IN THE	PO-13
NORTH PACIFIC.	
Bruce A. Warren and W. Brechner Owens	

Rings, Eddies

THE ENERGY BALANCE DURING THE POLYMODE LOCAL DYNAMICS EXPERIMENT INTENSIVE	PO-13
CTD SURVEYS.	
Nancy A. Bray, W. Brechner Owens and Bach-Lien Hua	
COMPARISON OF EDDIES OFF THE NORTH BRAZILIAN AND SOMALI COASTS.	PO-13
John G. Bruce	
ENERGETIC EDDIES AT THE NORTHERN EDGE OF THE ANTARCTIC CIRCUMPOLAR CURRENT	PO-14
IN THE SOUTHWEST PACIFIC.	
Harry L. Bryden and Ron A. Heath	
VELOCITY AND HYDROGRAPHIC STRUCTURE OF A GULF STREAM WARM CORE RING.	PO-14
Terrence M. Joyce	
RAPID EVOLUTION OF A GULF STREAM WARM-CORE RING.	PO-14
Terrence M. Joyce, Richard H. Backus, Karen S. Baker,	
Patricia L. Blackwelder, Otis B. Brown, Timothy J. Cowles,	
Robert H. Evans, Greta A. Fryxell, David G. Mountain,	
Donald B. Olson, Ronald J. Schlitz, Raymond W. Schmitt,	
Peter C. Smith, Raymond C. Smith and Peter H. Wiebe	
AN UPPER OCEAN CURRENT JET AND INTERNAL WAVES IN A GULF STREAM WARM CORE RING.	PO-15
Terrence M. Joyce and Marvel C. Stalcup	
THE SPATIAL AND TEMPORAL EVOLUTION OF A CLUSTER OF SOFAR FLOATS IN THE	PO-15
POLYMODE LOCAL DYNAMICS EXPERIMENT (LDE).	
H. Thomas Rossby, James F. Price and Douglas C. Webb	
ABYSSAL EDDY KINETIC ENERGY IN THE NORTH ATLANTIC.	PO-16
William J. Schmitz, Jr.	
ABYSSAL EDDY KINETIC ENERGY LEVELS IN THE WESTERN NORTH PACIFIC.	PO-16
William J. Schmitz, Jr.	
OBSERVATIONS OF THE VERTICAL STRUCTURE OF THE EDDY FIELD IN THE KUROSHIO EXTENSION.	PO-16
William J. Schmitz, Jr.	

Theoretical Models

THE GENERATION OF BAROTROPIC EDGE WAVES BY DEEP-SEA INTERNAL WAVES.	PO-17
David C. Chapman	
WARM WATER MASS FORMATION.	PO-17
Gabriel T. Csanady	

THE NONLINEAR SPIN-UP OF A STRATIFIED OCEAN.	PO-17
William K. Dewar, Peter B. Rhines and William R. Young	
TRACER DISPERSAL BY MID-OCEAN MESOSCALE EDDIES. I: ENSEMBLE STATISTICS. . . .	PO-18
Dale B. Haidvogel and Thomas Keffer	
A COMPARISON OF THE FORMULATION FOR EDDY DIFFUSION IN TWO ONE-DIMENSIONAL STRATIFICATION MODELS. . . .	PO-18
Brian Henderson-Sellers, M. H. McCormick and Donald Scavia	
EASTERN BOUNDARY VENTILATION AND THE STRUCTURE OF THE THERMOCLINE.	PO-18
Joseph Pedlosky	
THE EQUATIONS FOR GEOSTROPHIC MOTION IN THE OCEAN.	PO-19
Joseph Pedlosky	
THE GROWTH AND DECAY OF FINITE-AMPLITUDE BAROCLINIC WAVES.	PO-19
Joseph Pedlosky	
ON THE RELATIVE IMPORTANCE OF VENTILATION AND MIXING OF POTENTIAL VORTICITY IN MID-OCEAN GYRES.	PO-19
Joseph Pedlosky	
VENTILATION, POTENTIAL-VORTICITY HOMOGENIZATION AND THE STRUCTURE OF THE OCEAN CIRCULATION.	PO-20
Joseph Pedlosky and William R. Young	

Turbulence and Mixing, Fine Scale Processes

ESTIMATES OF THE RELATIVE ROLES OF DIAPYCNAL, EPIPYCNAL AND DOUBLE-DIFFUSIVE MIXING IN ANTARCTIC BOTTOM WATER IN THE NORTH ATLANTIC. . .	PO-20
Trevor J. McDougall and John A. Whitehead	
THE CHARACTERISTICS OF SALT FINGERS IN A VARIETY OF FLUID SYSTEMS, INCLUDING STELLAR INTERIORS, LIQUID METALS, OCEANS, AND MAGMAS. . .	PO-20
Raymond W. Schmitt	
FINESCALE VELOCITY-DENSITY CHARACTERISTICS AND RICHARDSON NUMBER STATISTICS OF THE EASTERN EQUATORIAL PACIFIC. . . .	PO-21
John M. Toole and Stanley P. Hayes	

Upper Ocean Surface Layer Processes and Meteorology

PRELIMINARY RESULTS FROM THE LONG-TERM UPPER-OCEAN STUDY (LOTUS).	PO-21
Melbourne G. Briscoe and Robert A. Weller	
INTERCOMPARISON OF AIRCRAFT AND SURFACE BUOY METEOROLOGICAL DATA DURING CODE-1.	PO-21
Carl A. Friehe, Robert C. Beardsley, Clinton D. Winant and Jerome P. Dean	
DIURNAL CYCLING: OBSERVATIONS AND MODELS OF THE UPPER OCEAN RESPONSE TO DIURNAL HEATING, COOLING, AND WIND MIXING. . . .	PO-22
James F. Price, Robert A. Weller and Robert Pinkel	
OBSERVATIONS OF THREE DIMENSIONAL FLOW IN THE SURFACE MIXED LAYER OF THE OCEAN.	PO-22
Robert A. Weller, Jerome P. Dean, James F. Price, Erika A. Francis, John Marra and David C. Boardman	

Technical Reports

NANTUCKET SHOALS FLUX EXPERIMENT (NSFE79) PART 2: MOORED ARRAY DATA REPORT. . .	PO-23
Robert C. Beardsley, Carol A. Mills, John A. Vermersch, Jr., Wendell S. Brown, Neal R. Pettigrew, James D. Irish, Steven R. Ramp, Ronald J. Schlitz and Bradford Butman	
MEAN EULERIAN SUBSURFACE CURRENTS MEASURED IN THE GULF OF MAINE AND ADJACENT SCOTIAN AND NEW ENGLAND SHELF AND SLOPE REGIONS, 1974-1980. . .	PO-23
Robert C. Beardsley, Carol A. Mills, Peter C. Smith and Bradford Butman	
HYDROGRAPHIC STATION DATA OF FIVE SURVEYS OF THE BETA-TRIANGLE IN THE EASTERN NORTH ATLANTIC, 1978-1981.	PO-23
David W. Behringer, George P. Knapp, Robert J. Stanley and Henry M. Stommel	
LONG-TERM UPPER-OCEAN STUDY (LOTUS) AT 34°N, 70°W: METEOROLOGICAL SENSORS, DATA, AND HEAT FLUXES FOR MAY-OCTOBER 1982 (LOTUS-3 AND LOTUS-4). . . .	PO-23
Clara Deser, Robert A. Weller and Melbourne G. Briscoe	
A COMPILATION OF MOORED INSTRUMENT DATA AND ASSOCIATED OCEANOGRAPHIC OBSERVATIONS FROM THE VEMA CHANNEL, 1979 - 1981, VOLUME XXXII.	PO-24
Ellen Levy	

A COMPILATION OF MOORED CURRENT METER DATA FROM THE WESTERN NORTH PACIFIC, VOLUME XXXI, 1980-1982.	PO-24
Ellen Levy and Susan A. Tarbell	
CODE-1: MOORED ARRAY AND LARGE-SCALE DATA REPORT.	PO-24
Leslie K. Rosenfeld, editor	
THE CAMS INTERACTIVE ATLAS PACKAGE: A COMPUTERIZED PROGRAM FOR THE ARCHIVING AND PRESENTATION OF OCEANOGRAPHIC DATA.	PO-25
Thomas A. Sgouros and Thomas Keffer	
USNS BARTLETT CRUISE 40-B, DATA REPORT.	PO-25
Marvel C. Stalcup, Terrence M. Joyce and R. Lorraine Barbour	
THE LONG-TERM UPPER-OCEAN STUDY (LOTUS) CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT OCEANUS 119 - MAY 1982.	PO-25
Richard P. Trask and Melbourne G. Briscoe	
THE LONG-TERM UPPER-OCEAN STUDY (LOTUS) CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT OCEANUS 129 - OCTOBER 1982.	PO-26
Richard P. Trask and Melbourne G. Briscoe	
THE LONG-TERM UPPER-OCEAN STUDY (LOTUS) CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT ENDEAVOR 97 - April 1983.	PO-26
Richard P. Trask and Melbourne G. Briscoe	
FINAL TECHNICAL REPORT CONTRACT N00014-81-C-0010 1 OCTOBER 1980 - 30 SEPTEMBER 1982.	PO-26
John A. Whitehead, Jr.	

THE SOCIOLOGY OF PRODUCTION IN RURAL MALAY SOCIETY.	MP-3
Conner Bailey	
SMALL SCALE FISHERIES OF SAN MIGUEL BAY, PHILIPPINES: OCCUPATIONAL AND GEOGRAPHIC MOBILITY.	MP-3
Conner Bailey	
SMALL SCALE FISHERIES OF SAN MIGUEL BAY, PHILIPPINES: SOCIAL ASPECTS OF PRODUCTION AND MARKETING.	MP-3
Conner Bailey	
SYSTEMS OF SHARING AND PATTERNS OF OWNERSHIP.	MP-4
Conner Bailey and E.D. Villafuerte	
MANAGING AN OPEN ACCESS RESOURCE: THE CASE OF COASTAL FISHERIES.	MP-4
Conner Bailey	
DEVELOPING A U.S. RESEARCH STRATEGY FOR MARINE POLYMETALLIC SULFIDES.	MP-4
James M. Broadus and Robert E. Bowen	
EVALUATING THE ECONOMIC SIGNIFICANCE OF POLYMETALLIC SULFIDES DEPOSITS.	MP-5
James M. Broadus and Robert E. Bowen	
CONFLICT RESOLUTION IN THE ASSIGNMENT OF AREA ENTITLEMENTS FOR SEABED MINING.	MP-5
James M. Broadus and Porter Hoagland, III	
THE MONTEGO BAY CONVENTION OF 1982 AND THE CUSTOMARY LAW OF THE SEA: OBSERVATIONS, A FRAMEWORK AND A WARNING.	MP-5
John King Gamble, Jr.	
THE SIGNIFICANCE OF SIGNATURE TO THE 1982 MONTEGO BAY CONVENTION ON THE LAW OF THE SEA.	MP-5
John King Gamble, Jr.	
MULTIATTRIBUTE ANALYSIS AND THE DEFINITION OF OPTIMUM YIELD.	MP-6
Michael C. Healey	
INTER- AND INTRAPOPULATION VARIATION IN THE FECUNDITY OF CHINOOK SALMON (<i>Oncorhynchus tshawytscha</i>) AND ITS RELEVANCE TO LIFE HISTORY THEORY.	MP-6
Michael C. Healey	
VARIATION IN FECUNDITY OF LAKE WHITEFISH (<i>Coregonus clupeaformis</i>) FROM LESSER SLAVE AND UTIKUMA LAKES IN NORTHERN ALBERTA.	MP-7
Michael C. Healey and K. Dietz	
FISH PREDATION ON AQUATIC INSECTS.	MP-7
Michael C. Healey	
THE U.S. FISHING INDUSTRY AND REGULATORY REFORM.	MP-7
Timothy Hennessey	
HOW U.S. OCEAN POLICY IS MADE AND UNMADE.	MP-7
Robert W. Knecht	
OCEAN POLICY-MAKING IN A CHANGING WORLD.	MP-8
Robert W. Knecht and Bilianna Cicin-Sain	
ECONOMIC INCENTIVES FOR ENVIRONMENTAL PROTECTION: OCEAN DUMPING CHANGES.	MP-8
William L. Lahey	
THE CASE FOR OCEAN WASTE DISPOSAL.	MP-8
William Lahey and Michael Connor	
AMERICA AND THE MIDDLE EAST: HOLDING THE CENTRE.	MP-9
Linda B. Miller	
ENERGY AT THE SUMMIT.	MP-9
Linda B. Miller	
THROUGH A GLASS DARKLY: WESTERN EUROPE AND THE MIDDLE EAST.	MP-9
Linda B. Miller	
OIL, SHRIMP, MANGROVES: AN EVALUATION OF CONTINGENCY PLANNING FOR THE GULF OF GUAYAQUIL, ECUADOR.	MP-9
Ivon d'Almeida Pires Filho	
PROSPECTIVE MARITIME JURISDICTIONS IN THE POLAR SEAS.	MP-10
Richard L. Price and Marc E. Vaucher	
EFFECTIVE USE OF THE SEA OVERCOMING THE LAW OF THE SEA PROBLEMS.	MP-10
David A. Ross	
THE SHELFBREAK: SOME LEGAL ASPECTS.	MP-10
David A. Ross and K.O. Emery	
U.S.-MEXICO MARINE SCIENCE BILATERAL AGREEMENTS: WHAT SHOULD BE INVOLVED?	MP-11
David A. Ross	
INTERNATIONAL JURISDICTION ISSUES IN THE ARCTIC OCEAN.	MP-11
Kurt M. Shusterich	

ASSESSING THE OCEAN'S ROLES IN ANTILLEAN PREHISTORY.	MP-11
David R. Watters	
HIGHLAND HOUSE, BARBUDA: AN 18th CENTURY RETREAT.	MP-11
David R. Watters and Desmond V. Nicholson	
ALTERNATIVE REGIMES FOR FUTURE MINERAL RESOURCE DEVELOPMENT IN ANTARCTICA. . . .	MP-12
William E. Westermeyer	

GRADUATE STUDENTS	GS-1
AN IMPROVED LORAN-C DRIFTING BUOY AND DROGUE FOR COASTAL APPLICATIONS.	GS-3
William J. Burke	
INVERSE METHODS AND RESULTS FROM THE 1981 OCEAN ACOUSTIC TOMOGRAPHY EXPERIMENT.	GS-3
Bruce D. Cornuelle	
FORAMINIFERAL TRACE ELEMENTS: UPTAKE, DIAGENESIS, AND	GS-3
100 M.Y. PALEOCHEMICAL HISTORY.	
Margaret L. Delaney	
ATMOSPHERIC INTERACTIONS WITH GULF STREAM RINGS.	GS-4
William K. Dewar	
PROCESSING AND INVERSION OF ARCTIC OCEAN REFRACTION DATA.	GS-5
Gregory L. Duckworth	
A CONTINENTAL SHELF BOTTOM BOUNDARY LAYER MODEL: THE EFFECTS OF WAVES,	GS-5
CURRENTS, AND A MOVEABLE BED.	
Scott M. Glenn	
PURIFICATION AND CHARACTERIZATION OF THE HEPATIC MICROSOMAL MONOOXYGENASE . . .	GS-6
SYSTEM FROM THE COASTAL MARINE FISH <u>Stenotomus Chrysops</u> .	
Alan V. Klotz	
STABLE ISOTOPE GEOCHEMISTRY OF NITROGEN IN MARINE PARTICULATES.	GS-7
Susan M. Libes	
EFFECTS OF SIZE, AGE AND PHOTOPERIOD ON HYPOSMOREGULATION IN BROOK TROUT, . . .	GS-8
<u>Salvelinus fontinalis</u> .	
Stephen D. McCormick	
LATE PALEOGENE (EOCENE TO OLIGOCENE) PALEOCEANOGRAPHY OF THE NORTHERN	GS-8
NORTH ATLANTIC.	
Kenneth G. Miller	
THE NUMERICAL SYNTHESIS AND INVERSION OF ACOUSTIC FIELDS USING THE	GS-9
HANKEL TRANSFORM WITH APPLICATION TO THE ESTIMATION OF THE PLANE WAVE	
REFLECTION COEFFICIENT OF THE OCEAN BOTTOM.	
Douglas R. Mook	
FLOW AND SKIN FRICTION OVER NATURAL ROUGH BEDS.	GS-10
Christopher Paola	
A STUDY OF THE SEISMIC STRUCTURE OF UPPER OCEANIC CRUST USING	GS-10
WIDE-ANGLE REFLECTIONS.	
Kristin M. M. Rohr	
WAVE-INDUCED TURBULENT FLOW NEAR A ROUGH BED: IMPLICATIONS OF A	GS-11
TIME-VARYING EDDY VISCOSITY.	
John H. Trowbridge	
THE OCEANOGRAPHIC AND GEOIDAL COMPONENTS OF SEA SURFACE TOPOGRAPHY.	GS-11
Victor Zlotnicki	

DEPARTMENT OF BIOLOGY

John M. Teal, Department Chairman

BIOLOGY

MICROSOMAL ELECTRON TRANSPORT AND
XENOBIOTIC MONOOXYGENASE ACTIVITIES
DURING THE EMBRYONIC PERIOD OF DEVELOPMENT
IN THE KILLIFISH, Fundulus Heteroclitus

Robert L. Binder and
John J. Stegeman

The developmental patterns of aryl hydrocarbon hydroxylase (AHH) activity and NADPH-cytochrome c reductase activity were followed during the embryonic period of development in Fundulus. AHH activity was localized in microsomal fractions prepared from whole Fundulus embryos and eleutheroembryos. On the basis of this subcellular localization, the requirements of O_2 and NADPH for activity and sensitivity to carbon monoxide and cytochrome c inhibition, the AHH activity in Fundulus embryos and eleutheroembryos appeared to be cytochrome P-450-dependent. AHH activity was measurable in stages prior to the appearance of the liver rudiment, and during subsequent embryonic development the extrahepatic tissues were likely to have contributed substantially to the AHH activity measured. At all stages assayed before hatching, microsomal AHH specific activity increased about 9-fold. This posthatching increase in AHH activity was not age-dependent, nor developmental stage-dependent, but rather required hatching, and was not due to the presence of endogenous inhibitors in pre-hatching stages. The levels of NADPH-cytochrome c reductase activity and AHH activity were not closely correlated in whole embryo and eleutheroembryo microsomes, but the AHH activity in these preparation apparently was not limited by the levels of the NADPH-cytochrome c (P-450) reductase. The presence of AHH activity in Fundulus embryos during the period of active organogenesis, prior to hatching, indicates that this species is likely to be susceptible to a variety of teratogens requiring metabolic activation, and this may be the case for other species of fish as well.

In press: Toxicology and Applied
Pharmacology.

Supported by: NSF Grants OCE77-24517
and OCE80-18569.

WHOI Contribution No. 5555.

BLUEFIN TUNA WARM THEIR VISCERA
DURING DIGESTION

Francis G. Carey, John W. Kanwisher
and E. Don Stevens

Acoustic telemetry showed that stomach temperature from large bluefin tuna, Thunnus thynnus, held in an impoundment, changes markedly during feeding. The stomach cools rapidly on ingesting cold food. It then warms to a maximum of 10 to 15°C above water temperature over a period of 12 to 20 hours. Temperature decreases slowly over the next 20 to 30 hours to a final state where it remains 3 to 6° above water temperature. The viscera are thermally isolated. Conductive heat losses are reduced by an overlying gas bladder and by the thick fatty muscle of the body wall. Convective heat losses are prevented by heat exchangers in the circulation. The temperature rise can be accounted for by heat released in the hydrolytic processes of digestion and by an increase in metabolic rate. The elevated temperatures should speed digestion and allow the tuna to feed frequently when food is abundant.

Supported by: NSF Grants PCM76-81612
and PCM78-03250; NOAA-NMFS; and
National Research Council of Canada.

WHOI Contribution No. 5515.

PECTORAL FIN LOCOMOTION:
SWIMMING ENERGETICS OF CUNNERS AND
TAUTOGS (LABRIDAE)

Marvin A. Freadman

Cunners (Tautogolabrus adspersus) and tautogs (Tautoga onitis) use their fanlike pectoral fins as a primary means of sustaining locomotion at slow and intermediate swimming velocities. Both species use asynchronous pectoral fin movements at slow speeds (up to approximately 1 bl's⁻¹), synchronous movements at intermediate speeds (up to approximately 2.5 bl's⁻¹) and body/caudal fin movements at high speeds (> 2.5 bl's⁻¹). Maximum sustainable swimming speeds for pectoral fin locomotion, determined in increasing velocity increment experiments, for cunners are 50 cm's⁻¹ (3.94 bl's⁻¹) in 30 minute swimming bouts and 41.5 cm's⁻¹ (3.27 bl's⁻¹) in 45 minute bouts. The values for tautogs are 52 cm's⁻¹ (4.1 bl's⁻¹) in 30 minute bouts and 43 cm's⁻¹ (3.1 bl's⁻¹) in 45 minute bouts.

The metabolic characteristics of cunners and tautogs during steady swimming are unique. During asynchronous and synchronous pectoral fin locomotion (slow and intermediate speeds), VO_2 increases linearly with increasing velocity. When the pectoral fins can no longer provide enough propulsive

force (at high speeds), the body/tail sweeps side to side. With body/tail propulsion, the line describing the change in VO_2 with swimming velocity increases in slope. When portions of the pectoral fins are ablated (bilaterally), the caudal fin is used more such that the more pectoral fin absent, the lower the velocity at which the body/caudal fin provides propulsive force. Measurements of VO_2 at velocities during which individual animals used pectoral fin locomotion then body/caudal fin locomotion after pectoral fin ablation, indicate less energy is expended at intermediate swimming velocities with pectoral fin locomotion. The results indicate cunners and tautogs routinely use pectoral fin locomotion as a primary means of generating sustained propulsive force at slow and intermediate swimming velocities. Values for maximum sustainable swimming speeds approach those of many fishes relying on body and caudal fin propulsion.

Supported by: NIH Grant 1 F32 AM05976
and WHOI.

WHOI Contribution No. 5510.

SWIMMING/LOCOMOTOR CAPACITY OF HUDSON RIVER STRIPED BASS

Marvin A. Freadman, Frederick P. Thurberg
and Anthony C. Calabrese

The life history of active fishes such as striped bass is characterized by significant periods of swimming activity. The process of maneuvering in water relates to the swimming capacity of a fish and is essential for events such as migration, escape from predators and food capture. Thus, lack of significant swimming capacity in active fish species could severely limit the survival, distribution and abundance of the species.

The present study was undertaken to investigate the swimming performance of striped bass from impacted and non-impacted waters. Striped bass were collected from two Hudson River sites (river mile 1 and river mile 39), from eastern Long Island Sound (Greenport-Orient Point area) and from the Edenton National Hatchery (N.C.). The indices of swimming performance measured were critical swimming speed and burst swimming speed. These experiments were performed using a water tunnel treadmill. The results indicate both groups of fishes from the Hudson River had significantly lower swimming capacity compared to animals collected from eastern Long Island Sound and the Edenton Hatchery.

A further aspect of our study dealt with exposures of striped bass to lead (20,100 and 200 ppb) for 60 days in the laboratory in a proportional diluter apparatus. Only at the highest exposure concentration, were

we able to detect significantly lower critical swimming speeds compared with control animals.

We obtained some measurements on the PCB content of various tissues of striped bass collected from the Hudson River. PCB's were present in the ppm range and the highest quantities were found in the red swimming musculature. All routine swimming activity in striped bass is powered by this type of musculature.

Supported by: NMFS, NOAA Office of
Marine Pollution Assessment and
WHOI.

WHOI Contribution No. 5558.

INTRODUCTION TO THE BIOLOGY OF HYDROTHERMAL VENTS

J. Frederick Grassle

The main features of hydrothermal vent communities are described. A brief discussion of the relationships between studies of metabolism, growth, feeding relationships, and life span is presented along with recent results from vent areas discovered since the Galapagos expedition.

Published in: Hydrothermal Processes at
Seafloor Spreading Centers, Plenum
Publishing Co., NY.

Supported by: NSF Grant OCE81-15251.

WHOI Contribution No. 5505.

AN ALTERNATIVE 7-ETHOXYRESORUFIN O-DEETHYLASE ACTIVITY ASSAY: A CONTINUOUS VISIBLE SPECTROPHOTOMETRIC METHOD FOR MEASUREMENT OF CYTOCHROME P-450 MONOOXYGENASE ACTIVITY

Alan V. Klotz, John J. Stegeman
and Chris Walsh

A procedure to directly measure the cytochrome P-450-dependent 7-ethoxyresorufin O-deethylase activity with a visible spectrophotometer is described and compared to the standard fluorometric method, which yielded identical results with β -naphthoflavone-treated rat liver microsomes. The assay takes advantage of a clean distinction in visible absorption spectra obtained for highly purified 7-ethoxyresorufin (substrate) and resorufin (enzymatic product). The purification and characterization of resorufin, the enzymatic product, is detailed and its extinction coefficient ($\epsilon_{572} = 73 \text{ mM}^{-1} \text{ cm}^{-1}$) provides for an accurate quantitation of enzyme activity. The large visible extinction coefficient of

the product chromophore provides a high sensitivity for low activity samples. The application of this enzyme assay in a visible spectrophotometer, along with the considerable evidence that a single aromatic hydrocarbon-inducible cytochrome P-450 isozyme is responsible for the catalysis, enhances the utility of this substrate in microsomal monooxygenase assays. The utility of the visible assay is demonstrated by the simple determinations of the coupling ratio for 7-ethoxyresorufin oxidation in Stenotomus chrysops liver microsomes and the K_1 for 7,8-benzoflavone inhibition of the enzymatic reaction.

Supported by: NSF Grant OCE80-18569 and NIH Grant GM21643.

WHOI Contribution No. 5544.

AN ARYL HYDROCARBON HYDROXYLATING
HEPATIC CYTOCHROME P-450 FROM THE
MARINE FISH Stenotomus Chrysops

Alan V. Klotz, John J. Stegeman
and Christopher Walsh

Hepatic microsomal cytochrome P-450 from the untreated coastal marine fish scup, Stenotomus chrysops, was solubilized and resolved into five fractions by ion exchange chromatography. The major fraction, cytochrome P-450E ($M_r = 54,300$), was further purified to a specific content of 11.7 nmol heme/mg protein and contained a chromophore absorbing at 447 nm in the CO-ligated, reduced difference spectrum. NH_2 -terminal sequence analysis of cytochrome P-450E by Edman degradation revealed no homology with any known cytochrome P-450 isozyme in the first nine residues. S. chrysops liver NADPH-cytochrome P-450 reductase, purified 225-fold ($M_r = 82,600$), had a specific activity of 45-60 U/mg with cytochrome c, contained both FAD and FMN, and was isolated as the one electron reduced semiquinone.

Purified cytochrome P-450 metabolized several substrates including 7-ethoxycoumarin, acetanilide and benzo[α]pyrene when reconstituted with lipid and hepatic NADPH-cytochrome P-450 reductase from either S. chrysops or rat. The purified reconstituted system was sensitive to inhibition by 100 μM 7,8-benzoflavone and analysis of products in reconstitutions with purified rat epoxide hydrolase indicated a preference for oxidation on the benzo-ring of benzo[α]pyrene consistent with the primary features of benzo[α]pyrene metabolism in microsomes. Cytochrome P-450E is identical to the major microsomal aromatic hydrocarbon-inducible cytochrome P-450 by the criteria of molecular weight, optical properties and catalytic profile. It is suggested that substantial quantities of this aromatic

hydrocarbon-inducible isozyme exist in the hepatic microsomes of some untreated S. chrysops. The characterization of this aryl hydrocarbon hydroxylase extends our understanding of the metabolism patterns observed in hepatic microsomes isolated from untreated fish.

Published in: Archives of Biochemistry and Biophysics, 226, 578-592.

Supported by: NIH Grant GM 21643 and NSF Grant OCE80-18569.

WHOI Contribution No. 5347.

PHYSIOLOGY OF THE WOOD BORING MOLLUSC
Martesia cuneiformis SAY

Roger Mann and Scott M. Gallager

Larvae of Martesia cuneiformis Say settled in greatest abundance (151.5 individuals/square meter surface area of wood substrate) during the months of August, September and October in a North Carolina estuary. Minimum estimates of post settlement shell growth rates are 0.19 mm/day during the summer months and 0.03 mm/day during the winter months. Gravid adults maintained in the laboratory spawn during physical disturbance of dissection from wood and, in isolated individuals, after thermal stimulation. The first shelled veliger has a mean shell size of 69.0 μm long and 60.0 μm high. At 22°C and 30-32 o/oo salinity the pediveliger stage, with a mean shell size of 260 μm long and 269 μm high, is reached in 35 days. Larval respiration rate is described by the relationship $R = 2.22 W^{2.42}$ where R is oxygen consumption (nl O_2 /larva/hr) and W is dry weight (μg). On starvation, larvae predominantly utilize protein and lipid as respiratory substrates. Adult respiration and ammonia excretion rates are described by the relationships $R = 17.95 W^{0.703}$ and $E = 0.133 W^{0.492}$ respectively with the units R ($\mu l O_2$ /individual/hr), E (μg at NH_3 -N/individual/hr) and W (gm live weight).

In press: Biological Bulletin.

Supported by: ONR Contract N00014-82-C-0071 NR 083-004.

WHOI Contribution Number 5473.

SWIMMING BEHAVIOUR OF LARVAE OF THE OCEAN
QUAHOG Arctica islandica IN RESPONSE TO
PRESSURE AND TEMPERATURE

Roger Mann and Charlotte C. Wolf

Trochophore larvae of the ocean quahog, Arctica islandica, swim continuously. At a constant temperature of 12°C they are negatively geotactic, have no phototactic response and exhibit no change in swimming behaviour in the pressure range 1-3 bars. In a vertical thermal gradient from 9 to 25°C, the trochophores swim throughout the experimental chamber and show no temperature preference. Veliger larvae of A. islandica alternate between periods of active upward swimming in vertically oriented helices, that is a negative geotaxis, and periods of passive sinking with the velum either trailing or retracted between closed valves. They do not swim in the horizontal plane and, in the length range 160 µm-202 µm, show no phototactic response. When exposed to sequential increases and decreases in hydrostatic pressure at 12°C larvae in the length range 160 µm-196 µm consistently exhibited a net upward movement following an increase in pressure and a net downward movement following a decrease in pressure. The threshold pressure change to elicit response is <0.5 bars. Larvae of 170 µm length respond to increased pressure by decreasing the diameter of the helix and increasing the height gain per rotation. Larvae of 196 µm length respond to increased pressure by increasing height per rotation and vertical velocity. Larvae of 202 µm length exhibit no significant change in swimming behaviour with increased pressure. In a vertical thermal gradient, early veligers swim in the range 7-23°C with preferential aggregation, depending upon size, in the range 12-18°C. Larvae of 204 µm length show no temperature preference in the range 6-20°C. The implication of the observed behaviour on seasonal depth distribution of A. islandica larvae in the Middle Atlantic Bight is discussed.

Published in: Marine Ecology - Progress Series, 13, 211-218.

Supported by: NOAA, Office of Sea Grant, Grant NA80-AA-D-00077 and a Summer Student Fellowship from WHOI.

WHOI Contribution No. 5276.

HYPOSMOREGULATION IN AN ANADROMOUS
TELEOST: INFLUENCE OF SEX AND MATURATION

Stephen D. McCormick and Robert J. Naiman

Physiological differences between sexes of the same species are generally related to primary and secondary sexual characteristics

or to adaptations resulting from differential strategies for reproductive success. Reports of sexual divergence in physiological processes are rare in lower vertebrates, except for those directly associated with reproduction or mate competition. We report decreased salinity tolerance and hypo-osmoregulatory ability in mature male anadromous brook trout (Salvelinus fontinalis) that does not occur in females or immature males. Lowered salinity tolerance of adult males becomes acute during autumn photoperiod when normal spawning occurs. Plasma Cl^- , Mg^{++} , osmolarity and hematocrit are significantly higher in mature males after transfer to seawater, relative to mature females. Reduced adult male hypoosmoregulatory ability explains skewed sex ratios in anadromous populations, may limit the extent of anadromy and may have been a significant phase in the evolution of salmonid anadromy.

Supported by: NOAA, Office of Sea Grant, Grant NA80-AA-D-00077 (R/A-14); a Tai-Ping Predoctoral Fellowship in Marine Biology; and the WHOI Education Office.

WHOI Contribution No. 5418.

SOME DETERMINANTS OF MATURATION IN BROOK
TROUT, Salvelinus fontinalis

Stephen D. McCormick and Robert J. Naiman

Size, age, growth rate and photoperiod were examined for their effects on the timing of maturation and the proportion of mature male and female brook trout (Salvelinus fontinalis). Photoperiod completely entrained the gonadosomatic index and the timing of functional maturation (spermiation). High feed and delayed photoperiod conditions resulted in a greater proportion of mature individuals of each sex in their first year, though the percent maturation in each experimental group was higher for males. Size by itself can explain almost all of the variation in the percent maturation across all experimental groups ($r^2 = 0.95$ and 0.90 for males and females, respectively). The results indicate that 1) given a positive growth rate, age and growth rate are less important than size in determining maturation of brook trout and 2) the maturation response to size is sexually divergent. Size dependent maturation has important implications for the control of maturation in domestic stocks, our understanding of brook trout demography, and for future investigations of the physiological and hormonal control of salmonid maturation. A conceptual model depicting the effects of environmental factors on size and the initiation of maturation is presented.

Supported by: NOAA, Office of Sea Grant,
Grant NA80-AA-D-00077 (R/A-14); the
WHOI Matamek Research Program; a
Tai-Ping Predoctoral Fellowship; and
the WHOI Education Office.

WHOI Contribution No. 5537.

DIFFERENTIAL REGULATION OF HEPATIC
XENOBIOTIC AND STEROID METABOLISM IN
MARINE TELEOST SPECIES

John J. Stegeman and Bruce R. Woodin

Sex differences in hepatic microsomal cytochrome P-450 content observed in three teleost species, winter flounder (Pseudopleuronectes americanus), killifish (Fundulus heteroclitus) and brook trout (Salvelinus fontinalis), were qualitatively similar, with the specific content in males being between 5 and 10-fold greater than that in females. In the winter flounder the specific content of hepatic cytochrome P-450 in males was particularly high, ranging up to 2.0 nmol/mg microsomal protein. Distinctions in specific content between gonadally mature and immature animals of the same sex, and similar size, indicated the differences were due principally to a suppression of the levels of P-450 in the ripe females in both Fundulus and winter flounder. Sex differences in the specific activity of ethoxyresorufin O-deethylase (EROD) and benzo[α]pyrene hydroxylase (B[α]PH) activities were correlated with the differences in cytochrome P-450 specific content in all three species, although a sex difference in the estimated turnover number (nmol/min/nmol P-450) for ER O-deethylase activities was measurable only in flounder. In contrast to these xenobiotic monooxygenase activities, hepatic testosterone hydroxylase activity per nmol of cytochrome P-450 was 5-10 fold greater in gonadally mature females than in gonadally mature males of both winter flounder and Fundulus. This was true of both 6 β -hydroxylase activity and of 16 β -hydroxylase activity (Table 1). The sex difference in testosterone 6 β -hydroxylase activity in brook trout was opposite to that seen in the marine species, but sex differences in testosterone 16 β -hydroxylase activity in the brook trout were like those in the marine species. The data indicate that some teleost hepatic cytochromes P-450 involved in steroid metabolism are regulated by hormonal factors in a way that is different from the regulation of the bulk of cytochrome P-450, including some cytochromes P-450 involved in xenobiotic metabolism.

In press: Marine Environmental Research.

Supported by: NSF Grant OCE80-18569.

FOOD SELECTION BY LARVAL GULF MENHADEN
(Brevoortia patronus)

Diane Stoecker and John J. Govoni

In laboratory experiments in which the dinoflagellate, Prorocentrum micans, the tintinnid, Favella sp., and N1 nauplii of the copepod, Acartia tonsa, were used as food organisms, small (average length 3.9-4.2 mm) laboratory-reared gulf menhaden (Brevoortia patronus) larvae ate Prorocentrum and Favella, but not nauplii. Larvae ate Favella sp. in significantly ($P < 0.001$) disproportionate numbers, indicating their preference for this tintinnid. Larger larvae (average length 6.4 mm) ate copepod nauplii as well as the other food organisms.

This feeding response is consistent with larval feeding in the northern Gulf of Mexico where small gulf menhaden larvae (3.01 to 5.00 mm) frequently ate large numbers of dinoflagellates (mostly Prorocentrum micans and compressum) and tintinnids (mostly Favella sp.), but did not eat copepod nauplii. As larvae grew, copepod nauplii and other food organisms became important, while dinoflagellates and tintinnids became relatively less important in the diet. Inasmuch as the Favella and nauplii used in the laboratory feeding experiments were roughly similar in size as well as carbon and nitrogen contents, the feeding selectivity and dietary ontogeny that we observed were likely due to a combination of prey capturability and larval fish maturation and learning.

Supported by: NOAA, Grant NA81RAD00014
and a Contract from the Ocean
Assessments Division, National Ocean
Services, NOAA.

WHOI Contribution No. 5575.

A Q U A C U L T U R E

BIVALVE MOLLUSC HATCHERIES: A CRITICAL
APPRAISAL OF THEIR DEVELOPMENT AND A
REVIEW OF THEIR POTENTIAL VALUE IN
ENHANCING THE FISHERIES OF DEVELOPING
NATIONS

Roger Mann

Over 100 years ago, Brooks succeeded in rearing and describing the development of Crassostrea virginica in the laboratory. This achievement marked the birth of the shellfish hatchery industry, and held the promise of a day when oyster seed would be grown by the million for use in commercial enterprise. Another 70 years were to pass

RESOURCE PARTITIONING AND AGGRESSIVE
INTERACTIONS BETWEEN BROOK TROUT
(Salvelinus fontinalis) FRY AND NINESPINE
STICKLEBACKS (Pungitius pungitius)

Alaine Gaudreault, Tandy Miller,
Gerry J. FitzGerald and W. Linn Montgomery

The food habits of trout fry (Salvelinus fontinalis) and ninespine sticklebacks (Pungitius pungitius) were compared in the Matamek River, Québec. The aggressive behavior of both species were observed in the field and under allopatric and sympatric conditions of varying densities in a stream tank. Our results showed a considerable overlap in food in August. The frequent aggressive interactions were usually begun by sticklebacks toward trout fry in the field but not in the stream tanks. Subsequent experimental manipulation of adult stickleback and trout fry densities in stream tanks showed increased intra- and interspecific aggression at higher than at lower densities although there was no simple monotonic relationship between aggression and density. The results suggest that in areas where both species are sympatric and are abundant, sticklebacks may have a serious impact upon the ecology of the brook trout.

Supported by: WHOI Matamek Research
Program and N.S.E.R.C. and F.C.A.C.
Grants.

Contribution No. 5514.

DYNAMICS OF PLANT LITTER DECAY IN
NORTHERN STREAMS

Jerry M. Melillo, Robert J. Naiman,
John D. Aber and Alan E. Linkins

In this paper we consider the influences of initial litter quality, exogenous nutrient supply, temperature and oxygen tension on decay rates and nutrient immobilization potentials of plant litter materials decomposing in northern streams. We review a new method of estimating nitrogen and phosphorus immobilization potentials in decaying litter and we discuss the mechanisms responsible for nitrogen immobilization.

Initial litter quality is the dominant factor that influences decay rate. Materials rich in nitrogen and poor in lignin generally decay most rapidly. Addition of nitrogen and/or phosphorus to the water often result(s) in increased decay rate as does elevation of water temperature. While low oxygen tension retards the decay of lignin and related compounds, it is not clear that low oxygen tension slows the rate of decay of more labile compounds.

Initial litter quality is also the dominant factor that influences nitrogen immobilization potential. Materials poor in both nitrogen and lignin generally have the highest immobilization potentials. Addition of nitrogen to the water increases the immobilization potential of a litter material. Plant litter decomposing under anaerobic conditions immobilizes less nitrogen than the same litter decomposing under aerobic conditions. And, all other conditions being equal, temperature changes will influence nitrogen immobilization rate but not its maximum amount.

The amount of nitrogen directly associated with microbial biomass makes up only a small fraction of the total nitrogen mass in decomposing litter. Microbes appear to be important agents in the nitrogen immobilization process on decaying litter. The microbes produce exoenzymes which in turn cause the degradation of large molecules such as polyphenolics and lignins. Some of these degradation products, such as the "reactive phenolics", recondense with nitrogen containing compounds. The degradation-recondensation cycle continues throughout litter decay, with the ultimate product being nitrogen-rich humic substances.

Supported by: NSF Grants DEB80-05081,
DEB81-05677 and DEB82-07055.

WHOI Contribution No. 5547.

NITROGEN BUDGET OF A SUBARCTIC STREAM
ALTERED BY BEAVER (Castor canadensis)

Robert J. Naiman and Jerry M. Melillo

Beaver (Castor canadensis) influence stream ecosystems through their wood cutting and dam building activities. To quantify this influence we have used measured rates of nitrogen dynamics to construct a nitrogen budget for a section of a second order stream in eastern Québec and a beaver dam in that stream. The budget demonstrates the importance of sediment accumulations and an expanded wetted area to the annual nitrogen economy and to pathways of nitrogen cycling. Major changes after impoundment (per unit area) include a reduction in allochthonous nitrogen inputs and an increase in nitrogen fixation by sediment microbes. Overall, the beaver-modified stream section accumulated $\sim 10^3$ times more nitrogen than before alteration. The ecosystem implications of beaver activity suggest that current concepts of patterns and processes in running waters require modification.

Supported by: NSF Grant DEB81-05677 and
the WHOI Matamek Research Program.

WHOI Contribution No. 5565.

SALT MARSH SYSTEMS

WATER UPTAKE BY ROOTS IN SHORT *Spartina* MARSH: MASS FLOW OF AIR INTO SEDIMENT AND INTO PLANTS

John W. H. Dacey and Brian L. Howes

Downward movement of the water table during day and night in the short grass zone of salt marshes is not drainage, but results from water uptake by roots. Removal of water from the sediment also leads to air entry into the sediment, suggesting a feedback between plant growth, water uptake, and sediment oxidation. The water balance of marsh grass appears to influence the internal morphology of its roots, giving rise to a new mechanism for mass flow of gas in plants.

Supported by: NSF Grants DEB80-21741, DEB80-04701, DEB79-05127 and DEB81-19819.

WHOI Contribution No. 5533

CARBON FLOW THROUGH OXYGEN AND SULFATE REDUCTION PATHWAYS IN SALT MARSH SEDIMENTS

Brian L. Howes, John W. H. Dacey
and Gary M. King

We measured surface O_2 uptake, $^{35}SO_4$ reduction and total sediment metabolism (CO_2 production) in sediments (0-30 cm) supporting stands of short *Spartina alterniflora* in a New England salt marsh. Sediment respiration parallels below-ground plant biomass. CO_2 production was highest at the surface where O_2 was present and declined exponentially with depth. In deeper sediments (greater than 5 cm), CO_2 production is equal to $^{35}SO_4$ reduction as determined by the Cr^{+2} reduction technique. Previous estimates of $^{35}SO_4$ reduction, based on aqua-regia digestion, have yielded rates several times higher than CO_2 production. Our time course experiments using CO_2 production and $^{35}SO_4$ reduction (by Cr^{+2} reduction and aqua regia digest) indicate that the aqua regia technique is not a reliable method for measuring SO_4 reduction and that the rate of SO_4 reduction is much less than previously reported for this marsh.

Our estimates of annual O_2 uptake agree with CO_2 production estimates and below-ground production estimates for this marsh. We estimate carbon remineralization to average about $200 \text{ mmol C m}^{-2} \text{ day}^{-1}$, among the highest measured for marine sediments. Our simultaneous measurements of O_2 uptake, CO_2 production and SO_4 reduction suggest that at least half of this decomposition occurs via SO_4 reduction, in the range

observed for marine sediments. We conclude that the mechanisms underlying carbon and sulfur cycling in salt marsh sediments are much more similar to subtidal sediments than previously recognized.

Supported by: NSF Grants DEB79-05127, DEB80-21741, DEB81-04701 and NSF EPSCOR.

WHOI Contribution No. 5468.

ANNUAL CARBON REMINERALIZATION AND BELOW-GROUND PRODUCTION OF *Spartina alterniflora* IN A NEW ENGLAND SALT MARSH

Brian L. Howes, John W. H. Dacey
and John M. Teal

The annual rates and depth distribution of organic carbon remineralization to CO_2 was determined in sediments supporting stands of short *Spartina alterniflora*. CO_2 production was estimated by two independent techniques. We constructed a CO_2 budget based on measurements of CO_2 emission from the sediment to the atmosphere, export of dissolved inorganic carbon in porewater exchange and changes in the porewater pool of dissolved organic carbon throughout the year. We also conducted the first known measurements of CO_2 production in salt marsh sediments by monitoring changes in total inorganic carbon in undisturbed cores. Both estimates yielded similar results. These data were used to estimate the total annual mineralization of organic carbon in these sediments ($72-75 \text{ mol C m}^{-2} \text{ yr}^{-1}$).

We also measured losses of organic matter as methane, dissolved organic carbon and burial in order to construct a carbon budget for these sediments. These data were combined with the estimates of carbon remineralization to yield an estimate of organic carbon loading to the sediment. We estimate that below-ground production in short *S. alterniflora* in this Massachusetts marsh is $79.5-83.7 \text{ mol C m}^{-2} \text{ yr}^{-1}$, of which about 90% decomposes to CO_2 in situ and less than 5% is exported from the sediment.

Supported by: NSF Grants DEB79-05127, DEB80-21741 and DEB81-04701.

WHOI Contribution No. 5467.

EFFECTS OF SAMPLING TECHNIQUE ON
MEASUREMENTS OF POREWATER CONSTITUENTS
IN SALT MARSH SEDIMENTS

Brian L. Howes, John W. H. Dacey
and Stuart Wakeham

We examined the effect of three methods of porewater extraction on the measured concentrations of seven porewater constituents in vegetated salt marsh sediments. Porewater extraction by standard core sectioning followed by squeezing or centrifuging is compared with two less destructive methods of water collection. The core sectioning technique significantly increased the concentration of total dissolved organic carbon and dimethylsulfide presumably due to root damage by core cutting. Sampling technique did not appear to influence concentrations of interstitial salinity, dissolved inorganic carbon, dissolved ammonium, soluble sulfides and dissolved sulfate as long as proper precautions were taken to prevent the loss of the reduced compounds.

Supported by: NSF Grants BSR80-21741,
BSR81-04701 and BSR83-08800.

WHOI Contribution No. 5561.

EFFECTS OF O₂ ON AMMONIUM UPTAKE
AND ROOT RESPIRATION BY
Spartina alterniflora

James T. Morris and John W. H. Dacey

Previously, the growth of Spartina alterniflora has been found to be limited by nitrogen and correlated with sediment redox potential. In this study, we have investigated a possible connection between these two factors. We have found that internal O₂ transport is insufficient to saturate NH₄ uptake in short S. alterniflora in hydroponic culture. Rates of NH₄ uptake and root respiration were very sensitive to O₂ concentration in the rhizosphere, saturating at about 5% O₂. However, NH₄ uptake continued at a reduced rate for at least 4 hr under anaerobic conditions. Plant to plant variations in anaerobic rates of NH₄ uptake and root respiration were significantly correlated to the diffusion rate of CH₄ tracer gas from the leaves to the roots of individual plants.

Supported by: NSF Grants DEB81-15503 and
DEB80-21741.

WHOI Contribution No. 5507.

THE ROLE OF ONE SALT MARSH IN
COASTAL PRODUCTIVITY

John M. Teal

We have been intensively studying the ecology of Great Sippewissett Salt Marsh, Massachusetts, for the last 12 years by experimentally fertilizing marsh plots at different levels of nutrients with fertilizers that both do and do not contain sewage sludge and its contaminants. Plant productivity is limited by nitrogen availability which is in turn modified by sediment oxidation-reduction potential. Species composition of the plant cover is also affected by nitrogen. At greater levels of nitrogen availability, nitrogen in the grasses increases. This leads to increased populations and/or productivity of herbivores, detritivores and decomposers. The nitrogen budget for the marsh shows that it imports nitrate, and exports nitrogen gas, particulate organic nitrogen and ammonia. The sulfur cycle is dominant in underground decomposition and transports energy to the sediment surface. A variety of fishes, especially young stages including those of commercially important species, make use of the marsh as a feeding place. Heavy metal pollutants which form insoluble sulfides tend to be immobilized in sediments and not accumulated in organisms; e.g., lead. Those such as cadmium which form soluble complexes in sea water do not accumulate in the marsh and do, at least, temporarily accumulate in organisms. Organic pollutants are bound to sediments and the one studied, Aldrin, disappeared about two years after it was added.

Supported by: NSF Grants DEB79-05127 and
DEB80-12437.

WHOI Contribution No. 5408.

THE IMPORTANCE OF CHEMICAL COMPOSITION
OF SALT MARSH LITTER ON DECAY RATES
AND FEEDING BY DETRITIVORES

Ivan Valiela, John Wilson,
Robert Buchsbaum, Carol Rietsma,
Donald Bryant, Kenneth Foreman
and John Teal

This paper uses experiments done in the field and laboratory with salt marsh organic matter to show that the chemical composition of the organic matter affects decay rates. Among the many components of the organic matter of dead S. alterniflora, nitrogen - especially soluble nitrogen - content, soluble phenolics, and lignin concentrations seem likely to be important in this respect.

The important effects of composition take place in the context of other environmental variables. Specifically, the effects of

chemical composition interact with other critical factors, such as redox and temperature. These mechanisms impose the controls on decay. Models of decomposition need therefore to include these control mechanisms if they are to realistically describe and make predictions about decay processes.

Although soluble nitrogen and phenolics do not make up much of the weight of detritus, they may be important as controls of various processes. We can show, for example, that, at least for one major detritivore in salt marshes, these classes of compounds are important. Soluble protein and ferulic acid make up part of a hierarchy of cues that determines feeding by this detritivore. Moreover, the growth achieved by detritivores feeding on detritus with differing composition differs. Again, the higher the available nitrogen, the lower soluble phenolic and lower lignin content seem to promote higher growth. Thus, there is a correspondence of chemical composition of litter, palatability to detritivores, and, apparently, assimilation of detritus and production by consumers.

Presented at: Detritus Dynamics in
Aquatic Ecosystems Symposium,
Skidaway Inst. of Oceanography,
November 1983

Supported by: NSF Grants DEB79-05127
and BSR82-17040.

WHOI Contribution No. 5559.

M A R I N E M A M M A L S

ANATOMY AND VASCULATURE OF A MINKE WHALE HEART

Christina Ochrmowych and
Richard Lambertsen

The heart from a 4 m long minke whale (*Balaenoptera acutorostrata*) was studied to determine the details of its anatomy and to consider cardiac adaptations to diving. The volume fraction (Vvc) of capillaries in the wall of the left ventricle was determined at different levels from base to apex and at different depths from epi- to endocardium using a light microscopic stereologic technique. Typical of cetaceans, this minke whale heart was distinctly flattened dorso-ventrally. A moderator band, characteristic of ungulate hearts, spanned the right ventricle. The right and left atrioventricular valves were tricuspid and bicuspid, respectively. The right coronary artery supplied the dorsal and right lateral myocardium. The left coronary artery supplied the ventral and left lateral

myocardium. An anastomosis between the dorsal and ventral interventricular arteries occurred in the dorsal interventricular groove. Stereologically, a decreasing transmural gradient in Vvc was identified between the epicardium and the subepicardium at 15 cm from the apex. Our results, however, did not reveal any significant deviations in the pattern of capillary distribution in the wall of the left ventricle between this baleen whale and terrestrial mammals. Measurements of the heart, great vessels, coronary vasculature and ventricle walls are also given, and suggest a physiologic and adaptive right ventricular hypertrophy. Based on these and other observations we propose that the relatively great thickness of the right ventricle and the distinctive shape of the cetacean heart are adaptations to the hemodynamic changes and collapse of the thorax associated with apneic diving.

In press: American Journal of Anatomy.

Supported by: New York Sea Grant
Institute and the WHOI Coastal
Research Center.

WHOI Contribution No. 5521.

STATUS OF *Eubalaena glacialis* OFF CAPE COD

William E. Schevill, William A. Watkins
and Karen E. Moore

Historic and current records of abundance and distribution of *Eubalaena glacialis* near Cape Cod are scanty and imprecise, and cannot be considered a census. The historical records since 1620 and ours from 1955 - 1981 both show a seasonal peak in April and few sightings in summer months. During the last quarter century, our counts show up to 131 individuals in one year; 70+ were the most seen in one day. We have seen 21 calves, up to 4 in one year, with a 3-year reproductive cycle indicated. Individual whales usually remain in the area only a few days at a time. The seasonal shifts do not compellingly indicate a traditional (N - S) migration pattern. The evidence available does not show that the *E. glacialis* in our waters nowadays are really very much fewer than they were three and a half centuries ago.

Supported by: ONR Contract N00014-82-
C-0019 NR 083-004.

WHOI Contribution No. 5419.

SONAR OBSERVATIONS EXPLAIN BEHAVIORS
NOTED DURING BOAT MANEUVERS FOR RADIO
TAGGING OF HUMPBACK WHALES
(Megaptera novaeangliae)
IN THE GLACIER BAY AREA

William A. Watkins and Camille A. Goebel

Humpback whales (Megaptera novaeangliae) in Glacier Bay, Alaska, were very difficult to approach closely for radio tagging. They generally seemed unconcerned with the boat beyond 150 m, but at about that distance they often reacted sharply as if the boats were suddenly more noticeable. In contrast, these whales in Icy Strait (off Point Adolphus, Chicagof I.) appeared to be aware of the boats at greater distances, but they could be approached quite closely, especially when the boats and whales were separated by tidal current lines. Sonar observations showed sharp downward refraction in the surface waters of Glacier Bay and good acoustic sound transmission in Icy Strait except across discontinuities in water masses that occurred at the boundaries of separate tidal currents. Although the whales in both areas submerged for periods up to 12 minutes and they usually raised their flukes out of water as they went down, they did not dive deeply. Only 5 out of more than 129 dives were deeper than 20 m - one of these was to a depth of 40 m.

In press: Cetology.

Supported by: NOAA, National Marine
Mammal Laboratory (82-ABA-2122) and
ONR Contract N000-14-82-C-0019 NR
083-004.

WHOI Contribution No. 5374.

SOUNDS OF THE ROSS SEAL, Ommatophoca rossi

William A. Watkins and G. Carleton Ray

In-air and underwater sounds were recorded from Ross seals, Ommatophoca rossi, during January 1966 in the western Ross Sea, Antarctica, from Cape Hallet to Cape Adare. The in-air vocalizations produced by seals hauled out on ice included short (0.05 to 0.1 sec) tonal pulses with downward sweeping frequency, and longer (1 to 1.5 sec) siren calls with downward and upward sweeping frequency, somewhat like that of a police siren. The frequencies of the in-air sounds ranged from 100 to 1000 Hz. Underwater vocalizations included pulses and siren calls and were similar to the in-air sounds except that they had greater and mostly higher-frequency ranges (1 to 4 kHz). The underwater siren calls were pulse-modulated at a consistent rate of about 650 per sec, and they usually were formed of two indepen-

dently varying tones each with separate sideband harmonics. Overlapping underwater sound sequences were heard from seals scattered throughout the areas.

Supported by: NSF, Office of Antarctic Programs, to the New York Zoological Society and The Johns Hopkins University, and ONR, Oceanic Biology Program, Contract N000-14-82-C-0019 NR 083-004.

WHOI Contribution No. 5471.

M A R I N E P O L L U T I O N

THE EFFECTS OF POLLUTANTS ON MARINE
ZOOPLANKTON POPULATIONS: SUBLETHAL
PHYSIOLOGICAL INDICES OF POPULATION STRESS

Judith M. Capuzzo

Sublethal responses are important in predicting disruptions in population stability and production associated with waste exposure. Reductions in feeding rates, respiration rates and production rates of copepods were observed with exposure to DuPont Grasselli waste, an industrial waste discharged at Deepwater Dumpsite 106, suggesting that components of the waste interfere with normal feeding behavior and energetics of exposed animals. The methanol and phenol content of Grasselli waste may exert an inhibitory effect on feeding mechanisms, whereas other trace organic constituents may exert toxic effects on oxidative metabolism. Although changes in oxygen consumption rates and feeding rates as a result of waste exposure were similar for the three species tested, there were distinct differences in subsequent egg production rates of the three copepod species with Centropages being the most sensitive and Temora the least. These differences may be related to differences in energetic tactics and the content of energy reserves among the three species and waste exposure may actually mimic starvation stress in its effects on reproduction and metabolism. Estimates of zooplankton production rates were based on production to biomass ratios and instantaneous birth rates. For Centropages and Pseudocalanus, both indices were comparable, and thus both the immediate metabolic responses and the delayed reductions in fecundity of copepods exposed to industrial wastes may be used as sensitive indices of sublethal stress. The indices were not comparable, however, in studies with Temora; this suggests that populations of copepods that continue to allocate energy reserves to egg production during periods of reduced food availability may be better able to withstand waste exposure.

Presented at: 18th European Marine
Biology Symposium, Oslo, Norway,
August 14-20, 1983.

Supported by: NOAA, Office of Marine
Pollution Assessment, Grant NA79AA-
D-00027.

WHOI Contribution No. 5531.

THE IMPACT OF WASTE DISPOSAL IN NEAR SHORE ENVIRONMENTS

Judith M. Capuzzo, Wayne V. Burt,
Iver W. Duedall, Dana R. Kester
and P. Kilho Park

The distribution, fate, and effects of wastes in nearshore coastal environments are governed by the physical, chemical, and biological processes that reduce the concentration, alter the chemical form or bioavailability of wastes, and remove such contaminants from the water column. Various anthropogenic contaminants, such as dredged materials, sewage sludge, and industrial and municipal effluents, are dispersed, degraded and deposited through physical processes and biogeochemical cycles. Contaminants of biological or human health concern, such as pathogenic microorganisms, trace metals and organic contaminants, are primarily associated with particulate matter and their transport coincides with the transport of natural sediments. Environmental concern with waste input to nearshore environments is focused on the accumulation of contaminants, both chemicals and pathogens, in commercial resources and the degradation of ecosystems through both nutrient enrichment and contaminant accumulation.

In press: Wastes in the Ocean, Vol. 6:
Near Shore Waste Disposal. B. H.
Ketchum, J. M. Capuzzo, W. V. Burt,
I. W. Duedall, P. K. Park and D. R.
Kester, eds. Wiley-Interscience,
New York.

Supported by: NOAA, Office of Marine
Pollution Assessment, Grant NA 79AA-
D-00027.

THE EFFECTS OF PETROLEUM HYDROCARBONS ON LIPID METABOLISM AND ENERGETICS OF LARVAL DEVELOPMENT AND METAMORPHOSIS IN THE AMERICAN LOBSTER (HOMARUS AMERICANUS MILNE EDWARDS)

Judith M. Capuzzo, Bruce A. Lancaster
and Glenn C. Sasaki

In studies with larval stages of the
American lobster (Homarus americanus) ex-
posed to sublethal concentration of South
Louisiana crude oil, the normal patterns of

lipid storage, utilization, and synthesis
during larval development and metamorphosis
were altered with oil exposure. Increased
rates of protein catabolism, delayed molting
and reduced growth were also evident among
oil-exposed lobster larvae and suggest
either deficiency or immobilization of lipid
reserves. From analyses of lipid class com-
position of control and oil-exposed lobster
larvae, it is evident that oil-exposed lar-
vae have consistently lower levels of tri-
acylglycerols (the major energy store) and
higher levels of sterols than control lar-
vae. Decreased ratios of non-essential
fatty acids (both saturated and unsaturated)
and increased ratios of essential fatty
acids were detected in the neutral lipid
pools of oil-exposed larvae that suggest
both decreased storage of fatty acids in
energy reserves and decreased mobilization
of essential fatty acids into phospholipid
pools. The fatty acid distribution in the
phospholipid pools of oil-exposed larvae,
however, varied little from control values
and reflect a tendency to preserve the
integrity of these structural lipids. In-
creases in sterol levels may be related to
decreased mobilization of cholesterol or
other sterols and/or transformation of
cholesterol into metabolic pathways includ-
ing the biosynthesis of ecdysterone.

Oil-exposed animals contained trace
quantities of benzene, thiophene, toluene,
alkylcyclohexane and alkylbenzenes, detected
with GC-MS analyses. No naphthalene,
dimethylnaphthalene, higher molecular weight
aromatics or their phenolic derivatives were
detected; their absence may be related to
either the detection limits of the
methodology used or extremely rapid
metabolism and turnover of these components.
Alterations in lipid metabolism may account
for the developmental and energetic
abnormalities observed in marine crusta-
ceans. It cannot be ruled out, however,
that decreased lipid utilization, synthesis,
and mobilization may be a defense mechanism
against incorporating lipophilic components
of petroleum hydrocarbons in metabolic path-
ways and disruption in energetics and
development is a consequence of the reduc-
tion in energy available for growth and
molting.

In press: Marine Environmental Research.
Presented at: Second International
Symposium on Responses of Marine
Organisms to Pollutants, Woods
Hole, MA, April 27-29, 1983.

Supported by: U.S. Dept. of the
Interior, BLM Contract AA551-CT9-5,
and NOAA, Office of Sea Grant,
Grant NA80-AA-D-00077 (R/P-5).

WHOI Contribution No. 5475.

EFFECTS OF EXPOSURE TO SUBLETHAL
CONCENTRATIONS OF CRUDE OIL ON THE
COPEPOD Centropages hamatus
II. ACTIVITY PATTERNS

Timothy J. Cowles

Female copepods of the species Centropages hamatus exposed to 80 ppb South Louisiana crude oil in seawater show alterations in their swimming activity, as expressed by the duration of pause intervals during swimming, mean swimming speed, and allocation of time to active food search. Behavioral patterns of oil-exposed copepods returned to patterns shown by control animals after only 30 minutes of depuration in uncontaminated seawater. The results indicate that sublethal concentrations of hydrocarbons interfere with mechanisms of food perception in copepods, as in decapod Crustacea, either via blockage of receptors or via narcotization of the animal's nervous system.

Published in: Marine Biology, 78, 53-57.

Supported by: NOAA Contract
NA80RAD00037.

WHOI Contribution No. 5362.

EFFECTS OF EXPOSURE TO SUBLETHAL
CONCENTRATIONS OF CRUDE OIL ON THE
COPEPOD Centropages hamatus
I. FEEDING AND EGG PRODUCTION

Timothy J. Cowles and
James F. Remillard

Female copepods of the species Centropages hamatus show decreased ingestion rates and decreased egg viability when exposed to crude oil/seawater dispersions having crude oil concentrations of 10-80 ppb. However, rates of egg production were not significantly affected by these exposure levels. In addition, we found no evidence for accumulation of petroleum hydrocarbons by copepods exposed to 200 ppb of South Louisiana crude oil. The results imply that biosynthetic pathways involved in oogenesis may be influenced by sublethal concentrations of crude oil or that petroleum hydrocarbons directly affect the viability of eggs. Recruitment into field populations of copepods could be severely reduced as a consequence of exposure to low levels of physically dispersed crude oil.

Published in: Marine Biology, 78, 45-51.

Supported by: NOAA Contract
NA80RAD00037.

WHOI Contribution No. 5361.

PHYSIOLOGICAL AND BIOCHEMICAL EFFECTS OF
THE POLYCYCLIC AROMATIC HYDROCARBON
BENZ(α)ANTHRACENE ON THE DEPOSIT
FEEDING POLYCHAETE Nereis virens

Anne E. McElroy

Physiological and biochemical effects of short-term exposure to sediment-sorbed benz(α)anthracene (BA) on Nereis virens, and in vivo BA metabolism were investigated using flow-through benthic chambers and ¹⁴C tracer techniques. Worms exposed to sediments containing 14 ppm (dry wt.) BA for six days accumulated 4.9 ppm (dry wt.) BA whereas no detectable decrease in the sediment concentration of BA was observed. Little or no ¹⁴C activity was found either in organic extracts of water samples or in carbon dioxide stripped from water samples. Analysis by high pressure liquid chromatography revealed substantial metabolism of BA in worms but not in sediments. In worm tissue extracts, 37% of total ¹⁴C activity remained unmetabolized, 13% appeared as primary metabolites, 47% as conjugated metabolites and 2% was not solvent extractable. Sediment-sorbed BA had no measurable effects on whole animal oxygen consumption or ammonia excretion. Although there was no apparent change in aryl hydrocarbon hydroxylase activity, significant increases in the ADP/ATP ratio in exposed individuals were observed.

Presented at: The 18th European Marine
Biology Symposium, Oslo, Norway,
August 14-20, 1983.

Supported by: NOAA, Grant 83-ABD-00012;
Andrew W. Mellon Foundation, to
WHOI/CRC; OIP, National Wildlife
Federation/American Petroleum
Institute Environmental Conservation
Fellowship Program; and MIT/WHOI
Joint Program in Oceanography.

WHOI Contribution No. 5506.

EFFECTS OF DRILLING FLUIDS ON LARVAE OF
THE AMERICAN LOBSTER, Homarus americanus

Jennifer G. Smith Derby
and Judith M. Capuzzo

Laboratory experiments were conducted to evaluate the impact of oil drilling operations on the American lobster (Homarus americanus). The lethal and sublethal effects of used, whole drilling fluids on the larval stages of the lobster were assessed in continuous-flow bioassay experiments. The five drilling fluids tested were shown to vary markedly in their toxicity, with LC₅₀ values ranging from 74 mg/L to 5000 mg/L. Sublethal exposures to drilling

fluids resulted in reduced respiration rates and O:N ratios and increased protein:lipid ratios, demonstrating a change in energetics of the larval lobsters. An additional serious consequence of the drilling fluid exposure was the impairment of growth and development of the larvae.

Exposure of larvae to barite (a major component of drilling fluids) and to a field-collected fine-grained sediment did not cause the effects seen with exposure to drilling fluid, indicating that the chemical components and not the physical properties of the drilling fluids were primarily responsible for the detrimental effects observed. Chemical analyses of the tested drilling fluids identified some of these potentially toxic compounds. Drilling fluids containing a diesel oil component proved to be more toxic than those without this component. Other toxic components in these drilling fluids included phenolic compounds and various metals.

In press: Canadian Journal of Fisheries and Aquatic Sciences.

Supported by: U.S. EPA Grant
CR807447010.

WHOI Contribution No. 5378.

OIL SPILL STUDIES: A REVIEW OF ECOLOGICAL EFFECTS

John M. Teal and Robert W. Howarth

We reviewed seven particularly well known and/or studied oil spills that have occurred since the National Academy of Sciences 1975 report, "Petroleum in the Marine Environment," or that occurred prior to that report but about which significant new information has since been acquired. The spills studied were from the barge FLORIDA, and tankers ARROW, ARGO MERCHANT, AMOCO CADIZ, and TSESIS and blowouts from the Bravo and Ixtoc I platforms.

These "best" studies yield only limited insight into effects because they lack controls and have a high degree of natural variability. The TSESIS, FLORIDA, and AMOCO CADIZ cases are exceptional since they occurred in areas of ongoing research programs and had nearby areas suitable for controls. Oil spills have produced measurable effects on ecosystems that have not been readily predictable from laboratory studies on isolated organisms. However, ecosystem-level interactions are poorly understood even without the complications resulting from effects of pollution. These generalizations emerge: oil regularly reaches sediments after a spill; oil in anoxic sediments is persistent; oil regularly contaminates zooplankton and benthic invertebrates; fish

are also contaminated, but to a lesser extent; oil contamination decreases the abundance and diversity of benthic communities.

In press: Environmental Management,
Vol. 7.

Supported by: Ford and Sherman
Foundations.

WHOI Contribution No. 5370.

UNDERSTANDING SUBLETHAL POLLUTANT EFFECTS IN THE MARINE ENVIRONMENT

John Vandermeulen and Judith M. Capuzzo

Responses of marine organisms to waste inputs can be manifested at four levels of biological organization: (1) biochemical and cellular; (2) organismal, including the integration of physiological and behavioral responses; (3) population, including alterations in population dynamics; and (4) community, resulting in alterations in community structure and dynamics. The interrelationships of these responses are presented in Table 1. All responses are not disruptive in nature and do not necessarily result in degeneration at the next level of biological organization. Only when the compensatory or adaptive mechanisms at one level begin to fail, do deleterious effects become apparent at the next level. For predictive purposes, one must be aware of the early warning signs of stress at each level of organization before compensatory mechanisms are surpassed. From the biochemical level to the community level, the degree of system complexity, the number of compensatory mechanisms available and the lag time to measure a response increase exponentially, thereby increasing the predictive difficulties at each level.

GESAMP has defined "pollution" as the concentration of a contaminant that causes some measurable impact at the population level (GESAMP, 1980). Induced perturbations in larval development, growth, or histopathological changes would be acceptable as long as the total population size of a species was not impaired. With the exception of populations of seabirds and a few well studied fish stocks, our ability to predict subtle changes in the populations of marine animals is severely limited. Further, we suggest that once changes have been observed at the population level, several compensatory mechanisms have already been surpassed. To understand the extent of pollution effects in the marine environment, we must understand the full range of adaptive and disruptive responses available at each level of biological organization and the importance of these responses in deter-

mining the structure and function of marine ecosystems.

In press: Management of Wastes in the Ocean. M. Champ and M. Trainor, eds. Wiley-Interscience, New York.

Supported by: NOAA, Office of Marine Pollution Assessment, Grant 79AA-D-00027.

M I C R O B I O L O G Y

Chloroherpeton thalassion GEN. NOV.
OF SPEC. NOV., A NON-FILAMENTOUS, FLEXING
AND GLIDING GREEN SULFUR BACTERIUM

Jane Gibson, Norbert Pfennig
and John B. Waterbury

A novel green sulfur bacterium was isolated in pure culture and characterized. Its growth characteristics and requirements, cell morphology and fine structure, pigment composition, DNA base composition and ability to fix nitrogen were described. The name Chloroherpeton thalassion was proposed and a formal description of the bacterium was included.

In press: Archives of Microbiology, for publication in a memorial issue for Roger Y. Stanier.

Supported by: NSF Grant OCE80-25241.

MICROBES IN THE OCEANIC ENVIRONMENT

Holger W. Jannasch

An invited discussion on the last decade's progress in certain facets of marine microbiology (100th Anniversary of the British Society of General Microbiology).

Published in: The Microbe 1984.
Symposium No. 36 of the Soc. Gen. Microbiol. Cambridge Univ. Press, 97-122.

Supported by: NSF Grants OCE80-24253 and OCE81-17560.

WHOI Contribution No. 5387.

RECENT PROGRESS IN THE MICROBIOLOGY OF HYDROTHERMAL VENTS

Holger W. Jannasch and D. C. Nelson

A report on recent advances of in situ measurements of microbial processes and of pure culture isolations and their specific activities with emphasis on thermophilic bacteria and chemosynthetic symbionts of vent invertebrates.

In press: Current Perspectives in Microbial Ecology. C. A. Reddy and M. J. Klug, eds. Amer. Soc. Microbiol. Publ., Washington.

Supported by: NSF Grants OCE80-24253 and OCE81-17560.

WHOI Contribution No. 5463.

DEEP SEA MICROBIOLOGY

Holger W. Jannasch and Craig D. Taylor

A detailed review of ecological, physiological and biochemical aspects of the effects of deep sea temperatures and pressures on microorganisms, covering research of approximately the last decade.

In press: Annual Reviews of Microbiology 38. L. N. Ornstron, ed. Palo Alto, 1984.

Supported by: Mellon Senior Study Award of the WHOI; NSF Grants OCE80-24253 and OCE82-14896.

WHOI Contribution No. 5557.

CHEMOAUTOTROPHIC GROWTH OF A MARINE Beggiatoa IN SULFIDE-GRADIENT CULTURES

Douglas C. Nelson and Holger W. Jannasch

A recently isolated strain of Beggiatoa, MS-81-6 (c.f. alba), was tested for chemoautotrophic growth in semi-solid (0.2 percent agar) sulfide/oxygen gradient cultures. The organism grew in a horizontal layer, the distance from the air/medium interface depending on sulfide concentrations and changing with time. Optimal growth as a gradient organism was based on a preference for reduced oxygen concentrations and a limited sulfide tolerance in combination with gliding motility. In gradient cultures chemoautotrophic growth was demonstrated by the following criteria: (1) biomass yield (protein) increased with increasing sulfide concentration, and estimated molar growth yields agreed with those for other sulfide-grown chemoautotrophs; (2) approximately

90% of total cell carbon and protein carbon were fixed from carbon dioxide; (3) the CO₂-fixing enzyme, RuBP-carboxylase, was present in cell-free extracts at a level typical of chemoautotrophs; and (4) acidification of the medium, apparently linked to utilization of internal S⁰ granules, accompanied the later phase of growth. The ability to grow on acetate in the absence of a source of reduced sulfur renders the organism facultatively chemoautotrophic.

In press: Archives of Microbiology.

Supported by: NSF Grants OCE80-24253 and PDF81-66059.

WHOI Contribution No. 5386.

CELLULOLYTIC NITROGEN-FIXING BACTERIUM
CULTURED FROM THE GLAND OF DESHAYES
IN SHIPWORMS (BIVALVIA: TEREDINIDAE)

John B. Waterbury, C. Bradford Calloway
and Ruth D. Turner

A novel bacterium has been isolated in pure culture from the gland of Deshayes in six species of teredinid bivalves. This bacterium is interesting in several respects: it is the first bacterium known to both digest cellulose and fix nitrogen, two physiological traits that genetic engineers have tried unsuccessfully to combine in a single bacterium, and it is a principal participant in a unique symbiotic relationship, that may be responsible for the ability of shipworms to use wood as their sole food source.

Published in: Science, 221, 1401-1403.

Supported by: ONR Contract N00014-76-C-D281, NR104-687 and NSF Grant OCE80-25241.

WHOI Contribution No. 5338.

IN SITU STUDIES ON DEEP-SEA AMPHIPODS
AND THEIR INTESTINAL MICROFLORA

Carl O. Wirsen and Holger W. Jannasch

Amphipods, trapped at depths of 1410 to 5330 meters and incubated in situ in the presence of a radiolabeled food source, were fractionated for measuring the distribution of label in the major biochemical constituents: low molecular pools, ethanol and base soluble proteins, lipids, polysaccharides and nucleic acids. The results suggest lipid energy storage and enhanced microbial growth within the amphipod guts as compared to growth on the uningested food material. The merits and

limitations of this general approach are critically evaluated.

In press: Marine Biology.

Supported by: NSF Grants OCE79-19178
and OCE82-17560; ONR Contract
N00014-79-C0071 NR 083-004.

WHOI Contribution No. 5464.

M O D E L I N G

AN INJUNCTIVE FORM OF THE AXIOMS
OF TRIALECTICS

Hal Caswell

Most discussions of such philosophical systems as trialectics take an ontological approach to their subject. The axioms of the system are taken as descriptions of the way things are; as such they are clearly metaphysical, unfalsifiable, and, in general, unpalatable to scientists with a natural tendency toward a logical positivistic dislike of metaphysics. However, it is now recognized that frankly metaphysical principles (paradigms, themata, metaphysical research programmes, etc.) can and do play important roles in directing scientific investigation. Such principles play an essentially injunctive, rather than a descriptive role, distinguishing important from trivial problems, suggesting approaches to those problems, and defining the criteria which identify successful solutions to those problems. In this paper, the axioms of trialectics are restated in injunctive form and contrasted with injunctive versions of the axioms of formal logic and dialectics. It is shown that such a restatement emphasizes their possible role in directing scientific research and fundamentally changes the grounds for their critical discussion.

Published in: Trialectics: Toward a
Practical Logic of Unity. R. E.
Horn, ed. Information Resources,
Lexington, MA, 1983, 41-46.

Supported by: NSF Grant OCE76-19278.

WHOI Contribution No. 5509.

TRIALECTICS, CYBERNETICS, AND ZADEH'S
THEORY OF STATE

Hal Caswell

"Trialectic logic" has been proposed by
Oscar Ichazo (1976, 1982) as a metaphysical

framework within which to study the unity underlying the diverse and continually changing processes of nature. Trialectics can be contrasted with two other major systems: the formal logic of Aristotle and the dialectic of Hegel and Marx. The goal of this paper is to examine some aspects of cybernetics and general system theory and to compare their principles with those of trialectics. It appears that they are convergent approaches to the study of dynamic processes. The results shed light on refinements of trialectic analysis. Examples are presented, drawn from the ecology of predator-prey interactions and from bioenergetics.

Published in: Trialectics: Toward a Practical Logic of Unity.
R. E. Horn, ed. Information Resources, Lexington, MA, 1983, 157-190.

Supported by: NSF Grant OCE76-19278.

WHOI Contribution No. 5436.

ZADEH'S THEORY OF STATE AND ICHAZO'S
TRIALECTIC ANALYSIS: CONVERGENT
APPROACHES TO UNDERSTANDING DYNAMICS

Hal Caswell

The prototypical scientific problem is that of understanding the way in which system behavior unfolds through time. Mathematical models designed to describe such dynamics are typically formulated as differential or difference equations. Zadeh's behavioral theory of state revealed the factors underlying this approach, formalized the idea that the state of a system summarizes its history at a given instant, and provided a set of criteria to aid in the identification of state variables. Quite independently, Ichazo proposed a metaphysical system, called trialectic logic, as a framework for understanding process. His system includes a procedure for describing dynamics which is isomorphic to Zadeh's formulation of state space analysis. While Zadeh presented his theory in the context of engineering systems analysis, Ichazo's approach is part of a much more general psychodynamic system, and is designed to provide an experimental appreciation of dynamics in the context of individual problem solving.

Published in: Proceedings of the Society for General Systems Research, 27, 481-485.

Supported by: NSF Grant OCE76-19278.

WHOI Contribution No. 5346.

PHYTOPLANKTON

COPPER COMPLEXATION DURING SPRING
PHYTOPLANKTON BLOOMS IN COASTAL WATERS

Donald M. Anderson, John S. Lively
and Ralph F. Vaccaro

Cupric ion bioassays were conducted throughout the spring phytoplankton bloom season at two stations - one estuarine, dominated by dinoflagellates, the other coastal, dominated by diatoms. Copper-complexing ligands were detected at both locations throughout this period. Ligand concentrations varied between 0.1 and 0.75 μM , with the estuarine concentrations typically 2-4 times higher than coastal values. Ligands from both locations were destroyed by UV-oxidation and had similar conditional stability constants (range $10^{8.3}$ to $10^{9.2}$) that were significantly correlated with pH, suggesting that the complexing material may be organic with weak acid functional groups. All measured or calculated parameters (DOC, ligand concentration, total copper concentration, salinity, and pH) remained relatively constant at the coastal station through time. The estuarine station was more dynamic, with DOC, total copper, and ligand concentrations varying 2-4 fold during the study. Although ligand concentrations were significantly different between the two locations, concomitant fluctuations in total dissolved copper and ligand strength resulted in a relatively constant estimate of the maximum free copper activity at both stations (near 10^{-11} M). This suggests that copper toxicity alone was not responsible for the distinctly different estuarine and nearshore phytoplankton assemblages, although sensitive species might have been inhibited at both locations. Major phytoplankton blooms at both sites were not accompanied by changes in DOC or complexation capacity. A significant inverse correlation between ligand concentration and salinity suggests a terrestrial or sedimentary origin for the copper-complexing compounds.

Supported by: NOAA, Office of Sea Grant, Grant NA80AA-D-00077 (R/B-41); NOAA, Ocean Assessments Division, Grant NA81RA-D-00012; and International Copper Research Association, Project No. 307.

WHOI Contribution No. 5318.

PRODUCTION AND ROLE OF HYALINE HAIRS IN
Ceramium rubrum

J. A. DeBoer and F. G. Whoriskey

Light intensity within the range of 40 to $210 \mu\text{E}\cdot\text{m}^{-2}\cdot\text{sec}^{-1}$ had relatively little effect on the abundance, length or distribution of hairs. External NH_4^+ concentrations less than $0.5 \mu\text{M}$ stimulated hair initiation and growth in apical regions of the thalli. Ammonium concentrations in excess of $20 \mu\text{M}$ inhibited hair formation on all regions of the thalli.

Ammonium uptake velocities of plants with hairs were approximately twice those of plants without hairs. These hairs may be an adaptive strategy to take advantage of intermittent bursts of nutrients by increasing the plant's surface area.

In press: Marine Biology.

Supported by: U.S. Energy Research and Development Administration, Contract E(11-1)-2948, and the Jessie Smith Noyes Foundation, Inc.

WHOI Contribution No. 5526.

UPTAKE AND ASSIMILATION OF AMMONIUM AND
NITRATE BY PHYTOPLANKTON:
INDICES OF NUTRITIONAL STATUS FOR
NATURAL ASSEMBLAGES

Patricia M. Glibert and
James J. McCarthy

Using direct measurements of nitrogen uptake by natural phytoplankton assemblages from the Chesapeake Bay, the Caribbean and Sargasso Seas, and a warm core Gulf Stream ring, we have examined the usefulness of some parameters as potential indices of cellular nutritional status. In different locations, as nutrient and biomass conditions permitted, we applied a variety of approaches, including the ratio of uptake of NH_4^+ and NO_3^- at high and low concentrations of these substrates, the rate of assimilation of NH_4^+ or NO_3^- into protein, and NH_4^+ enhancement of dark carbon fixation. Generalizations concerning the nutritional status of an assemblage were difficult to derive from any single index, but using a variety of approaches, we were able to demonstrate more severe nutritional stress for Caribbean Sea assemblages when NH_4^+ concentrations were $<0.3 \mu\text{g atom l}^{-1}$ than for Sargasso Sea assemblages when NH_4^+ concentrations were at least an order of magnitude lower. During the decline of phytoplankton blooms in both the Chesapeake Bay and a warm core Gulf Stream ring fairly severe nutritional stress was also demonstrated.

Supported by: NSF Grants DES75-14781, OCE77-26401, OCE78-26011, OCE80-22990 and OCE80-24445; awards from the Leopold Schepp Foundation and the WHOI Education Office.

WHOI Contribution No. 5493.

CONCEPTUAL ROLE FOR MICROAGGREGATES
IN PELAGIC WATERS

Joel C. Goldman

Organic microaggregates ($<100 \mu\text{m}$ in length) formed both by physical-chemical and biologically-mediated degradation processes in surface waters of nutrient-impovertished environments may play a vital role in the cycling of nutrients and organic carbon within microbial food chains. Based on recent evidence, it has become abundantly clear that small (up to several μm in diameter) flagellated autotrophs and phagotrophic grazers along with bacteria are among the dominant organisms in oligotrophic waters and that most of the nutrients that sustain these microbes are recycled. Colonization of recently formed microaggregates by bacteria, algae, and phagotrophic microflagellates through various types of association is one way that nutrients could be absorbed from the aqueous environment and concentrated against a gradient. Flagellates, because of their rapid swimming speeds, most likely can migrate among aggregates even though the density of aggregates may be low. Although swimming does little to increase the flux of nutrients to small flagellates, its main function may be to allow these microbes to seek out, through chemotaxis, enriched microenvironments located within aggregates. The physical and biochemical environment within microaggregates, undoubtedly, is very much different than what is perceived from measurements made on large-volume samples.

In press: Bulletin of Marine Research.

Supported by: NSF Grant OCE83-8578.

WHOI Contribution No. 5584.

INFLUENCE OF BOTTLE PHOTOINHIBITION ON
THE MEASUREMENT OF SEASONAL PRIMARY
PRODUCTION IN A TEMPORATE COASTAL WATER

Joel C. Goldman and Mark R. Dennett

Primary productivity in Vineyard Sound, Massachusetts was determined at monthly intervals during May - October, 1981. Over the course of each study day fresh samples first were obtained at three-hr intervals and then incubated until sunset with

time-course measurements of $\text{H}^{14}\text{CO}_3^-$ uptake made at frequent intervals. Two seasonal patterns of ^{14}C uptake were apparent. First, during the spring and fall rapid initial uptake of ^{14}C over each incubation was followed by reduced uptake during periods of strong sunlight. The duration over which rapid uptake was sustained was a function of the prevailing light intensity at the start of the incubation. Under these conditions, estimates of near surface primary productivity based on summing total production over the first three hrs of each back-to-back incubation were higher by 50-130% than estimates based on single end-point measurements from full-day incubations. During the summer, however, when both total daily irradiation and temperature were maximal, uptake of ^{14}C was linear throughout each incubation so that the two techniques for estimating productivity led to comparable results. The reasons for these seasonal differences in the pattern of ^{14}C uptake are unclear, but may be related to changes in temperature or phytoplankton species, or both. Our results are in general accord with the conclusion of Harris (1980) that photoinhibition may be far more pronounced during a bottle incubation than is occurring in the water column if vertical mixing is sufficient to prevent prolonged exposure of phytoplankton to photoinhibiting light intensities at the surface.

In press: Marine Ecology Progress Series.

Supported by: NSF Grant OCE80-24445.

WHOI Contribution No. 5437.

D I N O F L A G E L L A T E S

THE IMPORTANCE OF DORMANT CYSTS IN TOXIC DINOFLLAGELLATE BLOOMS AND SHELLFISH TOXICITY

Donald M. Anderson

Several of the dinoflagellates responsible for toxic blooms include a dormant cyst stage in their life cycles. These resistant cells have a variety of potential functions in the overall ecology of the toxic dinoflagellates. As Wall (1) discussed, cysts can theoretically act as "seed" populations to initiate blooms, as a survival mechanism through environmental extremes, as agents for species dispersal, and as means for genetic recombination through sexuality. Two additional functions can be added to this list: cysts can be

direct sources of toxicity, and their formation can be a major factor in bloom termination. The purpose of this paper is to examine our present knowledge of the cysts of the toxic Gonyaulax species to see whether these hypothetical functions are valid representations of the actual roles cysts play in toxic dinoflagellate blooms and shellfish toxicity.

Supported by: NOAA, Office of Sea Grant, Grant NA80AA-D-00077 (R/B-41) and NOAA, Ocean Assessments Division, Office of Oceanography and Marine Services, Grant NA81RAD00012.

WHOI Contribution No. 5541.

SEXUALITY AND CYST FORMATION IN THE DINOFLLAGELLATE Gonyaulax tamarensis: I. CYST YIELD IN BATCH CULTURES

Donald M. Anderson, David M. Kulis
and Brian J. Binder

Encystment of the toxic dinoflagellate Gonyaulax tamarensis Lebour was monitored in batch cultures exposed to a variety of nutritional and environmental treatments. Limitation by nitrogen (as NH_4 or NO_3) or phosphorus (as PO_4) resulted in cyst formation. When the initial concentration of limiting nutrient was varied, total cyst yield (mL^{-1}) was directly proportional to the peak motile population size at all but the highest nutrient concentrations (where encystment was minimal). Thus encystment efficiency was relatively constant ($0.1 - 0.2$ cysts cell^{-1}) over a range of cell densities. Since two motile cells fuse to produce each cyst, these efficiencies imply that 20 to 40% of the vegetative populations successfully encysted. Nutrient measurements during nitrogen-limited cyst formation suggest that sexuality occurred before internal nitrogen pools reached subsistence levels.

Cyst formation was negligible in nutrient-replete medium, even with a 50% reduction in growth rate due to non-optimal light. Light did alter cyst yield once encystment was initiated under nutrient limitation, however, but this was probably linked to smaller motile population sizes and not to a specific inhibition of encystment. In contrast, encystment was more sensitive to temperature than was growth rate, with optimal cyst production occurring over a relatively narrow temperature range and no encystment at some temperatures that permitted growth. Cyst yield could also be reduced by contaminants introduced during culture medium preparation. Crosses of most G. tamarensis isolates (including ordered tetrads and other presumably suitable mating types) produced virtually no cysts under

conditions known to favor encystment in the one strain used throughout this study. This may reflect genetic variability or the effects of bacteria or fungi as contaminants.

Supported by: NOAA, Office of Sea Grant,
Grant NA80AA-D-00077 (R/B-41) and
NSF Grant OCE82-08739.

WHOI Contribution No. 5523.

SEXUALITY AND CYST FORMATION IN THE
DINOFLAGELLATE *Gonyaulax tamarensis*:

II. TIME-COURSE MEASUREMENTS OF
PHOSPHORUS LIMITATION

Donald M. Anderson and Niels L. Lindquist

Intracellular and extracellular phosphorus concentrations were monitored at twice-daily intervals throughout the transition from asexual to sexual reproduction in phosphorus-limited, multi-tube batch cultures of the toxic dinoflagellate *Gonyaulax tamarensis* Lebour. Approximately 20% of all motile cells successfully completed sexuality through the formation of resting cysts (hypnozygotes). It was not possible to pinpoint a precise time when asexual division ceased and the sexual fusion of gametes began, but changes in the population's mean cell diameter, changes in cellular phosphorus content, and an apparent surge in division rate in the late stages of exponential growth, together suggest that sexuality was induced in response to actual or imminent nutrient limitation. Planozygotes (the swimming precursors to cysts) increased in abundance over a 4-day interval, followed 5-6 days later by a parallel increase in cyst numbers. These gradual increases indicate that gamete formation and/or fusion did not occur in a sudden, synchronized fashion, or alternatively that the duration of planozygote development varied as the batch cultures aged. A steady decrease in the phosphorus cell quota during this experiment indicates that *G. tamarensis* growth may exceed uptake at 20°C with external PO_4 concentrations as high as 3.5 μM . Encystment in natural waters may thus occur due to nutrient limitation at concentrations well above analytical detection limits.

Supported by: NSF Grant OCE82-08739 and
the Education Program at the Woods
Hole Oceanographic Institution.

WHOI Contribution No. 5524.

SEXUAL PROCESSES IN THE LIFE CYCLE OF
THE BLOOM FORMING DINOFLAGELLATE
Gyrodinium uncatenum:
A MORPHOGENETIC OVERVIEW

D. Wayne Coats, Mary A. Tyler
and Donald M. Anderson

Sexual processes in the life cycle of the dinoflagellate *Gyrodinium uncatenum* were investigated in manipulated field populations. Morphological and morphogenetic aspects of gamete production, planozygote formation, encystment, excystment, and planomeiocyte division are described from observations of living specimens, protargol silver impregnated material and scanning electron microscope preparations.

The sexual cycle is initiated by gamete formation which involves two asexual divisions of the vegetative organism. Gametes are fully differentiated following the second division and immediately capable of forming pairs. Either isogamous or anisogamous pairs are formed by the mid-ventral union of gametes. Gametes invariably join with flagellar bases in close juxtaposition. Complete fusion of gametes requires ~1 hr and involves plasmogamy followed by karyogamy, resulting in a quadriflagellated planozygote. Planozygotes encyst in 24-48 hrs to yield a hypnozygote capable of overwintering in estuarine sediments. After several weeks of dormancy under winter conditions, hypnozygotes readily excyst upon exposure to temperatures above 15°C. A single quadriflagellated planomeiocyte emerges from the cyst and under culture conditions, divides 1-2 days later. The four flagella are not evenly distributed at the first division and both bi- and tri-flagellated daughter cells are formed.

Supported by: NSF Grant OCE80-11039.

WHOI Contribution No. 5474.

P O P U L A T I O N E C O L O G Y

OPTIMAL LIFE HISTORIES AND AGE-SPECIFIC
COSTS OF REPRODUCTION: TWO EXTENSIONS

Hal Caswell

A recent paper in this journal (Caswell 1982a) derived a prediction for the age-specific costs of reproduction, conditional on the hypothesis of life history optimality. Costs of reproduction, assessed in terms of survival, were predicted to vary with age as the inverse of the reproductive value distribution and, thus, to increase markedly with age in most species. Costs,

assessed in terms of future parental fecundity, were predicted to vary as the ratio of successive terms in the stable age distribution; for many species this ratio is nearly age invariant. Both these predictions were supported by a variety of circumstantial evidence. However, the calculations were carried out separately for the two cases. The result of the simultaneous operation of both types of cost was left as an open problem. In this note, I extend the approach to consider constraint functions involving arbitrary simultaneous combinations of demographic parameters and show that the results obtained previously still hold. In addition, the present results are not restricted to the age-classified life cycle analyzed in the earlier paper, but can be applied directly to constrained optimization in complex life cycles.

Supported by: NSF Grants OCE76-19278 and BSR82-14583.

WHOI Contribution No. 5445.

EVALUATING THE CONSEQUENCES OF REPRODUCTION IN COMPLEX SALMONID LIFE CYCLES

Hal Caswell, Robert J. Naiman
and Rod Morin

Reproduction by individuals at different stages in the life cycle contributes differentially to population growth. This paper shows how to evaluate these contributions, using demographic analyses developed for the analysis of complex life cycles in which individuals may be classified by size or developmental stage as well as age. Population growth is measured by the maximum eigenvalue of a projection matrix corresponding to a life cycle graph. The sensitivity of this eigenvalue to changes in reproduction can be expressed as a simple function of the demographic parameters describing the life cycle. The sensitivity of growth rate to reproduction declines with age, at a rate which is inversely proportional to survival and directly proportional to growth rate itself. The importance of alternate developmental pathways is related to the reproductive values of the stages to which those pathways lead. The latter observation is related to the recent increase in precocious maturation by male parr in the Atlantic salmon (Salmo salar).

Supported by: NOAA, Office of Sea Grant, Grant NA83-AA-D-00049 (R/B-50) and the WHOI Matamek Research Program.

WHOI Contribution No. 5542.

PLANKTONIC LARVAE ACT LIKE PASSIVE PARTICLES IN TURBULENT NEAR-BOTTOM FLOWS

Cheryl Ann Hannan

The hypothesis that infaunal polychaete larvae (Mediomastus ambiseta) fall through the water like passive particles in turbulent near-bottom flows was tested in the field using two geometrically different sediment traps. Laboratory flume experiments dictated the a priori prediction that one trap geometry would significantly over-collect larvae relative to the other trap design geometry. In field experiments, this a priori prediction was always supported; Mediomastus larvae are distributed in near-bottom waters according to the hydrodynamical processes governing the distribution of fine suspended sediments near the seabed. Because larvae respond to small-scale fluid flows and turbulence like passive particles, it is possible that larvae are also initially deposited on the seafloor like passive particles. Larvae may first reach the seabed at sites where sediments, with fall velocities similar to larvae, would settle.

In press: Limnology and Oceanography.

Supported by: Coastal Research Laboratory at WHOI; the Diving Equipment Manufacturers Association; Association for Women in Science Predoctoral Award; NSF Dissertation Improvement Grant; National Ocean Survey/Sea Grant Fellowship, NOAA, Grant NA80-AA-D-00077; and the WHOI Education Program.

WHOI Contribution No. 5334.

SEASONAL CHANGES IN THE DEPTH DISTRIBUTION OF BIVALVE LARVAE ON THE SOUTHERN NEW ENGLAND SHELF

Roger Mann

The seasonal change in occurrence, depth distribution, size distribution and species composition of bivalve larvae at a station on the Southern New England Shelf was examined for the period April-December 1981 and related to temperature structure of the water column and chlorophyll a distribution. Bivalve larvae were most abundant during late August and September at depths greater than 10 m in association with water temperatures of 14-18°C and chlorophyll a concentrations of <2µg/L., and at the surface in October in association with a temperature of 15.5°C and chlorophyll a concentrations of ~3.0 µg/L. Larvae in excess of 200 µm length consisted predominantly of the species Modiolus modiolus L., Arctica islandica L. and Spisula solidissima Dilwyn.

M. modiolus was present in the depth range 10-40 m from late July through December with highest concentrations in August through October. A. islandica was present in May and, at depths of 20-40 m from late July through November. A. islandica larvae captured in May possibly originate from spawnings in late 1980. A. islandica larvae captured in November are first shelled veligers of 110 μ m length. These may form the basis of an overwintering population of larvae. S. solidissima larvae were present from late July through October and extended into shallower, warmer waters than either M. modiolus or A. islandica.

Supported by: NOAA Sea Grant, Grant NA80-AA-D-00077; Office of Naval Research, Contract N00014-82-C-0071 NR 083-004; and the Andrew W. Mellon Foundation.

WHOI Contribution No. 5508.

SEASONAL CHANGES IN THE DEPTH DISTRIBUTION
OF LARVAE Arctica islandica L:
A COMPARISON OF FIELD DATA FROM THE
SOUTHERN NEW ENGLAND SHELF AND A
LABORATORY GENERATED MODEL

Roger Mann

The seasonal change depth distribution of Arctica islandica larvae at a station on the Southern New England Shelf for the period April-December 1981 is compared with depth distribution as predicted by a laboratory generated model. In the field larvae in excess of 200 μ m length were present in May and, at depths of 20-40 m from late July through November. The majority of larvae captured in November were first shelled veligers of 110 μ m length. Good agreement exists with respect to absence of A. islandica larvae in the warm (>18°C) shallow (0-20 m) waters between July and early September, and the abundance of larvae throughout the depth range 20-40 m from July through October. The predicted occurrence of larvae in June and July is not seen in the field and may be due to the combination of reduced spawning by adult A. islandica and less than optimum conditions for larvae development. The model predicts aggregation of larvae at the surface following decay of a seasonal thermocline; however, such aggregations are not seen in the field.

Supported by: NOAA Sea Grant, Grant NA80-AA-D-00077; Office of Naval Research, Contract N00014-82-C-0071 NR 083-004; and the Andrew W. Mellon Foundation.

RECRUITMENT OF COPEPOD NAUPLII INTO THE
PLANKTON: THE IMPORTANCE OF
DIAPAUSE EGGS AND BENTHIC PROCESSES

Nancy Marcus

During the last ten years it has been shown that the eggs of many zooplankton (e.g., copepods, cladocerans, and tintinnids) occur in the bottom sediments of coastal regions. Accumulations of these eggs on the sea-bottom may represent an important source for the recruitment of individuals into the planktonic population. We examined the vertical distribution and viability of copepod eggs at several locations in Buzzards Bay, Massachusetts. Cylindrical cores were obtained by divers. In the laboratory each core was divided at 1.0 cm intervals to a maximum depth of 10.0 cm. From each layer, two subcores were taken. One was suspended in sea water and incubated for several days at 19°C. The nauplii that hatched were collected, counted, and reared for identification. Viable eggs (evidenced by the appearance of nauplii) were found at all depths in the sediment. The greatest densities (10^6 - 10^7 /m²) often occurred several centimeters below the water-sediment interface. Egg abundance was not significantly correlated with grain size of the sediments. The other subcore was sonified so that eggs of Labidocera aestiva could be removed, counted, and incubated (to determine viability). The results of laboratory feeding experiments indicate that diapause and subitaneous eggs of Labidocera aestiva remain viable after being consumed by either Capitella sp. 1 or Streblospio benedicti which are common benthic consumers in regions where eggs occur. The relative importance and influence of physical factors (waves, tidal and wind driven currents) and biological factors (feeding and bioturbating activities of benthic animals) on the distribution, abundance, and viability of eggs on the sea-bottom is discussed.

In press: Marine Ecology Progress Series.

Supported by: NSF Grants OCE80-24440 and OCE83-14882; NOAA, Office of Sea Grant, Grant NA80-AA-D-00077.

WHOI Contribution No. 5364.

VARIATION IN THE DIAPAUSE RESPONSE OF
Labidocera aestiva (COPEPODA: CALANOIDA)
FROM DIFFERENT LATITUDES AND ITS
IMPORTANCE IN THE EVOLUTIONARY PROCESS

Nancy H. Marcus

Labidocera aestiva were collected from

three distantly separated sites along the Atlantic coast of the United States. Their offspring were reared in the laboratory under several combinations of photoperiod and temperature to assess the diapause response and genetic similarity of the populations. The results indicate that the L. aestiva is not genetically homogeneous throughout its range. Differences in the percentage of subitaneous and diapause eggs produced by laboratory-reared and field-collected populations reflected the unique combination of environmental conditions occurring in the geographic regions from which the animals originated. In the Virginian province where planktonic stages of the species are seasonally present, diapause egg production is triggered by short daylength photoperiods and cool temperatures. On the other hand, in the Carolinian province where the planktonic stages tend to occur year-round, diapause eggs are rarely produced by either field-collected females or their laboratory-reared offspring. This suggests that most of these animals lack the genetic capacity to produce diapause eggs. Diapause may lead to the reproductive isolation of populations if gene flow is reduced or prevented owing to their separation on a temporal scale.

In press: Biological Bulletin.

Supported by: NSF Grant OCE80-24440.

WHOI Contribution No. 5513.

THE INFLUENCE OF BEAVER
(Castor canadensis) ON THE PRODUCTION
DYNAMICS OF AQUATIC INSECTS

Robert J. Naiman, Donald M. McDowell
and Barbara S. Farr

Few studies directly address the environmental alterations of lotic ecosystems by animals. The retention of sediment and organic matter by beaver dams alter the character of the substrate and, supposedly, the nature and dynamics of the invertebrate community. Rocky substrate is covered by fine particulate matter as the water current is reduced, with community composition shifting towards a larger proportion of invertebrates adapted to soft bottom. The principle objective of our research is to evaluate the role of beaver as agents influencing the dynamics of aquatic ecosystems. The purpose of this paper is to present preliminary data on how beaver influence the diversity and production dynamics of aquatic insects emerging from beaver ponds in Québec, Canada, and to compare these data to those from unmodified sections of the same streams and larger rivers.

In press: Verh. Internat. Verein.
Limnol.

Supported by: NSF Grant DEB81-05677 and
the WHOI Matamek Research Program.

WHOI Contribution No. 5479.

ENERGY BUDGET OF AN OMNIVOROUS ROCKY SHORE
AMPHIPOD, Calliopius laeviusculus (KROYER)

Judith B. Pederson and Judith M. Capuzzo

An energy budget was constructed for Calliopius laeviusculus (Kroyer), an opportunistic omnivore. Diets of ephemeral algae and harpacticoids provided the greatest amount of energy, whereas perennial algae that were abundant in the field were poor diets. In laboratory experiments, C. laeviusculus assimilated about 64% of ingested algae or harpacticoids. After feeding, mean respiration rates increased from 0.325 to 0.450 $\mu\text{g-at O}_2\text{h}^{-1}\cdot\text{mg dry weight}^{-1}$ and ammonia excretion rates from 0.017 to 0.024 $\mu\text{g-at NH}_4\text{-N}\cdot\text{h}^{-1}\cdot\text{mg dry weight}^{-1}$. Mean O:N ratios were about 25 for both starved and fed amphipods, suggesting protein stores were used as an energy source, but not exclusively.

In press: Journal of Experimental Marine
Biology and Ecology.

Supported by: NOAA, Office of Sea Grant,
Grant NA80-AA-D-00077 (R/P-5).

WHOI Contribution No. 5416.

Z O O P L A N K T O N

A TECHNIQUE FOR THE ENUMERATION OF
HETEROTROPHIC AND PHOTOTROPHIC
NANOPLANKTON USING EPIFLUORESCENCE
MICROSCOPY, AND A COMPARISON WITH
OTHER PROCEDURES

David A. Caron

A new method is described using the fluorochrome primulin and epifluorescence microscopy for the enumeration of heterotrophic and phototrophic nanoplankton (2-20 μm). Phototrophic microorganisms are distinguished from heterotrophs by the red autofluorescence of chlorophyll a. Separate filter sets are used which allow visualization of the primulin-stained nanoplankton without masking chlorophyll a fluorescence, thus allowing easy recognition of phototrophic cells. Comparison with existing epifluorescence techniques for counting heterotrophic and phototrophic nanoplankton shows

that primulin provides more accurate counts of these populations than the FITC or pro-flavine techniques. Accuracy is comparable to the acridine orange technique, but requires only one filter preparation for the enumeration of both phototrophic and heterotrophic populations.

Published in: Applied and Environmental Microbiology, 46, 491-498.

Supported by: Joint Program in Biological Oceanography between the Massachusetts Institute of Technology and the Woods Hole Oceanographic Institution; OIP; NSF Grants OCE81-12991 and OCE82-14928.

WHOI Contribution No. 5380.

PREDICTED AND OBSERVED FEEDING RATES OF
THE SPINOSE PLANKTONIC FORAMINIFER
Globigerinoides sacculifer

David A. Caron and Allan W. H. Bé

Planktonic foraminifera feed by using a sticky rhizopodial network to ensnare motile prey. Rates of capture of zooplankton prey were estimated from the frequency of prey items observed in the rhizopodia of SCUBA-collected specimens of the planktonic foraminifera Globigerinoides sacculifer (Brady). Copepods were the dominant prey observed, constituting 44% of the foraminiferal prey numerically and 95% of the biomass. A simple model was derived to describe the natural rates of predation by G. sacculifer on zooplankton, based on non-selective capture of appropriate-sized prey. The model was applied to predict feeding rates of G. sacculifer on copepod assemblages over a range of values. The model predicted a wide variability (two orders of magnitude) in the rate of capture of copepods for the range of variables chosen. Feeding rates, estimated from the frequency of copepods in the rhizopodia of SCUBA-collected foraminifera, also showed a wide variability, averaging 1 copepod captured every 3.3 days, and were generally within the range of feeding rates predicted by the model.

In press: Bulletin of Marine Science.

Supported by: NSF Grants OCE78-25450 and OCE81-17715.

WHOI Contribution No. 5525.

GEOGRAPHIC, VERTICAL, SEASONAL AND DIEL
DISTRIBUTIONS OF PHOTOTROPHIC AND
HETEROTROPHIC COMPONENTS OF THE
PICOPLANKTON AND NANOPLANKTON IN THE
NORTH ATLANTIC

Paul G. Davis, David A. Caron,
Paul W. Johnson and John McN. Sieburth

Heterotrophic and phototrophic members of the nanoplankton (2-20 μm) and picoplankton (0.2-2.0 μm) were enumerated by epifluorescence microscopy from 39 estuarine and oceanic stations to assess the trophic relationship of apochlorotic nanoplankton with other components of the microbial plankton. Relationships between plankton fractions were found over geographic separation as well as seasonal and diel cycles. Population densities of heterotrophic nanoplankton (Hnano) and phototrophic nanoplankton (Pnano) were similar, ranging from 10^4 cells/ml in estuarine environments to 10^2 cells/ml at oceanic stations. All picoplankton and nanoplankton populations in the euphotic zone showed significant inverse linear relationships with log of bottom depth. Vertical profiles showed decreasing abundance of all populations with depth. Studies in Narragansett Bay, Rhode Island, indicated that Hnano, Pnano and heterotrophic picoplankton (Hpico) followed similar trends over winter/spring and summer blooms. Hnano was positively correlated with Hpico populations one week earlier, suggesting the relationship of prey (Hpico) and predator (Hnano). Diel studies at oceanic stations and in mesocosms in Narragansett Bay revealed positive correlations between Hpico and Hnano populations which may represent small interval temporal changes. These data suggest an abundant and dynamic Hnano population in the marine plankton which reflects changes observed in other components of the microbial plankton.

Supported by: NSF Grants OCE81-21881, OCE74-01537A02, OCE76-81779, OCE80-24441, OCE78-26388 and OCE80-18514; EPA Grant R-806072; and the WHOI Education Program.

COMPARISON OF BEHAVIORAL PATTERNS IN THE
COPEPODS Acartia tonsa AND
Centropages typicus: A QUANTITATIVE
VIDEO ANALYSIS

Timothy J. Cowles

Quantification of the activity patterns of Acartia tonsa and Centropages typicus, through videotape analysis, shows that these copepods alter their allocation of time between periods of swimming and periods of no movement (pauses) in response to exposure to different food conditions.

Both A. tonsa and C. typicus allocate less time to swimming (active food search and capture) when food is scarce than when food is abundant. As a consequence of these shifts in time allocation, A. tonsa and C. typicus both sample, per unit time, less of the fluid for food cues when food is scarce than when it is abundant. While both copepod species show the same pattern of responses, C. typicus spends more time in active search than does A. tonsa under all conditions, reflecting its more omnivorous trophic status. The results indicate that quantitative analysis of behavioral patterns will help answer questions about the modes of food perception and the energetics of food search in zooplankton.

Supported by: NSF Grant OCE80-17271 and NOAA Grant NA80RAD0037.

WHOI Contribution No. 5317.

THE DEVELOPMENTAL STAGES OF Longipedia americana (COPEPODA:HARPACTICOIDA)
REARED IN THE LABORATORY

Takashi Onbé

The complete postembryonic development of Longipedia americana (Copepoda: Harpacticoida) is described and illustrated based on specimens cultured in the laboratory. Development from the first nauplius to the adult took about 25 days at 20-21°C, when animals were fed a mixture of phytoplankton consisting of Isochrysis galbana, Thalassiosira weissflogii, and T. pseudonana.

Supported by: Fellowship from the Ministry of Education, Science, and Culture of Japan.

WHOI Contribution No. 5500.

THE FUNCTIONS OF NEMATOCYSTS IN PREY CAPTURE BY EPIPELAGIC SIPHONOPHORES (COELENTERATE, HYDROZOA)

Jennifer E. Purcell

The nematocysts of 24 siphonophore species were examined by light and scanning electron microscopy (SEM) for differences that could relate to differences in the sizes and types of prey captured. The siphonophore species in the suborder Calycophorae had 4-30 microbasic mastigophores from 0.7-18.0 µl in volume and 50-2,000 smaller homotrichous anisorhizas in uncoiled nematocyst batteries. The physonect siphonophore species had 4-120 stenoteles or microbasic mastigophores from 1.8-40.7 µl in volume and 150-20,500 smaller homotri-

chous anisorhizas in coiled nematocyst batteries. The sizes of crustacean prey (primarily copepods) captured by species in both suborders increased with increasing nematocyst size and numbers. Examination by SEM of captured, but uningested, prey showed that the heavily-spined threads of these nematocysts adhered to the prey surface and primarily entangled the prey. In contrast, the tentacles of cystonect siphonophores, including Physalia physalis, have isorhizas of 1.0-18.0 µl volume with and without small spines. These nematocysts penetrate the soft-bodied prey (mostly fish larvae) of these siphonophores, but apparently do not penetrate or entangle hard-bodied prey. Thus, prey capture by siphonophores differs with the sizes, numbers, and types of nematocysts present in each species. The possible functions of nematocyst batteries in tentacle spreading and luring of large zooplankton prey are discussed.

Supported by: WHOI Postdoctoral Fellowship; NOAA, Office of Sea Grant, Grant NA80-AA-D-00077 (R/B-44); and NSF Grant OCE81-24441.

WHOI Contribution No. 5495.

PARTICLE PRODUCTION BY PLANKTONIC CILIATES

Diane K. Stoecker

Favella sp. (a tintinnid) and Balanion sp. (a non-loricate ciliate) produce fecal aggregates which overlap in size with their dinoflagellate food. These aggregates have a higher C:N ratio than the dinoflagellates. As the aggregates disintegrate, smaller-sized particles are produced. For Favella grazing on the thecate dinoflagellate, Heterocapsa triquetra, the volume of fecal material produced is 22-23% of the volume of algae consumed.

The fecal aggregates are similar to fine, amorphous aggregates which are observed in the ocean. Ciliate fecal aggregates remain suspended and should contribute to microbial production and nutrient regeneration in the upper water column. In contrast, most macrozooplankton fecal pellets sink rapidly and transport particulate organic matter downward.

The similarity in size of microplanktonic algae and ciliate fecal aggregates makes the Coulter counter an inappropriate instrument for use in determining clearance or ingestion rates in ciliate feeding experiments unless egestion rates can be estimated. Visual counts were compared with Coulter counter data for an experiment in which Favella and Balanion grazed Heterocapsa. Use of the Coulter counter resulted in an underestimation of ingestion rates by 30%, for Favella, and by 18%, for Balanion, and

confounded the results in respect to particle size selectivity. However, because electronic particle counting measures the net change in particle concentration (ingestion - egestion) it can be used to measure assimilation rates. The assimilation efficiencies for Favella and Balanion eating Heterocapsa were thus estimated to be 70 and 82%, respectively.

Supported by: NSF Grant OCE82-00030.

WHOI Contribution No. 5348.

FINE SCALE SPATIAL CORRELATIONS BETWEEN PLANKTONIC CILIATES AND DINOFLAGELLATES

Diane K. Stoecker, Linda H. Davis
and Donald M. Anderson

Fine scale spatial distributions of planktonic ciliates which prey on dinoflagellates were investigated in a small estuary. The horizontal distributions of Favella sp., Balanion sp., and Strobilidium sp. were positively correlated with the patchy distribution of dinoflagellates. The vertical distribution of Favella was positively correlated with the distribution of dinoflagellates. When dinoflagellate density was statistically held constant distribution of Favella was not related to temperature or salinity. Throughout the diurnal cycle, the vertical distribution of Favella was similar to the vertical distribution of dinoflagellates.

We speculate that behavioral responses are responsible for the association of ciliates with their prey in the water column. These coincident ciliate and algal vertical distributions then respond similarly to turbulent mixing and water displacement caused by wind stress, bathymetry, or frontal convergences, resulting in horizontal patches in which the ciliates and algae remain associated. This fine scale spatial coupling between ciliates and their prey should result in higher ciliate growth rates and greater impact of ciliate grazing on phytoplankton populations than would be predicted from average ciliate or algal densities. Coincident patches of algae and ciliates may also provide higher food concentrations for larger grazers which can use both resources.

Supported by: NSF Grant OCE82-00030
and NOAA, Grant NA 81RAD00014.

WHOI Contribution No. 5532.

DEPARTMENT OF CHEMISTRY

Robert B. Gagosian, Department Chairman

CHEMISTRY

ORGANIC AND
BIOLOGICAL CHEMISTRY

THE CONCH OF Limacina AND Peraclis
(PTEROPODA) AND A MODEL FOR THE
EVOLUTION OF PLANKTONIC GASTROPODS

K. Bandel, A. Almogi-Labin,
C. Hemleben and W. G. Deuser

Pteropod shell morphologies can be explained by neotenic extension of larval features observed in benthic forms, such as the sinistral coiling of Architectonicids and the ornamentation patterns of Tonnacea into the adult shell of permanently planktonic living snails. Presented models suggest derivation of pteropods from primitive prosobranch mesogastropods of the Late Paleozoic, rather than from opisthobranchs.

Supported by: NSF Grants OCE76-21280,
OCE78-19813 and OCE81-17002.

WHOI Contribution No. 5562

MEMBRANE LIPID FROM A DEEP-SEA HYDROTHERMAL
VENT METHANOGEN: IDENTIFICATION OF A
NEW MACROCYCLIC GLYCEROL DIETHER

Paul B. Comita and Robert B. Gagosian

The membrane lipid of a new deep-sea hydrothermal vent methanogen, Methanococcus jannaschii, has been structurally characterized. The hydrolyzed polar lipid of this archaeobacterium is primarily (95%) a macrocyclic glycerol diether, which has not been described previously. The structure was elucidated by a combination of chemical and spectroscopic techniques. An initial survey of selected methanogens failed to indicate the presence of this membrane lipid in any microorganism other than Methanococcus jannaschii.

Published in: Science, 222, 1329-1331.

Supported by: NSF Grants OCE80-24256
and OCE80-24895, and a NIH/WHOI
Postdoctoral Fellowship.

WHOI Contribution No. 5450.

STRUCTURAL ELUCIDATION OF A UNIQUE
MACROCYCLIC MEMBRANE LIPID FROM A NEW,
EXTREMELY THERMOPHILIC, DEEP-SEA
HYDROTHERMAL VENT ARCHAEABACTERIUM,
Methanococcus jannaschii.

P. B. Comita, R. B. Gagosian,
H. Pang and C. E. Costello

The membrane lipid of a new deep-sea

hydrothermal vent methanogen, Methanococcus jannaschii, was isolated, purified, and structurally characterized. The total lipid extract, amounting to 32.2 µg/mg dry cell weight, was fractionated on silica gel into a neutral lipid and polar lipid fraction. The neutral lipid fraction consisted of a series of isoprenoid hydrocarbons and free (non-phospholipid) alkylglycerol ethers. The polar phospholipid and glycolipid fraction (8.44 µg/mg dry cell weight) was hydrolyzed with methanolic HCl, and the resulting alkylglycerol ethers were analyzed by a combination of chemical and spectroscopic techniques. The hydrolyzed polar lipid was primarily (95%) a unique, macrocyclic glycerol diether heretofore unknown. High field (250 MHz) proton nuclear magnetic resonance and infrared spectra of this novel macrocyclic compound are nearly identical and overlapping those of the known bis-(phytanyl)glycerol diether and bis-(diphytanyl)diglycerol tetraether. Degradation of the macrocyclic ether with boron tribromide resulted in diphytanyl dibromide, and further reaction of this dibromide with lithium aluminum hydride resulted in diphytane as determined by gas chromatography-mass spectroscopy. The significance of the predominance of this structure in Methanococcus jannaschii is discussed. A survey of selected methanogenic Archaeobacteria, including three thermophiles, failed to indicate the presence of the macrocycle in any other microorganism, including two species of order Methanococcales, one species of Methanobacteriales, and three strains belonging to the order Methanomicrobiales.

In press: Journal of Biological
Chemistry.

Supported by: NSF Grants OCE80-24256
and OCE80-24895.

WHOI Contribution No. 5449.

SUSPENDED PARTICULATE ORGANIC MATERIAL
FROM HYDROTHERMAL VENT WATERS AT 21°N

Paul B. Comita, Robert B. Gagosian
and Peter M. Williams

Suspended particulate matter from warm hydrothermal vent waters (5-38°C) of the East Pacific Rise (21°N, 106°W) was sampled on the OASIS expedition with an in situ pumping system in April 1982. Particulate organic carbon (POC) and nitrogen (PON) as well as individual organic compound classes from the lipid fraction were analyzed in an effort to ascertain the sources and composition of the organic matter in this highly productive area. Our initial results show that the particulate organic material of the vent water was both quantitatively and qualitatively different from the non-vent

bottom water. The POC and PON measurements within the warm and non-vent waters vary by a factor of two to five, the higher values being in warm water. The organic matter in the non-vent water was found to be more resistant than in the vent water, as shown by two indices, C:N ratio and total lipid: POC ratio. Contributions to the lipid class compounds were assessed using well-known organic compound source biomarkers such as hydrocarbons, fatty alcohols, sterols, tri-acylglycerols and steryl and wax esters.

In press: Nature.

Supported by: NSF Grants OCE80-24256 and OCE80-24895, and a NIH/WHOI Postdoctoral Fellowship.

WHOI Contribution No. 5446.

FLY-ASH PARTICLES INTERCEPTED IN THE DEEP SARGASSO SEA

W. G. Deuser, K. Emeis, V. Ittekkot
and E. T. Degens

Results of a continuing series of sequential sediment-trap deployments in the Sargasso Sea, covering 5 yrs, suggest that an efficient mechanism exists for the removal of both biogenic and abiogenic particles from the surface layers of the ocean and for their rapid transfer to great depth. We report here the occurrence of fly-ash particles in material collected in that experiment. Available evidence indicates that these particles were carried by winds from their sources, probably located in North America, to the Sargasso Sea and subsequently transported to deep water as components of faecal pellets or other organic aggregates. The speed of their removal from the water column suggests that their geographical distribution on the sea floor is dictated mainly by their airborne transport. The conspicuous morphology and specific chemistry and mineralogy of fly-ash particles may make them useful indicators of the presence of industrial pollutants in remote parts of the ocean. As their production is mainly limited to the past 100 yr, their appearance in sediments may serve as a time marker or as a tracer for biological down-mixing of particles.

Published in: Nature, 305, 216-218.

Supported by: NSF Grants OCE76-21280, OCE78-19813 and OCE80-24130.

WHOI Contribution No. 5388.

PROCESSES CONTROLLING THE DISTRIBUTION OF BIOGENIC COMPOUNDS IN RECENT SEDIMENTS

Robert B. Gagosian

Several physical, biochemical and geochemical processes are responsible for the distribution of biogenic compounds found in Recent sediments. A discussion is presented concerning these processes with emphasis on 1) biomarker sources from both the water column above and the benthos in surface sediments and 2) the diagenetic pathways that source organism organic compounds undergo as this material is being vertically transported through the water column and at the sediment/water interface. The steroid class compounds will be used as examples of these processes since they encompass several compound types such as saturated hydrocarbons and olefins, alcohols, ketones and ethers. Results from coastal and open ocean environments as well as the highly productive upwelling zone off Peru will be presented.

In press: American Association of
Petroleum Geologists/SEPM. W.
Meinschein, ed.

Supported by: NSF OCE82-14695.

REVIEW OF MARINE ORGANIC GEOCHEMISTRY

Robert B. Gagosian

Over the past four years the field of marine organic chemistry has expanded in many directions. In order to cover the major areas of this expansion in a review of the type presented here, several major topic areas have been selected focusing on the processes responsible for the transport and transformation of organic material. Some of these topics have been discussed previously in a review of future research problems in marine organic chemistry. They include the organic geochemistry of 1) the atmosphere over the oceans and the air/sea interface, 2) dissolved material in seawater, 3) the composition and vertical flux of particulate material as it is transported through the water column, 4) recent marine sediments, 5) manuscripts focusing on transformation reactions, 6) metal-organic interactions including adsorption/desorption processes in the ocean, and 7) selected subjects in new areas of research.

Published in: Review of Geophysics and
Space Physics 21, 1245-1258.

Supported by: NSF Grants OCE80-18436,
OCE82-14695 and OCE81-11947.

WHOI Contribution No. 5315.

SEASONALITY IN THE FLUXES OF SUGARS,
AMINO ACIDS AND AMINO SUGARS
TO THE DEEP SARGASSO SEA

V. Ittekkot, W. G. Deuser and E. T. Degens

The fluxes of sugars, amino acids and amino sugars were determined in the $<37\ \mu\text{m}$ fraction of samples collected during successive two-month sediment trap deployments in the deep Sargasso Sea ($3200 \pm 100\ \text{m}$) from April 1978 until December 1981. All fluxes varied seasonally and in phase with the flux of $<37\ \mu\text{m}$ fraction, which was shown previously to vary in phase with primary productivity in the surface layers. During the period of investigation, the fluxes of sugars and amino acids and amino sugars were in the range of 0.03 to $0.64\ \text{mg m}^{-2}\ \text{d}^{-1}$ and 0.12 to $1.7\ \text{mg m}^{-2}\ \text{d}^{-1}$, respectively. They contributed 18 to 20% of the measured organic carbon. About 40 to 50% of the measured total nitrogen could be accounted for by amino acids and amino sugars. The distribution patterns of sugars and amino acids were, in general, similar to those reported for biomineralized tissues of carbonate and silica producers and the cell walls of biomineralizing organisms. However, amounts of non-protein amino acids, i.e., β -alanine and γ -aminobutyric acid, relative to aspartic and glutamic acids, and those of amino sugars relative to total amino acids varied seasonally. They appear to indicate the nature and source of organic matter arriving at the sediment trap.

In press: Deep-Sea Research.

Supported by: NSF Grants OCE76-21280,
OCE78-19813 and OCE80-24130.

PARTICULATE AMINO ACIDS IN THE SEA:
EFFECTS OF PRIMARY PRODUCTIVITY AND
BIOLOGICAL DECOMPOSITION

Cindy Lee and Carolyn Cronin

The flux of amino acids associated with sinking particles collected by sediment traps shows a strong relationship with primary productivity at eight sites sampled. This relationship exists in spite of differences in the oceanic regimes sampled and in the sediment traps, bactericides, and amino acid analysis techniques used. A greater proportion of particulate amino acids leaves the euphotic zone in areas of higher productivity, and a larger amount of particulate amino acids reaches deeper waters in these areas. However, the particulate amino acids leaving the euphotic zone decompose faster with depth in more productive areas. That decomposition below the surface waters can occur faster in areas of high productivity

suggests that (1) remineralization may be mediated more by zooplankton and less by microbial processes than in areas of low productivity, or (2) phytoplankton growing in more productive areas are more easily remineralized than those growing in less productive areas.

Supported by: Office of Naval Research
Contract N00014-82-C-0019; NSF
Grants OCE79-25080, OCE80-24282 and
OCE83-00018.

WHOI Contribution No. 5566.

DISSOLVED, EXCHANGEABLE, AND BOUND
ALIPHATIC AMINES IN MARINE SEDIMENTS:
INITIAL RESULTS

Cindy Lee and Brenda L. Olson

Initial studies of aliphatic amines were undertaken to investigate the potential importance of these compounds in marine sediments. Marine sediments from a shallow bay on the coast of Massachusetts and from the continental margin of Mexico (Eastern Tropical North Pacific Ocean) were analyzed for short-chain aliphatic amines by gas chromatography with chemiluminescent detection. Two amines, dimethyl and trimethyl amine, were found in relatively large amounts, as much as 1% of the total organic nitrogen. Both amines were present in the interstitial waters, on exchangeable sites, and bound within the sediments. Similarities between amine and ammonia behavior suggest that, like ammonia, the amines are also a product of the decomposition of organic nitrogen compounds. However, the amines are enriched in anoxic sediments compared to oxygenated sediments over values expected from relative carbon and nitrogen content. Amines like dimethyl and trimethyl amine may be important in reducing the C/N ratio with depth in the sediment.

Supported by: Office of Naval Research
Contract N00014-82-C-0019; NSF
Grants OCE80-24282 and OCE83-00018.

WHOI Contribution No. 5567.

TRANSFORMATION REACTIONS AND RECYCLING
OF CAROTENOIDS AND CHLORINS IN THE
PERU UPWELLING REGION (15°S , 75°W)

Daniel J. Repeta and Robert B. Gagosian

Total chlorins and carotenoids were measured in suspended particulate matter, sediment trap, and zooplankton fecal pellet samples collected in the Peru upwelling region. Individual carotenoids were analyzed by high pressure liquid chromatography and

mass spectrometry. Fucoxanthin, diadinoxanthin, diatoxanthin, carotene, and peridinin constitute >95% of the total carotenoid pigments in suspended particulate matter samples. However, sediment trap and zooplankton fecal pellet samples contained significant amounts of the carotenoid transformation products fucoxanthinol, fucoxanthin 5'-dehydrate, fucoxanthinol 5'-dehydrate, peridininol and peridininol 5'-dehydrate. These samples were further characterized by high values of the ratio fucoxanthinol/total fucopigments (f/F_t) and low values of the ratio total carotenoids/total chlorins. Three transformation reactions are proposed to account for the formation of observed transformation products: heterotrophic ester hydrolysis, chemically mediated epoxide rearrangement, and microbially mediated dehydration. The distribution of specific transformation products within the samples suggests that heterotrophic ester hydrolysis is a general transformation pathway operative in the water column on a wide suite of organic esters. Epoxide rearrangement and dehydration appear to occur over a longer time scale and are more typical of sedimentary environments.

In press: *Geochimica et Cosmochimica Acta*.

Supported by: NSF Grants OCE79-25352, OCE81-18436 and OCE83-14695; ONR Contract N00014-74-CO-262 NR 083-004; and the Coastal Research Center, WHOI.

WHOI Contribution No. 5548.

PARTICLE SIZE DISTRIBUTIONS OF N-ALKANES
AND ^{210}Pb IN AEROSOLS OFF THE
COAST OF PERU

J. K. Schneider, R. B. Gagosian,
J. K. Cochran and T. W. Trull

Lipid class compounds have been used to determine the sources of organic material in atmospheric aerosols from remote marine areas. Recently, it has been shown that the temporal variability of terrestrial source marker organic compounds in aerosols from the tropical North Pacific is directly related to the seasonality of dust storm activity in China. However, very little is known about the detailed emission and transport mechanisms of these compound classes from the continents. We report here, as part of the Sea/Air Exchange Program (SEAREX), the particle size distributions of aliphatic hydrocarbons and ^{210}Pb in aerosols in the upwelling zone off the coast of Peru to further elucidate these mechanisms.

Published in: *Nature*, 304, 429-432.

Supported by: NSF Grants OCE77-12914, OCE81-11947 and OCE81-13153; and Swiss National Science Foundation 82-903-081.

WHOI Contribution No. 5349.

FREE AND ESTERIFIED STEROLS OF THE
MARINE DINOFLAGELLATE Gonyaulax polygramma

J. K. Volkman, R. B. Gagosian
and S. G. Wakeham

Free and esterified sterols of the common marine dinoflagellate Gonyaulax polygramma were identified using capillary gas chromatography-mass spectrometry. Fractions containing free 4α -methyl and 4-desmethyl sterols were isolated by column chromatography and shown to consist of at least 20 components. Major sterols included $4\alpha,23,24$ -trimethyl- 5α -cholestan- 3β -ol (dinostanol), $4\alpha,23,24$ -trimethyl- 5α -cholest- 22E -en- 3β -ol (dinosterol), cholest- 5 -en- 3β -ol (cholesterol), $23,24$ -dimethyl- 5α -cholest- 22E -en- 3β -ol and $23,24$ -dimethylcholesta- $5,22\text{E}$ -dien- 3β -ol. Although the same suite of sterols was found in the free and esterified sterol fractions, the proportions of individual sterols were quite different. The complexity of the sterol distributions, together with the predominance of dinostanol, distinguishes the sterol composition of this alga from those of other members of the Gonyaulacaceae.

In press: *Lipids*.

Supported by: NSF Grants OCE80-18436 and OCE82-14695; and ONR Contract N00014-82-C-0019.

WHOI Contribution No. 5528.

BIOGEOCHEMISTRY OF PARTICULATE ORGANIC
MATTER IN THE OCEANS: RESULTS FROM
SEDIMENT TRAP EXPERIMENTS

S. G. Wakeham, C. Lee, J. W. Farrington
and R. B. Gagosian

Particulate organic matter collected in sediment traps from various oceanic regimes - Sargasso Sea, equatorial north Atlantic, north central Pacific, California Current, and Peru coastal upwelling - have been analyzed for their lipid and amino acid composition and flux. Despite rapid settling of the large particles through the water column and a relatively small depth gradient for total organic carbon flux, there are major changes in the composition and flux of lipids and amino acids associated with the particles. The rapid disappearance of

the more labile compounds, such as amino acids and polyunsaturated fatty acids, with increasing depth, indicates that the major sources of these compounds are in the upper part of the water column and that they are readily degraded as the particles sink. On the other hand, the intermittent appearance of large amounts of wax ester, along with the changing fatty acid composition of the particles, points to deep-water sources for some of these compounds.

In press: Deep-Sea Research.

Supported by: ONR Contract
N00014-79-C-0071; NSF Grants
OCE77-26084 and OCE80-18436.

WHOI Contribution No. 5367.

STERENES IN SUSPENDED PARTICULATE MATTER IN THE EASTERN TROPICAL NORTH PACIFIC

S. G. Wakeham, R. B. Gagosian,
J. W. Farrington and E. A. Canuel

Particulate matter in the oceans is operationally partitioned into two pools. Large ($<32\mu\text{m}$) particles, such as zooplankton fecal pellets and marine snow, sink rapidly and dominate the vertical flux of organic matter in the oceans. These relatively rare particles, usually collected in sediment traps, sink at rates of tens to hundreds of meters/day leading to a significant flux of labile particulate organic matter from sites of production in the euphotic and mesopelagic zones to the deep ocean. Finer ($<2\mu\text{m}$) suspended material, on the other hand, dominates the standing stock of particulate matter. Sinking rates of only a few meters/day for small particles result in long residence times in the water column, and major transformations of the suspended organic matter may occur in situ. Source and transformation processes involving particulate organic matter in seawater and recent sediments may be traced by studying various classes of specific organic marker compounds, including the steroids. We report here the apparent production of sterenes (steroidal alkenes) on suspended particulate matter within the oxygen minimum zone in the equatorial Pacific Ocean off central Mexico. This observation provides further evidence of the importance of water column processes on early stages of the transformation of biogenic sterols to the sterenes present in mature sediments.

In press: Nature.

Supported by: ONR Contract
N00014-79C-0071.

WHOI Contribution No. 5535.

SEA WATER AND GEOCHEMISTRY

DSDP HOLE 504B, THE FIRST REFERENCE SECTION OVER 1 KM THROUGH LAYER 2 OF THE OCEANIC CRUST

R. N. Anderson, J. Honnorez, K. Becker,
A. C. Adamson, J. C. Alt, R. Emmermann,
P. D. Kempton, H. Kinoshita, C. Laverne,
M. J. Mottl and R. L. Newmark

Hole 504B provides a reference section for correlation of in situ petrological and geophysical studies with regional crustal models based on dredged samples and remote surveys; and, at least for the medium-spreading-rate Costa Rica Rift, Hole 504B confirms that the main features of the upper part of an idealized ophiolite sequence occur in the oceanic crust.

Published in: Nature, 300, 589-594.

Supported by: NSF Grant OCE80-19056.

WHOI Contribution No. 5323

IN SITU ELECTRICAL RESISTIVITY AND BULK POROSITY OF THE OCEANIC CRUST COSTA RICA RIFT

K. Becker, R. P. Von Herzen,
T. J. G. Francis, R. N. Anderson,
J. Honnorez, A. C. Adamson, J. C. Alt,
R. Emmermann, P. D. Kempton, H. Kinoshita,
C. Laverne, M. J. Mottl and R. L. Newmark

In situ electrical resistivity was measured in DSDP Hole 504B to a depth of 1,013 m into oceanic basement. Apparent resistivities are about $10\Omega\text{m}$ in about 600 m of pillow lavas, sharply increasing to nearly 1,000 Ωm in the underlying dykes. Bulk porosities calculated from resistivities show a distinct layering, apparently corresponding to geophysical layers 2A, 2B and 2C.

Published in: Nature, 300, 594-598.

Supported by: NSF Grant OCE81-17693.

WHOI Contribution No. 5322.

RESIDUAL GEOID ANOMALIES IN ATLANTIC OCEAN BASIN: RELATIONSHIP TO MANTLE PLUMES

Carl Bowin and Geoffrey Thompson

The relations of geoid data from radar

altimeter measurements (GEOS-3) to water depth and geochemical variations along the Mid-Atlantic Ridge are investigated. Using an optimum estimation method, eight geophysical lines across the ridge in the Atlantic Ocean were constructed approximately normal to the local ridge trend. Data were interpolated at equally spaced intervals along each line for application of time-series analysis techniques. Profiles of geoid, water depth, and crustal age were assembled for each line. Slopes of geoid per kilometer of topographic relief show an asymmetry of geoid/topography slopes between the two sides of the ridge that generally is not matched by topography, which has better symmetry between the two flanks. The geoid asymmetry is due to a regional field unrelated to the structure of the spreading ridge system that obscures the local anomaly relations. Subtracting a regional field, based on previous work, produces residual geoid anomalies whose source must be in the earth's outer 600 km. The residual geoid anomalies of the North Atlantic show striking correlation with the axis of the Mid-Atlantic Ridge, but those in the South Atlantic do not. Instead, major positive residual geoid anomalies appear over the Rio Grande and Walvis Ridges. The residual geoid anomaly provides further support for the removal and interpretation of the regional geoid field.

Determination of slopes, S , of residual geoid per million years crustal age reveals that S increases with greater residual geoid anomaly and with higher (shallower) crestal ridge topography. The slope of 0.27 m/m/y over the Azores High suggests that the crestal residual geoid and depth anomalies are not of thermal origin. Correlation of residual geoid anomalies with depth and geochemical (especially FeO^*/MgO , La/Sm , Nb/Zr ratios, and TiO_2) variations along the length of the Mid-Atlantic Ridge suggest that the residual geoid and crustal depth anomalies are largely a function of crustal thickness and density variations induced by plume volcanism.

In press: Journal of Geophysical Research.

Supported by: NSF Grant OCE82-08407

CHEMICAL OCEANOGRAPHY OF THE PERSIAN GULF

Peter G. Brewer

Measurements of basic hydrographic properties, nutrient elements, alkalinity and total carbon dioxide were made in the Persian Gulf in February and March 1977, on R/V ATLANTIS II Cruise 93. The results show clearly the penetration into the Gulf of surface Indian Ocean water, initially of

36.5‰ salinity which is carried northwards along the Iranian Coast, gradually increasing in salinity to more than 40‰. The inflowing surface water is high in phosphate (1.0–0.5 $\mu\text{moles/kg}$) due to upwelling induced by the northeast monsoon. However nitrate levels are at, or below, the detection limit and nitrogenous nutrients appear to be limiting.

The seawater carbonate system is greatly perturbed by the precipitation of large quantities of calcium carbonate. The alkalinity/salinity relationship shows that the greatest carbonate losses occur in the shallow waters of the Trucial Coast, where up to 125 $\mu\text{moles/kg}$ of CaCO_3 have been lost from the water. Dense, saline low specific alkalinity water formed in the shallows appears to migrate down slope to augment the outflowing waters in the deeper part of the Gulf. The outflow of the Tigris, Euphrates and Karun rivers to the north provides an intense, but low capacity signal, of low salinity, high specific alkalinity waters. Large amounts of CO_2 are also fixed by photosynthetic processes.

The classical Mediterranean type of circulation is plainly revealed. The residence time of water within the Persian Gulf is approximately 2.5 years.

In press: Progress in Oceanography, Swallow Festschrift.

Supported by: National Science Foundation and Dept. of the Environment, Iran.

WHOI Contribution No. 5529.

A CLIMATIC FRESHENING OF THE DEEP NORTH ATLANTIC (NORTH OF 50°N) OVER THE PAST 20 YEARS

P. G. Brewer, W. S. Broecker,
W. J. Jenkins, P. B. Rhines,
C. G. Rooth, J. H. Swift,
T. Takahashi and R. T. Williams

Observations made in the summer of 1981 show a significant and widespread decrease in the salinity of deep waters, averaging 0.02‰, in the sub-polar North Atlantic over the past two decades. This implies a relatively rapid response of deep-water formation to climatic perturbation.

Published in: Science, 222, 1237–1239.

Supported by: DoE, NSF Grants
OCE80-08160, OCE80-21378,
OCE79-25888 and OCE81-10646.

WHOI Contribution No. 5516.

THE MIXING PROFILE OF ORGANIC CARBON AND
SEVERAL RADIOISOTOPES IN
ABYSSAL RED CLAY SEDIMENTS

E. R. M. Druffel, P. M. Williams,
H. D. Livingston and M. Koide

Red clay sediments from the North Pacific Ocean were collected using standard box core techniques. Ionium/thorium analyses revealed that the clay sedimentation rate in three cores collected within a 50 km radius ranged from 0.75 to 1.05 mm/10³ yrs. The mixed layer appeared to be in the top 10 cm in all cores. However, analyses of the organic carbon in thin layers of sediment revealed that radiocarbon was present much deeper in the cores than was expected from the I₀/Th distribution. In addition, both the stratigraphy and inventory of radiocarbon was significantly different between box cores. The distributions and inventories of cesium-137 and plutonium-239+240 were similar to that found for radiocarbon, further illustrating spatial variability of radionuclides in the oligotrophic North Pacific red clays. These data imply that the major process controlling the distribution of organic carbon in red clay sediments is bioturbation. Diffusion may also be important for transporting organic carbon to depths below the mixed layer. A likely scenario involves variability in the activity of burrowing organisms within the sediments, which could be partially controlled by the infrequent "large falls" of dead organisms to the sediment/water interface. Infrequent high energy bottom currents may be responsible for redistribution of particulate matter at the sediment/water interface, which would account for the varying inventories of the bomb-produced radionuclides.

In press: Earth and Planetary Science
Letters

Supported by: NSF Grant OCE79-17652;
DoE Contracts DE-AT03-82ER and
DE-AC-1-81EV10694; and Sandia
Laboratories Contract 37-3164.

WHOI Contribution No. 5517.

ALKALI AND Sr ISOTOPE GEOCHEMISTRY OF
WATERS COLLECTED FROM BASALTIC BASEMENT,
DEEP SEA DRILLING PROJECT HOLE 504B,
COSTA RICA RIFT

S. R. Hart and M. J. Mottl

Basalt formation waters collected from Hole 504B at sub-basement depths of 194, 201, 365, and 440 meters show inverse linear relationships between ⁸⁷Sr/⁸⁶Sr and Ca, ⁸⁷Sr/⁸⁶Sr and Sr, and K and Ca. If the Ca

content of a fully reacted formation water end-member is assumed to be 1340 ppm, the K, Sr, and ⁸⁷Sr/⁸⁶Sr values for the end-member are 334 ppm, 7.67 ppm, and 0.70836, respectively. With respect to contemporary seawater at Hole 504B, K is depleted by 13%, Sr is enriched by 2.7%, and ⁸⁷Sr/⁸⁶Sr is depleted by 0.8%. The Sr/Ca ratio of the formation water (0.0057) is much lower than that of seawater (0.018) but is similar to the submarine hot spring waters from the Galapagos Rift and East Pacific Rise and to geothermal brines from Iceland. At the intermediate temperatures represented by the Hole 504B formation waters (70°-105°C), the interaction between seawater and the ocean crust produces large solution enrichments in Ca, the addition of a significant basalt Sr isotope component accompanied by only a minor element at Sr component, and the removal from solution of seawater K. The Rb, Cs and Ba contents of the formation waters appear to be affected by contamination, possibly from drilling muds.

Published in: Initial Reports of the
Deep Sea Drilling Project, 69,
487-494.

Supported by: NSF Grants OCE79-09457
and OCE80-19056.

WHOI Contribution No. 5425

GEOCHEMISTRY OF RARE-EARTH ELEMENTS IN
BASALTS FROM THE WALVIS RIDGE:
IMPLICATIONS FOR ITS ORIGIN AND EVOLUTION

Susan E. Humphris and Geoffrey Thompson

Selected basalts from a suite of dredged and drilled samples (IPOD sites 525, 527, 528 and 530) from the Walvis Ridge have been analyzed to determine their rare earth element (REE) contents in order to investigate the origin and evolution of this major structural feature in the South Atlantic ocean. All of the samples show a high degree of light rare earth element (LREE) enrichment, quite unlike the flat or depleted patterns normally observed for normal mid-ocean ridge basalts (MORBs). Basalts from Sites 527, 528 and 530 show REE patterns characterized by an arcuate shape and relatively low (Ce/Yb)_N ratios (1.46-5.22), and the ratios show a positive linear relationship to Nb content. A different trend is exhibited by the dredged basalts and the basalts from Site 525, and their REE patterns have a fairly constant slope, and higher (Ce/Yb)_N ratios (4.31-8.50).

These differences are further reflected in the ratios of incompatible trace elements, which also indicate considerable variations within the groups. Mixing hyperbolae for these ratios suggest that simple

magma mixing between a "hot spot" type of magma, similar to present day volcanics of Tristan da Cunha, and a depleted source, possibly similar to magmas being erupted at the Mid-Atlantic Ridge, was an important process in the origin of parts of the Walvis Ridge, as exemplified by Sites 527, 528 and 530. The Site 525 and dredged basalts cannot be explained by this mixing process, and their incompatible element ratios suggest either a mantle source of a different composition or some complexity to the mixing process. In addition, the occurrence of different types of basalt at the same location suggests there is vertical zonation within the volcanic pile, with the later erupted basalts becoming more alkaline and more enriched in incompatible elements.

The model proposed for the origin and evolution of the Walvis Ridge involved an initial stage of eruption in which the magma was essentially a mixture of enriched and depleted end member sources, with the N-MORB component being small. The dredged basalts and Site 525, which represent either later stage eruptives or those close to the hot spot plume, probably results from mixing of the enriched mantle source with variable amounts and variable low degrees of partial melt of the depleted mantle source. As the volcano leaves the hot spot, these late stage eruptives continue for some time. The change from tholeiitic to alkalic volcanisms is probably related either to evolution in the plumbing system and magma chamber of the individual volcano or to changes in the depth of origin of the enriched mantle source melt similar to processes in Hawaiian volcanoes.

In press: Earth and Planetary Science Letters.

Supported by: NSF Grants OCE74-2297, OCE78-26842 and OCE80-24629.

WHOI Contribution No. 5420.

TRACING OCEAN CIRCULATION BY MEANS OF CHEMICAL AND RADIOCHEMICAL DATA

William J. Jenkins

The oceans will play an important role in both the control of anthropogenic CO₂ levels in the atmosphere and the maintenance or modification of climatic response to the perturbation. To realistically predict and understand the nature and magnitude of the oceanic response to and impact on the future climate-CO₂ system, we must understand quantitatively the mode and mechanisms of transport (of both heat and CO₂) by the oceans and, ultimately, how those mechanisms respond to climatic perturbations.

Although we are some distance from this understanding, observation of anthropogenic (transient) tracers is giving us new insights into this area. Past and present ¹⁴C distributions tell of CO₂ exchange rates and surface water residence times. Tritium and its daughter ³He show us that the nature of thermocline "ventilation" is diffusive or exchange-like rather than advective. In this regard, the Transient Tracers Program and GEOSECS are giving us critical information that allow us to design, test and calibrate ocean transport models.

Finally, the time record of oceanic response afforded us by hydrographic series (e.g. Panulirus) and ¹⁴C-coral records (e.g. Belize) are showing us the actual oceanic response to climatic perturbation. The two kinds of records appear to be consistent with one another, and the former appears (when calibrated with transient tracers) to be consistent with the mode of thermocline ventilation deduced from TTO-GEOSECS type data.

In press: The Proceedings of the Sixth ORNL Life Sciences Symposium - The Global Carbon Cycle: Analysis of the Natural Cycle and Implication of Anthropogenic Alterations for the Next Century.

Supported by: NSF Grant OCE79-21378.

THE SEDIMENTATION RATES OF TRACE ELEMENTS IN THE SARGASSO SEA MEASURED BY SEDIMENT TRAP

T. D. Jickells, W. G. Deuser and A. H. Knap

The sedimentation rates of aluminium, cadmium, copper, iron, manganese, nickel, phosphorus, lead, vanadium and zinc in the Sargasso Sea have been determined from samples collected in a sediment trap 3200 m below the ocean surface. The fluxes are temporally variable being driven by the seasonal cycle of primary production in the surface waters. There is evidence to suggest anthropogenic perturbation of the copper, lead and zinc fluxes. The measured fluxes imply short (<100 day) lifetimes of particulate trace elements in the surface ocean.

In press: Deep-Sea Research

Supported by: NSF Grants OCE78-19813 and OCE80-24130.

WHOI Contribution No. 5553.

TROPOSPHERIC NO_x AND O₃ BUDGETS
IN THE EQUATORIAL PACIFIC

S. C. Liu, M. McFarland, D. Kley,
O. Zafiriou and B. Huebert

By combining the results of model calculations and simultaneous measurements of O₃, NO, HNO₃, particulate NO₃⁻, and meteorological parameters in the mid-Pacific, we have investigated the budget and photochemistry of near surface atmospheric ozone and odd nitrogen. The observed decrease of O₃ toward the Intertropical Convergence Zone is consistent with the transport of O₃ from higher latitude and a photochemical sink for O₃ by the reaction between H₂O and O(¹D). The total deposition flux of odd nitrogen deduced from the data is about $1.3 \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$, or 5 MT(N)/yr averaged globally. Most of the deposition flux is due to HNO₃ and particulate NO₃⁻. This deposition flux is probably representative of values over most of the Pacific because the HNO₃ and particulate NO₃⁻ densities measured over the Pacific during the GAMETAG experiments are consistent with our values. Since the residence time of odd nitrogen in the lower troposphere is short, a widely distributed, diffuse source of odd nitrogen is required to balance the deposition flux. It seems that the NO formed by cloud to cloud lightning in the tropical upper troposphere is the most likely source.

Published in: Journal of Geophysical
Research, 88, 1360-1368.

Supported by: NSF Grant ATM-8108399.

WHOI Contribution No. 5081.

PROGRAMMED PYROLYSIS OF ORGANIC MATTER FROM
THERMALLY ALTERED CRETACEOUS BLACK SHALES

K. E. Peters, J. K. Whelan,
J. M. Hunt and M. E. Tarafa

Organic-rich Cretaceous black shales from the Cape Verde Rise in the eastern Atlantic were penetrated by hot diabase sills during the Miocene. Programmed pyrolysis and pyrolysis-gas chromatography of organic matter from core samples taken at various distances from a major sill were used to show: (1) the type of kerogen, (2) its relative level of thermal maturity, (3) the fraction of pyrolyzable organic compounds which are free, and (4) the compositions of the bitumen and kerogen.

The dominant factor affecting these sediments appears to be thermal maturity rather than differences in the quality of the organic matter. Systematic changes in the pyrolyzate resulted from thermal cracking of volatile organics from the oil-

prone Type II kerogen. Generation of these products cause progressive aromatization of the residual kerogen closer to the sill. These conclusions are supported by changes in kerogen elemental compositions, quantity of bitumen extract, and vitrinite reflectance.

Although the major sill is 15 m (~50 ft) thick, solvent extraction and pyrolysis results show that hydrocarbon generation was restricted to within about 10 m (~33 ft) of the shale/sill contacts. At equal distances, the maximum temperature reached by the shales was higher above than below the sill. The reflectance of vitrinite responds more rapidly than bitumen composition to high temperatures imposed over a short time.

Published in: The American Association
of Petroleum Geologists Bulletin,
67, 11, 2137-2146.

Supported by: NSF Grant OCE80-19508 and
DoE Contract No. EG-77-S-02-4392.

HYDROTHERMAL ACTIVITY IN THE
TAG HYDROTHERMAL FIELD, MID-ATLANTIC
RIDGE CREST AT 26°N

P. A. Rona, G. Thompson, M. J. Mottl,
J. A. Karson, W. J. Jenkins, D. Graham,
M. Mallette, K. Von Damm and J. M. Edmond

The first submersible observations of the only known active submarine hydrothermal field on a slow-spreading oceanic ridge are incorporated with results of ten prior years of investigation to derive an understanding of periodicity, duration, and intensity of cyclic, episodic, multi-stage hydrothermal activity at the TAG Hydrothermal Field on the mid-Atlantic Ridge crest near latitude 26°N. Hydrothermal activity has persisted at this location for at least 1×10^6 y based on the distribution of hydrothermal and hydrogenous mineralization with respect to crustal age. High-temperature black smoker-type hydrothermal venting with a periodicity of the order of 1×10^4 y and duration of the order of 10^1 y is deduced from the ages of discrete sedimentary layers anomalously enriched in Cu, Fe, and Zn, and corresponds with the independently determined periodicity of volcanic eruptive cycles on the Mid-Atlantic Ridge. The most recent episode of black smoker-type venting is inferred to have ceased within the past 20 years based on metal enrichment in the surficial sediment layer and the occurrence of intact bivalve-like forms with the characteristics of vent clams on bottom photographs. A low-temperature hydrothermal stage of the order of 1×10^4 y in duration is inferred from thick deposits of layered and earthy manganese oxide, iron oxide, hydroxide and silicate.

The fresh appearance of intact tubules that acted as vents on relict deposits and temporal variations in the record of temperature anomalies and excess ^3He in the near-bottom water column indicate that the low-temperature hydrothermal discharge is intermittent at individual vents on a time scale of years.

In press: Journal of Geophysical Research.

Supported by: NSF Grant OCE82-00812.

WHOI Contribution No. 5454.

ALTERATION OF THE OCEANIC CRUST:
IMPLICATIONS FOR GEOCHEMICAL CYCLES
OF LITHIUM AND BORON

W. E. Seyfried, D. R. Janecky
and M. J. Mottl

Fresh tholeiitic basalt glass has been reacted with seawater at 150°C (water/rock mass ratio of 10), and fresh diabase has been reacted with a Na-K-Ca-Cl fluid at 375°C (water/rock mass ratios of 1, 2 and 5) to understand better the role of temperature, basalt composition, and water/rock mass ratio on the direction and magnitude of B and Li exchange during basalt alteration. At 150°C, slight but nevertheless significant amounts of B and Li were removed from seawater and incorporated into a dominantly smectite alteration phase. At 375°C, however, B and Li were leached from basalt. B behaved as a "soluble" element and attained concentrations in solution limited only by the B concentration in basalt and the water/rock mass ratio. Li, however, was less mobile. For example, at water/rock mass ratios of 1, 2 and 5, the percent of Li leached from basalt was 58, 70 and 92%, respectively. This suggests some mineralogic control on Li mobility during hydrothermal alteration of basalt, especially at low water/rock mass ratios. In general, these results, as well as those for B, are consistent with the temperature-dependent chemistry of altered seafloor basalt and the chemistry of ridge crest hydrothermal fluids.

Based on the distribution and chemistry of products of seafloor weather, low (<150°C) and high-temperature hydrothermal alteration of basalt, and the chemistry of ridge crest hydrothermal fluids, it was estimated that alteration of the oceanic crust is a Li source for seawater. This is not true for B, however, since the hot spring flux estimated for B is exactly balanced by low-temperature basalt alteration. These data, coupled with B and Li flux estimates for other processes (e.g., continental weathering, clay mineral adsorption, authi-

genic silicate formation and formation of siliceous skeletal material), yield new insight into the B and Li geochemical cycles. Calculations performed here indicate relatively good agreement between the magnitude of B and Li sources and sinks. The geochemical cycle of B, however, may be affected by serpentinization of mantle derived peridotite in oceanic fracture zones. Serpentinites are conspicuously enriched in B and if the B source for these rocks is seawater, then an additional B sink exists which must be integrated into the B geochemical cycle. However, until more data are available in terms of areal extent of serpentinization, serpentinite chemistry and isotopic composition, the importance of B in these rocks with respect to the B geochemical cycle remains speculative at best.

In press: Geochimica et Cosmochimica Acta.

Supported by: NSF Grants OCE81-10913 and OCE80-18644.

WHOI Contribution No. 5522.

BASALT - SEAWATER INTERACTION

Geoffrey Thompson

The upper part of the igneous oceanic basement consists of basaltic lava. This basalt undergoes reaction with seawater over a range of temperatures, time and locale. This reaction is a major source and sink for various ions in seawater and is a major process in buffering seawater composition and forming metalliferous ores in the marine environment.

The nature of the chemical reaction and the fluxes of ions exchanged between the oceans and the igneous basement are mostly dependent on the temperature of reaction and the relative proportion of the reactants. These vary in respect to the location of the water circulation and distance from the heat source. Four examples of seawater-basalt interactions are considered and the net exchange fluxes are calculated; these examples cover the range of temperature and water:rock ratios typically found in the ocean floor.

Low temperature, high water:rock ratio is typical of the exchange in the upper few meters of the oceanic basement. Only about 0.1% of the new oceanic crust undergoes this reaction which extends over a time period of tens of millions of years. The annual fluxes produced are not very large. Low temperature, low water:rock ratio is typical of the reaction in the deeper parts of the oceanic basement. About 8% of newly formed oceanic crust can be expected to undergo this reaction over a period of a

few million years. Fractions and fluxes on the flanks of spreading centers are at moderate temperatures and water:rock ratios. These reactions are relatively short lived, but the fluxes produced are quite high. High temperature reaction of seawater and basalt (in excess of 100°C) takes place at spreading center axes. These reactions are fast but result in very high fluxes and formation of polymetallic sulfides or iron and manganese oxides. The products of this reaction and the direction of exchange for some ionic species are quite different compared to the lower temperature reactions.

The net effect of the basalt-seawater exchange is the sum of all the reaction fluxes over the full temperature range. This calculated net flux indicates that the basalt is a source of ions such as Si, Ca, Ba, Li, Fe, Mn, Cu, Ni, Zn and hydrogen ions. It also is the sink for ions such as Mg, K, B, Rb, H₂O, Cs and U. The annual fluxes calculated for some of these species is of the same order of magnitude as the annual river influxes into the ocean.

In press: Hydrothermal Processes at Seafloor Spreading Centers. Plenum Publishing Corp., NY.

Supported by: NSF Grants OCE82-24930 and OCE82-00812.

WHOI Contribution No. 5360.

MORPHOLOGY, MINERALOGY AND CHEMISTRY OF HYDROTHERMAL DEPOSITS FROM THE TAG AREA, 26°N MID-ATLANTIC RIDGE

Geoffrey Thompson, Michael J. Mottl and Peter A. Rona

Metalliferous hydrothermal deposits have been observed and sampled by submersible from the TAG Hydrothermal Field on the Mid-Atlantic Ridge at 26°N. The deposits include surficial, metal-rich staining on the surface of carbonate ooze deposits over considerable areas of the seafloor as well as discrete, massive layered deposits. The latter include hydrothermal precipitates of Fe-silicate (nontronite), Mn-oxide (birnessite), and Fe-oxide (amorphous). The deposits range in size from less than a meter to about 15x20 m. The composition and morphology of individual deposits varies from thick, laminated crystalline birnessite precipitates, to iron-rich tubular vents, to loose, earthy birnessite, nontronite or amorphous iron oxide interlayered deposits. Fractionation and crystallinity are presumably controlled by local Eh conditions and the hydrodynamics of circulation in the upper crust. They are presumed to represent deposition from low temperature hydrothermal solutions which had originally reacted with

basaltic oceanic crust at depth at high temperature and had precipitated metal-rich sulfides within the crust. Occasional short duration, high temperature, black-smoker type venting is inferred to have occurred based on Cu-Fe-Zn rich layers in the sediments and observations of clam shells in bottom photographs.

In press: Chemical Geology.

Supported by: NSF Grant OCE82-00812.

PHOTOCHEMISTRY AND THE SEA-SURFACE MICROLAYER: NATURAL PROCESSES AND POTENTIAL AS A TECHNIQUE

Oliver C. Zafiriou

The largest effects of photochemical processes in the Top Boundary Layer probably involve reactions at the sea surface of species formed by atmospheric processes. The ozone influx is probably the largest and most significant one. Ozone reacts with dissolved iodide and with other unknown components. One consequence is the formation of volatile iodine species, perhaps in sufficient amounts to balance the atmospheric I budget. However, most of the ozone reaction products undergo other fates.

These reactions drive chemical fluxes that may be coupled to other Top Boundary Layer Processes in three ways: (1) modification of air-sea gas exchange processes and fluxes, (2) chemical modification of trace components of the Top Boundary Layer, especially of species with long residence times there (indigenous biota, surface-active molecules, particles), and (3) chemical modification of the physical properties of the interface via reaction of surface active materials, leading to changes in surface physical properties. These mechanisms may link the photochemically-driven fluxes to trace element fluxes, surface biology, the fate of particles at the interface, and the damping of capillary waves by surface tension effects.

In order to substantiate some of these fluxes and effects, better methods of studying and of modeling the air-sea interface on scales even below 1 micron are needed. One possible tool in the required arsenal of new techniques may be the use of photochemical processes to generate materials at known rates and with known spatio-temporal distributions.

In press: NATO Conference Series IV: Marine Sciences. Plenum Press.

Supported by: NSF Grants OCE81-11947 and OCE82-15614.

WHOI Contribution No. 5520.

SEASONALITY IN THE FLUX OF NATURAL
RADIONUCLIDES AND PLUTONIUM
IN THE DEEP SARGASSO SEAMichael P. Bacon, Chih-An Huh,
Alan P. Fleer and Werner G. Deuser

A record of radionuclide fluxes at a deep-ocean station near Bermuda (32°05'N, 64°15'W) was obtained from analysis of a three-year collection of sediment-trap samples. The trap was placed at a depth of 3200 m, 1000 m above the seafloor, and the samples were recovered at two-month intervals. Concentrations of ^{238}U , ^{234}U , ^{232}Th , ^{230}Th , ^{228}Th , ^{231}Pa , ^{210}Pb , ^{210}Po and $^{239+240}\text{Pu}$ were measured in the trapped material. Most of the radionuclide activity was found in the <37- μm sieved fraction. All of the radionuclide fluxes showed seasonal variations that were in phase with the variations in total sediment flux, which were shown in earlier work to be closely tied to the annual cycle of primary production in the overlying surface water. The seasonal variations are especially noteworthy for ^{230}Th and ^{231}Pa , considering that most of their production occurs in the water column below the euphotic zone. Evidently the seasonal influence is transmitted downward by the varying particle flux so that radionuclide scavenging rates at depth, as well as on the surface, are affected. It is suggested that this could be brought about by seasonal variations in the flux of marine snow or in the rate of fecal-matter production in the deep water column. Fluxes of ^{230}Th and ^{231}Pa integrated over the annual cycle yielded a trapping efficiency of 105±17% for the PARFLUX sediment trap used in this investigation.

Supported by: NSF Grants OCE82-19583,
OCE76-21280, OCE78-19813 and
OCE80-24130; and ONR Contract
N00014-82-C0019 NR 083-004.

WHOI Contribution No. 5452.

 ^{210}Po AND ^{210}Pb DISTRIBUTIONS IN THE
CENTRAL AND EASTERN INDIAN OCEANJ. K. Cochran, M. P. Bacon,
S. Krishnaswami and K. K. Turekian

Disequilibrium between ^{210}Po and ^{210}Pb and between ^{210}Pb and ^{226}Ra has been mapped in the eastern and central Indian Ocean based on stations from Legs 3 and 4 of the GEOSECS Indian Ocean expedition. $^{210}\text{Po}/^{210}\text{Pb}$ activity ratios are less than 1.0 in the surface mixed layer and indicate a residence time for Po of 0.6 years. ^{210}Po

and ^{210}Pb are generally in radioactive equilibrium elsewhere in the water column except at depths of 100–500 m, where Po may be returned to solution after removal from the surface water, and in samples taken near the bottom at a few stations. ^{210}Pb excesses relative to ^{226}Ra are observed in the surface water but these excesses are not as pronounced as in the North Pacific and North Atlantic. The difference is attributable to a lower flux of ^{210}Pb from the atmosphere to the Indian Ocean. Below the main thermocline, ^{210}Pb activities increase with depth to a broad maximum before decreasing to lower values near the bottom. Departures from this pattern are especially evident at stations taken in the Bay of Bengal (where $^{210}\text{Pb}/^{226}\text{Ra}$ activity ratios as low as 0.16 are observed) and near the Mid-Indian Ridge. The data suggest that removal of ^{210}Pb at oceanic boundaries, coupled with eddy diffusion along isopycnals, can explain gradients in ^{210}Pb near the boundary. Application of a simple model including isopycnal diffusion, chemical removal, production and radioactive decay produces fits to the observed $^{210}\text{Pb}/^{226}\text{Ra}$ gradients for eddy diffusion coefficients of $\sim 10^7 \text{ cm}^2/\text{s}$. High productivity in surface waters of the Bay of Bengal makes this region a sink for reactive nuclides in the northern Indian Ocean.

Published in: Earth and Planetary
Science Letters, 65, 433–452.

Supported by: NSF Grants OCE77-20783
and OCE77-21258.

WHOI Contribution No. 5389.

NEUTRON ACTIVATION ANALYSIS OF RARE
EARTH ELEMENTS IN SEAWATER

Hein J. W. DeBaar

Novel methods were developed for determination of picomolar concentrations of 12 Rare Earth Elements (REE) in seawater. Rigorous separation of sodium (Na) and uranium (U) by three consecutive ion exchange steps is essential to avoid interferences. Ultraclean procedures and reagents are crucial to prevent contamination with REE, Na or U. The purified REE extract is irradiated for 8 hours with a high neutron flux. The various gamma spectra are recorded and processed by computer. An internal standard (^{144}Ce) is carried through the procedure to compensate for variations in counting-geometry.

In press: Proceedings of International Symposium on the Use and Development of Low and Medium Flux Research Reactors, MIT, Cambridge, 1983. Special Volume of Atomkern-energy and Atomtechnik.

Supported by: DoE Contract
DE-AS02-76EV03566.

WHOI Contribution No. 5519.

DETERMINATION OF LEAD-210 AND POLONIUM-210 IN SEAWATER AND MARINE PARTICULATE MATTER

Alan P. Fleer and Michael P. Bacon

Natural ^{210}Pb and ^{210}Po are determined in seawater and suspended particulate matter by alpha spectrometry with silicon surface-barrier detectors. Procedures used at WHOI include a pre-concentration, based on co-precipitation with cobalt-APDC, which can be carried out at sea. Source preparation is by spontaneous deposition of Po on silver, with ^{210}Pb being determined by measurement of ^{210}Po ingrowth. Yields are monitored by the addition of ^{208}Po and stable Pb tracers. A desk-top computer is programmed to process the alpha spectra and make radioactive growth and decay computations. The accuracy of oceanic ^{210}Pb data is estimated to be 10 percent, which is significantly worse than the attainable precision.

In press: Nuclear Instruments and
Methods.

Supported by: DoE Contract
DE-AS02-76EV03566.

WHOI Contribution No. 5399.

HELIUM ISOTOPIC VARIATIONS IN VOLCANIC ROCKS FROM LOIHI SEAMOUNT AND THE ISLAND OF HAWAII

M. D. Kurz, W. J. Jenkins,
S. R. Hart and D. Clague

Helium isotopic ratios ranging from 20 to 32 times the atmospheric $^3\text{He}/^4\text{He}$ (R_A) have been observed in a study of 15 basaltic glasses from the Loihi Seamount. These ratios, which are up to four times higher than those of MORB glasses and more than twice those of nearby Kilauea, are strongly suggestive of a primitive source of volatiles supplying this volcanism. The Loihi glasses measured span a broad compositional range, and the $^3\text{He}/^4\text{He}$ ratios were found to be generally lower for the alkali basalts than for the tholeiites. The component with a lower $^3\text{He}/^4\text{He}$ ratio appears to be associated with olivine xenocrysts, within which fluid inclusions are probably the carrier of contaminant helium. One Loihi sample has a much lower isotope ratio ($<5 R_A$), but a combination of low He concentration, high vesicularity, and presence of cracks lined with clay minerals suggests that the low

ratio is due to gas loss and contamination by atmospheric helium.

Crushing and melting experiments show that for modest vesicularities ($<5\%$ by volume) the Loihi glasses obey a MORB type partitioning trend, but at higher vesicularities the data show considerably more scatter due to volatile mobilization. The high vesicularities, low extrusion pressure and generally low helium concentrations are consistent with a considerable degree of degassing. Analyses of dunites, plus a correlation between total helium concentrations with xenocryst abundances, also suggest that xenocrysts are a significant carrier of contaminating (low $^3\text{He}/^4\text{He}$) helium.

$^3\text{He}/^4\text{He}$ ratios from samples of other Hawaiian volcanoes (Kilauea, Mauna Loa, Hualalai, and Mauna Kea) show a smooth decrease in $^3\text{He}/^4\text{He}$ with increasing volcano age and volume. We interpret this to be a synoptic picture of the time evolution of a hot-spot diapir. The earliest stage is characterized by primitive ($>30R_A$) helium with some (variable) component of lithospheric contamination added during "break-through", while the later stages are characterized by a relaxation toward lithospheric $^3\text{He}/^4\text{He}$ ratios ($\sim 8 R_A$) due to isolation of the diapir from the mantle below (as the plate moves on) and subsequent mining of the inherited helium and contamination from the surrounding lithosphere. The abrupt contrast in $^3\text{He}/^4\text{He}$ ratios between Kilauea and Loihi, despite their close proximity, is indicative of the small lateral extent of the plume.

In press: Earth and Planetary Science
Letters.

Supported by: NSF Grant OCE81-00812;
WHOI Education Program; and WHOI
Ocean Industries Program.

WHOI Contribution No. 5393.

ANTHROPOGENIC TRITIUM IN SOUTH ATLANTIC BOTTOM WATER

W. J. Jenkins, D. E. Lott, M. W. Pratt
and R. D. Boudreau

By observation of the invasion of man-made substances into the oceans, we learn about the nature and rates of mixing and flow. One of the more useful of these tracers is tritium (^3H), which was produced by the atmospheric thermonuclear weapons tests of the 1950s and 1960s and which entered the oceans in a pulse-like fashion, totally dominating the natural background. The production and entry of this radioisotope (half-life ca. 12.5 y) into the ocean took place largely in the Northern Hemisphere, so that corresponding concentra-

tions of tritium in the Southern Hemisphere are five- to ten-fold smaller. Unfortunately, this means that the dynamic range (the ratio of maximum observable to minimum detectable concentrations) of tritium is limited for southern hemispheric work. If the minimum detectable quantity (that is, the detection limit) of tritium could be reduced, the evolution of this transient tracer would still provide valuable insight into mixing and advection. By increasing sample size and altering sample handling techniques, we have now extended the detection limit by more than an order of magnitude, and have detected what we believe to be man-made tritium in Antarctic bottom water at about 40°S in the western South Atlantic.

Published in: Nature, 305, 45-46.

Supported by: NSF Grant OCE78-21734

WHOI Contribution No. 5470.

IN SITU CHEMISORPTION OF TRANSURANICS FROM SEAWATER

Don R. Mann, Lolita D. Surprenant
and Susan Casso

Measurement of transuranics in open ocean seawater generally has required the collection and processing of relatively large volume samples. A system based on in situ transuranic adsorption on manganese dioxide is presented which (1) eliminates the need for collecting and transporting very large volumes of seawater; and (2) permits substantial improvements in analytical precision and sensitivity because of the increased volume of water sampled. Water to be stripped of transuranics is passed sequentially through a pre-filter and two MnO₂ impregnated fiber filter cartridges. The system has been used to collect both surface and, via an in situ pumping system, subsurface samples as deep as 5,800 m. Details of the operation of this system and examples of the improved data quality realized by this technique are presented and compared to bulk water methods. Of particular note is the demonstration of the capability of measuring ²³⁸Pu/^{239,240}Pu ratios in seawater of low Pu concentration.

In press: Nuclear Instruments and
Methods

Supported by: DoE Contracts
DE-AC02-81EV10694 and
DE-AC02-76EV03563; and Sandia
National Laboratories Contract
37-3164.

WHOI Contribution No. 5398.

PLUTONIUM OXIDATION STATE DISTRIBUTIONS IN THE PACIFIC OCEAN DURING 1980-1981

D. M. Nelson, A. E. Carey and V. T. Bowen

The concentration of plutonium and its oxidation state distribution have been measured as a function of depth at three locations in the North Pacific Ocean. Concentration profiles were similar to those observed in the same area during 1974 with primary maxima at depth of a few hundred meters below the surface and secondary maxima near the bottom. Oxidation state distribution profiles were similar at the three locations with plutonium about equally divided between the reduced and oxidized forms except near the bottom. There the oxidized form was more abundant and comprised ~90% of the total plutonium. No major change in oxidation state occurred at the depths of the shallow concentration maxima suggesting that their formation and persistence were not dependent upon a redox change. The concentration maxima near the bottom coincided with the enhanced abundance of the poorly sorbed oxidized form suggesting that they resulted from the loss of oxidized plutonium from the sediments.

In press: Earth and Planetary Science
Letters.

Supported by: DoE Contract
DE-AC02-81EV10694 and Sandia
National Laboratories Contract
16-3112.

WHOI Contribution No. 5481.

THE DISTRIBUTION OF ^{239,240}Pu, ¹³⁷Cs, AND ⁵⁵Fe IN CONTINENTAL MARGIN SEDIMENTS: RELATION TO SEDIMENTARY REDOX ENVIRONMENT

F. L. Sayles and H. D. Livingston

A series of sediments from the East and West Coasts of the U.S. have been analyzed for fallout nuclides in conjunction with redox sensitive parameters in the pore waters of the sediments. Depth distributions do show correlation with redox environments. In particular, depth of penetration of Pu, ¹³⁷Cs and ⁵⁵Fe are greater under more oxidized conditions than in reduced sediment. Modeling of distributions indicates that in a number of sediments dispersion of nuclides at depth is enhanced above that pertaining to the uppermost 2-5 cm of sediment. Comparison of ^{239,240}Pu with ²¹⁰Pb indicates that migration of Pu occurs independently of particle mixing. Modeling of distributions indicates that the occurrence of enhanced migration at depth (below 2-5 cm) may be widespread. A simple model of the migration of a small soluble

fraction of the sedimenting nuclides produces distributions consistent with those observed in the sediments.

In press: Proceedings of the 4th
International Ocean Disposal
Symposium, Plymouth, England.

Supported by: NOAA/Sea Grant.

WHOI Contribution No. 5413.

MEASUREMENT OF CURIUM IN MARINE SAMPLES

David L. Schneider and Hugh D. Livingston

Measurement of environmentally small but detectable amounts of curium requires reliable, accurate, and sensitive analytical methods. The radiochemical separation developed at Woods Hole is briefly reviewed with specific reference to radiochemical interferences in the alpha spectrometric measurement of Cm nuclides and to the relative amounts of interferences expected in different oceanic regimes and sample types. Detection limits for ^{242}Cm and ^{244}Cm are ultimately limited by their presence in the ^{243}Am used as Cm yield monitor. Environmental standard reference materials are evaluated with regard to Cm. The marine literature is reviewed and Cm measurements are discussed in relation to their source of introduction to the environment. Sources include ocean dumping of low-level radioactive wastes and discharges from nuclear fuel reprocessing activities. In particular, the question of a detectable presence of ^{244}Cm in global fallout from nuclear weapons testing is addressed and shown to be essentially negligible. Analyses of Scottish coastal sediments show traces of ^{242}Cm and ^{244}Cm activity which are believed to originate from transport from sources in the Irish Sea.

In press: Nuclear Instruments and
Methods.

Supported by: DoE Contract
DE-AC02-81EV10694 and Sandia
National Laboratories Contract
37-3164.

WHOI Contribution No. 5400.

THE COAGULATION OF DISSOLVED $^{239,240}\text{Pu}$ IN ESTUARIES AS DETERMINED FROM A MIXING EXPERIMENT

Glen T. Shen, Edward R. Sholkovitz
and Don R. Mann

A mixing experiment, using large volumes (100 l) of filtered (<1 μm) fresh-

water and seawater end-members, was performed to simulate and investigate the chemical reactivity of $^{239,240}\text{Pu}$ during estuarine mixing. An organic-rich freshwater with a relatively high concentration (0.8 dpm/100 l) of dissolved $^{239,240}\text{Pu}$ was used as one end-member; Buzzards Bay seawater (dissolved $^{239,240}\text{Pu}$ =0.04 dpm/100 l) was the other. The results demonstrate that dissolved $^{239,240}\text{Pu}$ in the freshwater undergoes extensive and rapid coagulation under simulated estuarine conditions. There is a strong correlation between the amount of coagulation of dissolved $^{239,240}\text{Pu}$, humic acids, and Fe. The extent of coagulation of all three constituents increases with increasing salinity and the net extent of their removal is 53%, 57% and 100% for Pu, HA, and Fe respectively. As has been documented for Fe in freshwater, dissolved $^{239,240}\text{Pu}$ appears to be stabilized by naturally occurring humic substances to form negatively charged colloids which are then coagulated by seawater cations. The extrapolation of these experimental results to real estuaries will require additional research.

Published in: Earth and Planetary
Science Letters, 64, 437-444.

Supported by: DoE Contract
DE-AC02-81EV10694.

WHOI Contribution No. 5325.

REDOX-RELATED GEOCHEMISTRY IN LAKES - ALKALI METALS, ALKALINE EARTH ELEMENTS AND ^{137}Cs

Edward R. Sholkovitz

Redox-driven reactions are important components of the biogeochemical cycles in many lake waters and sediments. The coupling (if any) between redox conditions and the aquatic chemistry of alkali (Na, K, Cs), alkaline-earths (Mg, Ca, Ba), and radioactive ^{137}Cs are discussed. Water column (dissolved and suspended particulate phases) and pore water data are used to show that anoxic conditions indirectly result in enhanced fluxes of alkali and alkaline-earth elements across the sediment/water interface. An important reaction appears to be the release of these elements from an adsorbed phase as the reduction of sedimentary Fe(III) oxides yields the more soluble Fe(II) species under anoxic conditions in lakes. As such, the alkali and alkaline-earth elements undergo a seasonal cycle in the hypolimnion of seasonally anoxic lakes.

Recent studies in lakes are used to hypothesize that anoxic conditions in certain types of lake sediments may result in enhanced mobilities of ^{137}Cs across the sedi-

ment/water interface and within the sediment column. The indirect coupling between ammonia production, which accompanies anoxic conditions, and the exchange of ammonium cations for ^{137}Cs may be an important process in sediments of certain mineralogical compositions. Before any generalities about ^{137}Cs mobility can be made, much more data on ^{137}Cs in anoxic sediments and waters are required.

In press: Chemical Processes in Lakes.
W. Stumm, ed. Wiley Press.

Supported by: DoE Contract
DE-AC02-83ER60172.

WHOI Contribution No. 5536.

THE PORE WATER CHEMISTRY OF $^{239,240}\text{Pu}$
AND ^{137}Cs IN SEDIMENTS
OF BUZZARDS BAY, MASSACHUSETTS

E. R. Sholkovitz and D. R. Mann

The collection of large volumes of pore water (1-2 liter per 2 cm horizon of sediment) and low level radiochemical measurements of $^{239,240}\text{Pu}$ and ^{137}Cs have been combined to produce the first systematic study of these artificial radionuclides in marine pore waters. Profiles from box cores taken in June and September 1982 from Buzzards Bay, MA, are reported along with profiles of many diagenetic constituents (i.e. SO_4^{2-} , alkalinity, Fe, Mn, DOC and nutrients).

The $^{239,240}\text{Pu}$ pore water profile is characterized by a subsurface maximum of about 0.28 dpm/100 kg lying between 3-11 cm. Overlying seawater, in contrast, has an activity of 0.01 ± 0.02 dpm/100 kg. Below about 11 cm, the pore water $^{239,240}\text{Pu}$ distribution follows that of the solid phase which decreases rapidly with depth. The pore water profile of $^{239,240}\text{Pu}$ results from a combination of diffusion and equilibrium with the solid phase. The pore water profiles of ^{137}Cs are characterized by a broad and deeply penetrating maximum where activities of about 42 dpm/100 kg extend from 3 to 20 cm. Overlying seawater, in contrast, has an activity of 21-28 dpm/100 kg. The ^{137}Cs and $^{239,240}\text{Pu}$ pore water data provide direct evidence that there is preferential downward transport of ^{137}Cs and that $^{239,240}\text{Pu}$ does not have an active diagenetic chemistry and is not significantly mobile in these coastal sediments. Incubation studies and flux calculations support these conclusions and point to ^{137}Cs as having a significant mobility.

In press: Geochimica et Cosmochimica
Acta.

Supported by: DoE Contract
DE-AC02-81EV10694.

WHOI Contribution No. 5426.

TRIPOD MODIFICATION OF A SPHINCTER CORER:
CONSTRUCTION, OPERATION, CORE EXTRUSION
AND SAMPLING EFFICIENCY

J. C. Burke, R. E. Hamblin and S. A. Casso

A 21-cm diameter sphincter corer (Burke, 1968) has been modified by mounting it in a tripod frame. This modification results in more dependable recovery of undisturbed surficial sediment and greater penetration into firm sediment. The device is useful in water depths from 2-6000 m, is adaptable for use on small boats, is very easily employed on large research vessels, and can be readily disassembled for more convenient transporting and storing.

In press: WHOI Technical Report.

Supported by: DoE Contract
EY-76-C-3563; Sandia National
Laboratories Contract 37-3164; and
Nuclear Regulatory Commission
Contract NRC-04-76-349.

DEPARTMENT OF GEOLOGY AND GEOPHYSICS
Richard P. von Herzen, Department Chairman

G E O L O G Y

NEOGENE GEOCHRONOLOGY AND CHRONOSTRATIGRAPHY

William A. Berggren, Dennis V. Kent
and John A. Van Couvering

We present a revised Neogene geochronology based upon a best fit to selected high temperature radiometric dates on a number of identified magnetic polarity chrons (within the late Cretaceous, Paleocene, and Neogene) which minimizes apparent accelerations in seafloor spreading. An assessment of first order correlations of calcareous plankton biostratigraphic datum events to magnetic polarity stratigraphy yields the following estimated magnetobiochronology of major chronostratigraphic boundaries: Oligocene/Miocene (mid-anomaly 6C): 23.7 Ma; Miocene/Pliocene (slightly younger than Gilbert/Chron 5 boundary): 5.3 Ma; Pliocene/Pleistocene (slightly younger than Olduvai Event): 1.6 Ma.

Changes to the marine time scale are relatively minor in terms of recent and current usage. Radiometric dates correlated to bio- and magnetostratigraphy in continental sections support the revised marine magnetobiochronology presented here. Major changes, however, are made in marine - non-marine correlations in the Miocene in Eurasia which indicate African-Eurasian migrations through the Persian Gulf as early as 20 Ma. The 12 Ma estimate of the Hipparion datum is supported by recent taxonomic revisions of mammals in magnetostratigraphically studied sections in the Siwaliks of Pakistan.

In press: Geological Society of
London Special Paper.

Supported by: NSF Grants OCE80-23728
and OCE80-19052; and the Senior
Studies Award.

WHOI Contribution No. 5496.

MORPHOLOGY, SEDIMENTS AND LATE QUATERNARY HISTORY OF THE EAST CHINA SEA

Jorge Butenko, Ye Yincin
and John D. Milliman

Surficial sediments and fine-scale morphology in the Chang Jiang-East China Sea region directly reflect a complex interaction between Late Quaternary history, sediment input and oceanographic conditions. Fine-grained sediments on the inner shelf are derived mainly from the Chang Jiang in the south and Huangho in the

north; rates of accumulation locally exceed 4 cm/yr. The resulting topography is generally smooth and featureless, although local pock marks from biogenic gas blow outs are noted. The middle shelf is dominated by relict sands which are partly to completely rippled (local occurrence of megaripples), indicating physical reworking of sediments by waves and tidal currents. The outer shelf south of Cheju Island is covered with modern fine-grained muds and muddy sands, apparently derived from the Huangho.

The East China Sea is underlain by a network of late Pleistocene-early Holocene drainage channels formed by the Chang Jiang and Huangho as well as other rivers draining the mainland. In the area of study these channels are overlain by as much as 20 m of late Holocene sediment. Thicknesses are greatest in the nearshore area south of the Chang Jiang, where the modern submarine delta is prograding over a relict delta.

Published in: Proceedings to the International Symposium on Sedimentation on the Continental Shelf. China Ocean Press.

Supported by: ONR Contract N00014-81-C-009 and NOAA Cooperative Agreement No. NA81AA-H-00008.

WHOI Contribution No. 5356.

MAGNETIC CHRON 12R AT FLAGSTAFF RIM, WYOMING

Charles R. Denham

Early Oligocene magnetic Chron 12R is expressed over a 79 m interval in radio-metrically-dated, fossiliferous Chadronian strata at Flagstaff Rim, Wyoming. The Chron 12R/13N boundary occurs at Ash F, dated 34.6 Ma (Evernden et al., 1964). The stepwise AF demagnetization data were interpreted by an objective method to obtain statistically significant polarities based on the latitudes of the corresponding virtual geomagnetic poles.

Supported by: NSF Grants DES76-82255
and NSF DES78-23687.

WHOI Contribution No. 5390.

TECTONICS AT THE INTERSECTION
OF THE EAST PACIFIC RISE
WITH THE TAMAYO TRANSFORM FAULT

D.G. Gallo, W.S.F. Kidd, P.J. Fox,
J.A. Karson, K. Macdonald, K. Crane,
P. Choukroune, M. Seguret, R. Moody
and K. Kastens

During the Fall of 1979, a manned submersible program, utilizing DSRV ALVIN, was carried out at the intersection of the East Pacific Rise (EPR) with the Tamayo Transform boundary. A total of seven dives were completed in the vicinity of the EPR/Tamayo intersection depression and documented the geologic relationships that characterize the juxtaposition of these types of plate boundaries. The young volcanic terrain of the EPR axis can be traced into and across the Tamayo Transform valley but becomes rapidly buried by sedimentary talus that is being shed from sediment scarps along an unstable sediment slope that defines the north side of the intersection depression. Within 4 km of the transform boundary, the dominant trend (000°) of the fissures and faults that disrupt the rise generated volcanics are markedly oblique to the regional direction of sea floor spreading (120°). Since no evidence was found to suggest that these structures accommodate significant amounts of strike-slip displacement, they are taken to reflect a distortion of the EPR extensional tectonic regime by a transform generated shear couple. The floor of the Tamayo Transform valley in this area is inundated by mass-wasted sediment, and the principal transform displacement zone is characterized by a narrow (<1.5km) interval of deformed sediment that trends parallel with the transform valley and, if extrapolated to the west, links with zones of transform deformation investigated during earlier submersible studies (CYAMEX and Pastouret, 1981). Evidence of low-level hydrothermal discharge was seen at one locality on the EPR axis and another 8 km west of the axis at the edge of the zone of transform deformation.

In press: Marine Geophysical
Researches.

Supported by: NSF Grant OCE79-13144.

WHOI Contribution No. 5488.

SEDIMENTATION UNDER DEEP SEA-STORMS

Charles D. Hollister and Ian N. McCave

Fast currents, high concentrations of suspended sediment and grooved mud beds are attributed to erosion in frequent abyssal

storms where a fast deep mean flow is augmented (or reversed) by intense intermittent currents. This occurs about 5 km below the Gulf Stream or its rings. The waning phase of a storm results in development of bedforms and rapid deposition of a mud blanket. Several other regions of the world ocean display evidence of abyssal storm activity.

In press: Nature.

Supported by: ONR Contract N00014-82-C-0019.

WHOI Contribution No. 5570.

ABYSSAL TELECONNECTIONS II.
INITIATION OF ANTARCTIC BOTTOM WATER
FLOW IN THE SOUTHWESTERN ATLANTIC

David A. Johnson

Three principal factors are potentially important in governing the initiation of Antarctic Bottom Water (AABW) flow through the deep ocean basins during the Paleogene, and subsequent fluctuations in its intensity during the Neogene and Quaternary: (a) The beginning of circumpolar flow, leading to thermal isolation of the Antarctic continent and decreasing surface water temperatures around the Antarctic perimeter; (b) Injection of high-salinity waters into the Antarctic region at intermediate depths; and (c) Rifting and subsidence of sills and passages below critical threshold depths. Of these factors, salinity has most probably been the limiting one. Formation of bottom water in the northern Atlantic began near the Eocene/Oligocene boundary via overflow from the Norwegian Sea through deepening sills in the Iceland-Scotland Ridge. This overflow, or proto-North Atlantic Deep Water (NADW), injected high-salinity water into the central and southern Atlantic at intermediate depths and was probably essential to AABW production. A prominent seismic discontinuity (Horizon A of Ewing) marks the initiation of AABW flow through the southwestern Atlantic; drilling through this discontinuity in the Brazil Basin at DSDP Site 515 shows thick, rapidly-accumulating (<20 m/m.y.) bottom current deposits beginning in the middle Oligocene (32 Ma) and continuing through the Quaternary. The effects of AABW in the deep North Atlantic were also well established by the middle Oligocene (~30 to 33 Ma), with the advection of older and ¹³C-depleted bottom waters as far north as the Bay of Biscay. Beginning in the middle Miocene (~16 Ma), an alternative source region developed in the northwestern Indian Ocean for supplying high salinity waters to high southern latitudes. The gradual closing of the

eastern Tethyan seaways during the Neogene has dramatically altered the evaporation/precipitation budget of southwest Asia, allowing the Arabian Sea and its marginal basins to become source areas for southward-flowing hypersaline intermediate and deep waters. During the climatic cycles of the late Neogene and Quaternary, the most recent episodes of intense AABW flow appear to precede the strongest glacial maxima at 20 Kyr and 150 Kyr BP. These episodes are not directly associated with maxima in NADW production which may occur at a 40 Kyr period. Thus the production of high salinity deep water in the northernmost Atlantic (i.e., NADW) is not necessarily the only precondition to the formation of AABW, provided that an alternative source of hypersaline water (e.g., the north-western Indian Ocean) is available.

In press: Cambridge University Press
(symposium volume).

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5489.

THE VEMA CHANNEL: PHYSIOGRAPHY, STRUCTURE AND SEDIMENT-CURRENT INTERACTIONS

David A. Johnson

The east flank of the Vema Channel in the southwestern Atlantic is an opportune location for interpreting the Pleistocene/Holocene record of deep thermohaline circulation. The channel and the adjacent sediment-covered Rio Grande Rise intersect each of the principal sub-thermocline water masses of the western Atlantic. We analyzed a suite of 25 giant gravity cores in transects extending between 1400 m and 4150 m depth; this interval includes the upper and lower branches of Circumpolar Water, North Atlantic Deep Water, and Antarctic Bottom Water. The late Pleistocene/Holocene sediments along these transects are pelagic carbonates with excellent stratigraphic continuity; shallower and deeper regions are relatively unsuitable due to poor core recovery or widespread erosional unconformities. Extensive hydrographic and direct current observations near the location of our prime coring transect at 31°S indicate a notable cross-channel asymmetry in the abyssal hydrography and velocity structure which is reflected in the underlying lithofacies distribution. The near-bottom flow field shows marked temporal and spatial variability on a local scale, even in the channel axis where one would anticipate relatively steady, uniform flow. Much of this variability appears to reflect the effects of transient mesoscale eddies rather than regional variations in the integrated

northward transport of AABW. In spite of temporal variations in the flow field, the NADW/AABW transition zone appears to have remained remarkably stable over time scales of 10^5 to 10^6 years, based on geological evidence discussed elsewhere in this volume (Jones, 1984; Jones and Johnson, 1984). This observation is somewhat surprising, because in the modern oceans the density contrast between NADW and AABW is relatively small. If the transition zone between the two water masses has indeed remained relatively stationary, then the density-determining water properties (T and S) may also have remained relatively constant, with only the rates of production changing. If this interpretation proves correct upon further studies, it implies that paleoceanographic interpretations of deep thermohaline flow should in some instances utilize geological indices of non-conservative water properties (e.g. CO_2 , $\delta^{13}\text{C}$) for monitoring paleo-circulation changes.

In press: Marine Geology.

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5490.

DISPLACED ANTARCTIC DIATOMS IN VEMA CHANNEL SEDIMENTS: LATE PLEISTOCENE/ HOLOCENE FLUCTUATIONS IN AABW FLOW

Glenn A. Jones and David A. Johnson

Displaced Antarctic diatoms were identified and counted in a suite of 23 giant gravity cores from the east flank of the Vema Channel, providing a semi-quantitative index of Late Pleistocene/Holocene fluctuations in Antarctic Bottom Water (AABW) flow. Twelve species of Antarctic diatoms were counted, and three (*Nitzschia kerguelensis*, *Eucampia antarctica*, and *Coscinodiscus lentiginosus*) occurred in sufficient quantity for statistically meaningful interpretations. The number of diatom specimens per gram of carbonate-free sediment was determined for core top, last glacial maximum, last interglacial, and penultimate glacial maximum intervals. Antarctic diatoms are abundant in cores from depths in excess of 4100 m and disappear abruptly in core tops between approximately 4125 and 3975 meters water depth. No specimens of Antarctic diatoms were found in cores shallower than 3744 m. The top of this transition zone corresponds with the isotherm $\theta = 1.9^\circ\text{C}$ and the isopycnal $\sigma_t = 45.93$ that Hogg et al. (1982) selected as a suitable reference level between northward and southward flow in the Vema Channel axis. No significant shallowing or deepening of this level is found in the sediments from the Vema Channel during

any of the glacial or interglacial intervals examined. In general, displaced Antarctic diatoms are less abundant during glacial than during interglacial periods. Down-core counts of displaced Antarctic diatoms reveal two peaks of dramatically increased abundances at the isotopic stage 6/7 and 3/2 boundaries. In other cores within the Vema Channel from below 3965 m, the incidence of hiatuses increases and is inferred that these hiatuses are the result of increased flow velocities of AABW.

In press: Marine Geology.

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5527.

GEOLOGIC AND SEISMIC VELOCITY STRUCTURE OF THE CRUST/MANTLE TRANSITION IN THE BAY OF ISLANDS OPHIOLITE COMPLEX

Jeffrey A. Karson, John A. Collins and J.F. Casey

Geological investigations of the Bay of Islands Ophiolite Complex show that while a typical layered ophiolite suite is present, the thickness of major lithologic units is extremely variable from place to place. The composition and internal structure of map-scale (kilometers across) lithologic units and the contacts which bound them are also laterally variable. Inferred velocity-depth functions of the crust/mantle ('MOHO') transition reconstructed for this terrane as oceanic lithosphere suggest an extremely complex internal seismic structure. The crust/mantle transition from mafic to ultramafic lithologies varies across the ophiolite from a sharp geologic and seismic velocity discontinuity to a complexly inter-layered transition zone as much as 3 km thick. The mafic/ultramafic transition is characterized by laterally discontinuous lithologic units on the order of hundreds of meters to a few kilometers thick and up to several kilometers long. These units (megasequences) may be composed of lithologies with higher or lower seismic velocities than those of the surrounding units. Solid-state deformation has produced seismically anisotropic materials in the upper mantle and lower crustal units. Substantial relief and slopes of up to at least 12° over lateral distances of about 10 km occur on the top of the upper mantle as well as other geologic unit contacts. Although these types of geological features are striking to the field geologist working in ophiolites, many of them occur on such a small scale that they might go undetected in different types of seismic experiments in contemporary oceanic lithosphere.

In press: Journal of Geophysical Research (Red).

Supported by: NSF Grants EAR80-26445 and OCE80-025206.

WHOI Contribution No. 5560.

GEOLOGY OF THE OCEANOGRAPHER TRANSFORM: THE RIDGE-TRANSFORM INTERSECTION

J.A. Karson, P.J. Fox, K. Crane, H. Sloane, E. Bonatti, W. Kidd, J. Stroup, D. Fornari, D. Elthon, P. Hamlyn, J. Casey, D. Gallo, D. Needham and R. Sartori

Seven dives in the submersible ALVIN and four deep-towed camera lowerings (ANGUS) have been made at the eastern ridge-transform intersection of the Oceanographer transform with a segment of the Mid-Atlantic Ridge. These data constrain our understanding of the processes that create and shape the distinctive morphology that is characteristic of slowly-slipping ridge-transform ridge plate boundaries. Although the geology of the rift valley floor in the study area is similar to relationships reported for the FAMOUS area, we observe a distinct change in the character of the rift valley floor with increasing proximity to the transform. Over a distance of approximately 10 kilometers the volcanic constructional terrain becomes increasingly more disrupted by faulting and degraded by mass wasting. Moreover, proximal to the transform boundary faults with orientations oblique to the trend of the rift valley and parallel to the transform are recognized. The morphology of the eastern rift valley wall is characterized by inward-facing scarps that are ridge-axis parallel but the western rift valley wall, which faces the transform, is characterized by a complex fault pattern defined by faults exhibiting a range of orientations (Fig. 3). Irrespective of orientation, however, no evidence for strike-slip displacement is observed throughout the rift valley and evidence for normal (dip-slip) displacements is ubiquitous regardless of scarp orientation. Basalts, semi-consolidated sediments (carbonate, debris fans) and serpentinized ultramafic rocks are recovered from localities within the rift valley. The axis of accretion - principal transform displacement zone intersection is not clearly established but appears to be located along the E-W trending south flank of the intersection deep (Fig. 8).

In press: Marine Geophysical Researches.

Supported by: ONR Contract N00014-81-C-0820.

WHOI Contribution No. 5351.

TECTONICS OF RIDGE-TRANSFORM INTERSECTIONS
AT THE KANE FRACTURE ZONE

J.A. Karson and H.J.B. Dick

The Kane Transform offsets spreading-center segments of the Mid-Atlantic Ridge by about 150 km at 24°N latitude. In terms of its first-order morphological, geological, and geophysical characteristics it appears to be typical of long-offset (>100 km), slow-slipping (2 cm/yr) ridge-ridge transform faults. High-resolution geological observations were made from deep-towed ANGUS photographs and the manned submersible ALVIN at the ridge-transform intersections and indicate similar relationships in these two regions. These data indicate that over a distance of about 20 km, as the spreading axes approaches the fracture zone, the two flanks of each ridge axis behave in very different ways. Along the flanks that intersect the active transform zone, the rift valley floor deepens and the surface expression of volcanism becomes increasingly narrow and eventually absent at the intersection where only a sediment-covered 'nodal basin' exists. The adjacent median valley walls have structural trends that are oblique to both the ridge and the transform and have as much as 4 km of relief. These are tectonically active regions that have only a thin (<200 m), highly fractured, and discontinuous carapace of volcanic rocks overlying a variably deformed and metamorphosed assemblage of gabbroic rocks. Overprinting relationships reveal a complex history of crustal extension and rapid vertical uplift. In contrast, the opposing flanks of the ridge axes that intersect the non-transform zones appear to be similar in many respects to those examined elsewhere along slow-spreading ridges. In general, a near-axial horst and graben terrain floored by relatively young volcanics passes laterally into median valley walls with a simple block-faulted character where only volcanic rocks have been found. Along strike toward the fracture zone, the youngest volcanics extend into constructional volcanic ridges that transect the entire width of the fracture zone valley. These volcanics are continuous with the older-looking, slightly faulted volcanic terrain that floors the non-transform fracture zone valleys. These observations document the asymmetric nature of seafloor spreading near ridge-transform intersections. An important implication is that the crust and lithosphere across different portions of the fracture zone will have different geological characteristics. Across the active transform zone, two lithosphere plate edges formed at ridge-transform corners are faulted against one another. In the non-transform zones, a relatively younger section of lithosphere that formed at a ridge-non-transform corner is welded to an

older, deformed section that initially formed at a ridge-transform corner.

Published in: Marine Geophysical
Researches, 6, 51-98.

Supported by: NSF Grants OCE77-
09155, OCE77-26842, OCE79-26302 and
OCE80-24930.

WHOI Contribution No. 5343.

HIGH RESOLUTION ISOTOPIC STUDY OF THE
LATEST DEGLACIATION FROM BERMUDA RISE CORES

L. D. Keigwin, Bruce H. Corliss, Ellen M.
Druffel and E. P. Laine

Stable oxygen and carbon isotopic results and radiocarbon results from three cores on the Bermuda Rise reveal the most detailed record of deglaciation yet obtained from deep sea cores. High resolution study of these cores is possible because deep sea currents in this region maintain high sedimentation rates through transport and deposition of clay and silt whereas the sand fraction, mostly planktonic foraminifera, accumulates slowly from overlying surface waters. Radiocarbon age determinations on bulk CaCO_3 yield Holocene sedimentation rates of $\sim 25 \text{ cm}/10^3 \text{ yrs}$. Sedimentation rates are estimated to be $41 \text{ cm}/10^3 \text{ yrs}$ for the interval 16 to 75,000 yrs and as high as $200 \text{ cm}/10^3 \text{ yrs}$ during the latest glacial maximum.

In core GPC-5 oxygen isotopic results on the planktonic foraminiferal species Globorotalia inflata and Globigerinoides ruber follow similar patterns, suggesting they record changes in the volume of continental ice as well as local temperature and salinity changes. Maximum $\delta^{18}\text{O}$ values occurred $\sim 19,000$ yrs ago and proceeded to Holocene values ($\sim 1.75^\circ/\text{oo}$ lower) in at least two steps, which probably correlate with isotopic features within the deglaciation seen elsewhere in the North Atlantic. All three oxygen isotopic records from the Bermuda Rise contain an abrupt interval enriched by $1^\circ/\text{oo}$ in ^{16}O immediately following maximum values of the glaciation. This is tentatively interpreted as a melt-water pulse from sea ice and icebergs in the northern North Atlantic.

Supported by: Sandia Contract
26/6605.

WHOI Contribution No. 5422.

L. D. Keigwin, S.V. Margolis
and B.U. Haq

Samples from an 8 m, 400,000 year interval (~6.1 to 6.5 MY) in the latest Miocene of DSDP Site 158 (Cocos Ridge, Panama Basin) have been examined at 5,000 to 10,000 year spacing to study the detail of stable isotopic, faunal and floral change. At this location the Late Miocene carbon isotopic shift in benthic foraminifera occurred in about 75,000 years. At the time of the carbon shift (~6.2 MY) planktonic foraminiferal $\delta^{13}\text{C}$ decreases but nannofossil $\delta^{13}\text{C}$ increases, and nannofossil $\delta^{18}\text{O}$ increases but planktonic foraminiferal $\delta^{18}\text{O}$ does not change. Oxygen isotopic results are consistent with a surface water cooling of 2°C and migration upward in the water column by planktonic foraminifera. Faunal and floral evidence likewise indicates cooler surface water temperatures as well as increased CaCO_3 dissolution beginning at the carbon shift. Benthic foraminiferal $\delta^{18}\text{O}$ does not correlate with the planktonic $\delta^{18}\text{O}$ records, suggesting a greater influence of temperature over ice volume effects.

Supported by: NSF Grants OCE79-14594
and OCE81-17590.

WHOI Contribution No. 5242.

OLIGOCENE CALIBRATION OF THE
MAGNETIC POLARITY TIMESCALE

Donald R. Prothero, Charles R. Denham
and Harlow G. Farmer

Magnetostratigraphic studies of the Oligocene White River Group in Wyoming, Colorado, Nebraska, and the Dakotas yield a radiometrically dated polarity stratigraphy that provides mid-Tertiary calibration points for the magnetic polarity time scale. Anomaly 12-13 reversal is bracketed by dates of 32.4 and 34.6 m.y., in best agreement with the time scale of LaBrecque and colleagues. The magnetostratigraphy also helps calibrate the Oligocene North American land mammal "ages" and allows correlation with the European marine microfossil zonation. This correlation suggests that the age of the Eocene-Oligocene boundary is 37.0 m.y., contrary to younger dates obtained from glauconites and microtektites.

Published in: *Geology*, 10, p. 630-653.

Supported by: NSF Grants EAR76-82255
and EAR78-23687.

WHOI Contribution No. 5321.

David A. Ross

The Red Sea is situated between the Arabian Peninsula and the African continent and is an elongated basin that is being formed by sea-floor spreading. Bottom topography consists of smooth shelves separated by a large, central main trough that itself has a deep and deformed axial valley. It is bounded by topographically high areas that influence the weather and generally restrict atmospheric circulation to winds parallel to the axis of the sea. Climate is arid and hot and evaporation is of the order of 2000 mm yr^{-1} . North of 19°N , the winds are generally from a north to northeasterly direction; to the south, their direction changes according to the monsoons. From October to May, winds are mainly from the southeast, while during June to September, they are mainly from the northwest. The combination of climate, low river runoff and rainfall, high evaporation rate and restricted circulation has produced relatively dense, high temperature (up to 30°C) and high salinity ($40^\circ/\text{oo}$ or more) water.

The Red Sea is about 1800 km long; about 250 km in average width and about 560 m in average depth, with a maximum depth of 2920 m. Thus, the surface area is about $450,000 \text{ km}^2$ and the volume of the Red Sea is about $251,000 \text{ km}^3$ (Muromtsev, 1960). It is connected to the Mediterranean Sea via the Gulf of Suez and the Suez Canal (although relatively little water exchange occurs there), and to the Indian Ocean via the Strait of Bab al Mandab.

The general circulation pattern of the Red Sea shows very distinct seasonal changes that are influenced primarily by prevailing winds and to a lesser extent by seasonal changes in density and evaporation. In the winter (October-May) surface waters from the Gulf of Aden flow to the north and the subsurface flow is to the south. This pattern reverses in the summer. Upwelling occurs in the northern part of the sea during the summer months, and sinking of dense water occurs in the winter period. The renewal time for the upper 150 m of Red Sea water may be as little as six years, whereas, for the entire Red Sea, a value of about 200 years may be more appropriate.

Published in: *Estuaries and Enclosed Seas*. B.H. Ketchum, ed. Elsevier Scientific Publishing Co., Amsterdam, 1983.

Supported by: ONR Contract N00014-74-C-0262.

WHOI Contribution No. 4092.

CENOZOIC REGIONAL EROSION OF THE
ABYSSAL SEAFLOOR OFF SOUTH AFRICA

Brian E. Tucholke and Robert W. Embley

The Cape, Agulhas, and Mozambique Basins off South Africa have a well defined and mostly continuous erosional zone along their perimeters between 4 and 5 km water depth. This zone lies beneath a deep boundary current of Antarctic Bottom Water. However, current speeds are generally less than 15-20 cm/s and part of the erosional zone is armored by authigenic manganese deposits, so that only limited erosion presently is occurring. Erosion and corrosion of sea-floor sediments by abyssal currents probably are in dynamic equilibrium with sediment supply. The present erosional zone is largely a relict feature inherited from late Miocene time when strongly increased glaciation of West Antarctica produced large volumes of bottom water that scoured the sea-floor. A deeper unconformity dating to the early Oligocene marks the onset of significant abyssal circulation in the basins, and current-controlled deposition of sediments is well defined above this unconformity. In contrast, an underlying, basal Eocene unconformity shows no marked effects of control by abyssal circulation, although erosion/corrosion by weak, deep currents could have occurred. The unconformity formed primarily because of reduced sediment supply caused by elevated sea level and probably low productivity in surface waters.

In press: AAPG Memoir.

Supported by: ONR Contract N00014-79-C-0071 and NSF Grant OCE76-21782.

WHOI Contribution No. 5331.

ABYSSAL CURRENT CHARACTER DETERMINED
FROM SEDIMENT BEDFORMS
ON THE NOVA SCOTIAN CONTINENTAL RISE

B.E. Tucholke and C.D. Hollister

Bottom photographs across the Nova Scotian continental rise show effects of an intense bottom current at 4800-5000 m that decreases in strength upslope to tranquil near-bottom conditions at 3200-4000 m. Apparently weak currents are present above 3200 m. Long-term, geologically significant flow parallels the bathymetric contours toward the southwest, but significant short-term variability in direction is indicated by orientations of small-scale bedforms. The Western Boundary Undercurrent, documented elsewhere along the continental margin at depths less than 4000 m, either is poorly developed on the Nova Scotian rise or has relatively little effect in

modifying the sea floor there.

Repeated photography of small sea-floor areas suggests that many bedforms, especially those of smaller scale, are constructed and destroyed by strong, variably directed, current events ("benthic storms") on time scales of 3 to 6 months or less. Radiometric studies of short-lived isotopes in the sediments agree with this interpretation. Thus, a significant part of the geologic response of the seabed to currents, particularly along western margins of basins, may be event-controlled on time scales not very different from those in coastal environments.

Supported by: ONR Contracts N00014-79-C-0071 and N00014-82-C-0019.

WHOI Contribution No. 5158.

G E O P H Y S I C S

DEPTH OF PRINCIPAL MASS ANOMALIES
CONTRIBUTING TO THE EARTH'S GEODAL
UNDULATIONS AND GRAVITY ANOMALIES

Carl Bowin

To help decipher the distribution of the principal mass anomalies within the earth, an analysis is made of the earth's potential field by simultaneously considering both gravity and geoid anomalies. Ancillary knowledge about the earth from astronomical, angular momentum, seismological, and plate tectonic studies aids in deciding between deep or broad shallow-mass anomalies as the causative source of certain potential field anomalies.

The geoid (or gravity) anomaly for an individual gravity potential field feature as a function of increasing spatial resolution (decreasing wavelength) is obtained by accumulating stepwise the contributions of spherical harmonic coefficients for each degree and order for any selected location. This is equivalent to displaying the spectral content of an anomaly. The ratio of gravity to geoid (g/N) for each individual harmonic degree is independent of the harmonic coefficient values, being determined only by the degree and values for normal gravity and the earth's radius. This g/N ratio value has an equivalent point mass depth. Thus, the gravity and geoid contributions from each individual degree have a fixed ratio value in proportion to that of point masses at a depth of some fraction of the earth's radius, and this depth places a greater constraint on the depth of the source mass anomaly than can be obtained from wavelength considerations. Curves are presented that quantify the g/N relation-

ships for two simple geometric degrees 2 and 3 geoid and gravity anomalies.

A preliminary decomposition of the earth's gravity field is made using these new considerations. This decomposition comprises four main sources for mass anomalies: (1) mass anomalies at the core-mantle boundary region expressed principally in coefficients of harmonic degrees 2* and 3*; (2) mantle anomalies, and the mass excess of the deeper portions of plate convergent zones, expressed principally in coefficients of harmonic degrees 4* through 10*; (3) upper mantle mass anomalies in the outer 600 km of the earth revealed by residual geoid anomalies from radar altimeter observations; and (4) crustal anomalies revealed by residual free-air gravity anomalies. The observational data cited do not distinguish perfectly between these four sources of mass anomalies as well as between other possible lesser contributors. However, the evidence is sufficiently reasonable to give credence to the decomposition. The degree 10 geoid thus provides an estimate of the combined contributions from mass anomalies at the core-mantle boundary region and from the deep mantle and deeper parts of plate convergent zones. Its subtraction from radar altimeter data indicates mass anomalies in the outer 600 km, which in turn provides new information regarding the nature of convection within the upper mantle.

Published in: Marine Geodesy, 7, 61-100, 1983.

Supported by: ONR Contract N00014-79-C-0071.

WHOI Contribution No. 4612.

GLOBAL GRAVITY MAPS AND THE STRUCTURE OF THE EARTH

Carl Bowin

It is now possible to view global maps depicting long-wavelength and short-wavelength features of the earth's gravity potential field, both as geoid anomalies and as free-air anomalies. This progress has resulted from (1) an increased number of surface gravity measurements, (2) orbiting of artificial satellites, (3) installation of radar altimeters in three spacecraft, and (4) improved computing and display capabilities. The ratio of gravity to geoid anomalies g/N has been utilized to suggest a decomposition of the earth's gravity field. The g/N ratio for each individual spherical harmonic degree is independent of the harmonic coefficient values, being determined only the degree and values for normal gravity and the earth's radius. This g/N ratio value has an associated point mass

depth that is more limiting of source depth than that provided by wavelength considerations. Mass anomalies that may exist in the earth at great depth will be best represented in the coefficients of low degree spherical harmonics, not only because of their longer wavelength anomalies at the surface, but also to better match the resulting ratio of gravity to geoid at the center of the anomaly feature.

Shallow (less than 600 km) broad masses that might be considered possible sources of the degree 2* and 3* Sri Lanka geoid low and the New Guinea geoid high are considered geologically unlikely, whereas 1 to 2 km of relief on the core mantle boundary is plausible and can explain the combined degree 2 and 3 geoid and gravity anomalies. The core mantle boundary may become warped as a result of stresses developed by hydrodynamic motions in the outer part of the core, possibly related to those that produce the earth's magnetic field. Combined contributions from harmonic degrees 4 through 10 identify a narrow positive mass anomaly beneath plate convergent zones. These positive anomalies are not due to the anomaly of a downgoing slab because they are much broader and have larger magnitude. The fact that association of this positive mass anomaly with plate convergent zones is more sharply identified in the selected packet of harmonic coefficients than in the full field is a strong indication that the preliminary decomposition is proceeding in the correct direction.

Cumulative geoid degree contribution curves for the low harmonic terms of the gravity field of Venus have a different pattern than that for the earth. Maps of the low harmonic coefficients for Venus indicate correspondence with surface topographic features in marked contrast to the earth. Venus rotates very slowly and has no internal magnetic field. Perhaps coherent motion within the core is negligible, so that there is no magnetic field and no distortion of the core-mantle boundary. These relations support the conclusion that long-wavelength undulations at the earth's core-mantle boundary may contribute significantly to the lowest-degree harmonic coefficients of the earth's potential field.

In press: Society of Exploration
Physicists.

Supported by: ONR Contract N00014-79-C-0071.

WHOI Contribution No. 5549.

GRAVITY, TOPOGRAPHY AND
CRUSTAL EVOLUTION OF VENUS

Carl Bowin

The gravity anomalies of Venus, although small by comparison with those on Mars and the Moon, are still much larger than those on Earth for large features. On Venus, even the low-degree spherical harmonic terms for the Venusian gravity field indicate a close association of broad positive gravity anomalies with major topographic highs. This is in striking contrast to the situation on Earth, where the broad regional gravity anomalies show little correlation with continental masses or plate tectonic features, but instead appear to be caused by deep mass anomalies.

A method for estimating radial gravity anomalies from line-of-sight acceleration data, their interpolation, and use of iteration for improved radial anomaly estimates is outlined. A preliminary gravity anomaly map of Venus at spacecraft altitude prepared using first estimate values is presented. A profile across the western part of Aphrodite along longitude 85°E was analyzed using time-series techniques. An elastic plate model would require a plate thickness of about 180 to 200 km to match the general amplitude of the observed gravity anomaly (about 33 mgal) -- a thickness much greater than that found for earth structures and, because of high surface temperatures, unlikely for Venus. An Airy isostatic model convolved with the topography across Aphrodite, however, provides a better match between the predicted and observed gravity anomalies if the nominal crustal thickness is about 70 to 80 km. This thickness is over twice that for continental crust on the earth and considerably greater than that of the earth's basaltic ocean crust (only 5 km). Thus, a different differentiation history for Venus than that of the earth is anticipated.

High gravity anomalies (+110 mgal) occur over Beta Regio and over the topographic high in eastern Aphrodite; both highs are associated with regions where detected lightning is clustered, and thus, the topographic features may be active volcanic constructs. The large gravity anomalies at these two sites of volcanic activity require an explanation different than that indicated for western Aphrodite.

Published in: *Icarus*, 56, 345-371,
1983.

Supported by: NASA Contract NAG2-91.

WHOI Contribution No. 5485.

STRUCTURE AND VARIABILITY OF OCEANIC
LITHOSPHERE ON THE FLANKS OF THE
EAST PACIFIC RISE BETWEEN 11° and 13°W

Steven R. Bratt and G.M. Purdy

We use over 500 seismograms collected by five ocean-bottom seismometers to examine the structure and variability of 0.5 m.y. old crust in the ROSE area of the East Pacific Rise between 11° and 13°N. Though there are no significant differences between the ten first arrival travel-time curves from the rise-flanking line, amplitude patterns recorded on the individual lines suggest that the structure of the crust varies by hundreds of meters over lateral distances of tens of kilometers under this 200 km segment of lithosphere. Based on the modeling of amplitude patterns at ranges of 5 to 10 km and on comparisons with the upper crust sampled at DSDP hole 504B and with the structure of ophiolite complexes, layer 2 can be divided into A, B, and C units which correspond to: A) extrusive volcanics with a steep velocity gradient, B) extrusives with a near-zero velocity gradient, and C) a complex region of transition from extrusives to sheeted dikes. The total thickness of layer 2 varies by about 0.7 km over the length of the line. Another amplitude peak which varies in range between 14 and 21 km may be related to the transition from sheeted dikes to isotropic gabbros within layer 3. The variability of these two amplitude patterns provides direct evidence that the along strike structure of oceanic crust in the ROSE area is definably heterogeneous and that the process of crustal accretion is likewise variable on the scale resolved by amplitude and travel-time information.

In press: *Journal of Geophysical Research*.

Supported by: ONR Contracts N00014-79-C-0071 and N00014-82-C-0019.

WHOI Contribution No. 5550.

SEISMIC REFLECTION PROFILING WITH A DEEP-
TOWED HYDROPHONE ARRAY

John A. Collins, Donald E. Koelsch,
G. M. Purdy

A seismic reflection profiling system consisting of a 264 m long, deep-towed, 15-element, vertical array and a 40 cubic inch airgun was successfully used to profile a sediment pond in the trough of the inactive segment of the Kane Fracture Zone close to its intersection with the Mid-Atlantic Ridge. The increased signal to noise ratio achieved with the array suggests that it is

a useful tool for detailed seismic profiling in areas of rough topography.

In Press: Marine Geophysical
Researches.

Supported by: NSF Grant OCE80-025206.

WHOI Contribution No. 5443.

MODELING PALEOMAGNETIC REMANENCE

Charles R. Denham and Loren Shure

We have modeled paleomagnetic remanence from its stepwise demagnetization behavior by using arbitrary magnetization decay functions that are constrained only to be monotonically-decreasing and non-intersecting with increasing demagnetizing field strength. In comparison with the two-vector exponential model of Stupavsky and Symons (1978), our weakly-constrained decomposition better describes the remanence. The two-vector weakly-constrained model is illustrated for some actual data where the demagnetization behavior is substantially non-exponential.

Supported by: NSF Grants DES 76-82255
and 78-23687; and ONR Contract
N00014-82-C-0019.

WHOI Contribution No. 5435.

THE AMERICA-ANTARCTIC RIDGE

Lawrence A. Lawver and Henry J.B. Dick

The America-Antarctic Ridge forms the boundary between the South America and Antarctic plates in the South Atlantic. A recent cruise on R/V MELVILLE surveyed and dredged the America-Antarctic Ridge between the Conrad Fracture Zone at 56°S, 4°W and a spreading center at 60°S, 18.5°W. This section of the plate boundary consists of five transform faults and six spreading center segments. Prior to our work, the plate boundary was generalized on the basis of seismicity and was thought to consist of only two or three spreading centers offset by as many transform faults. Magnetic anomaly correlations indicate that the spreading rate on the America-Antarctic Ridge near the Bouvet triple junction is about 9 mm/yr (half-rate). The ridge has the rough topography and great relief characteristic of slow-spreading ridges. The average relief of the median valleys ($2.55 \pm .33$ km) is even greater than that of the Mid-Atlantic Ridge ($1.36 \pm .47$ km, Shih, 1980) consistent with the slower spreading rate of the America-Antarctic Ridge. Dredging of the fracture zones recovered diabase, green-

stone, gabbro, and abundant peridotites (in addition to pillow basalt) -- rocks which are rare from fast-spreading ridges.

On the basis of the new bathymetric and magnetic data, we have revised the pole rotation for the Antarctic and South American plates. Depending on the actual instantaneous plate velocities for the American-Antarctic Ridge at the Bouvet triple junction, the newly determined Antarctic-South America pole of rotation is either 85°S 22°E, if the half-spreading rate on the American-Antarctic Ridge is 9.2mm/yr, or is 84.5°S 14°E if the rate is 9.0mm/yr.

Published in: Journal of Geophysical
Research.

Supported by: NSF Grants DPP80-19769
and DPP77-21208.

WHOI Contribution No. 5376.

HYPERBOLIC STACKING OF WIDE-ANGLE REFLECTIONS FROM DEEP-SEA SEDIMENTS

Kristin Rohr

Wide-angle reflections from deep sea sediments were analyzed by stacking along hyperbolae. The details in the sediment structure that are achieved here are greater than any that have previously been made with wide-angle reflections on the Bermuda Rise. A 40 in³ airgun source was used in the western North Atlantic; the reflections were recorded with six ocean bottom hydrophones. The arrivals are stacked and semblance calculated along hyperbolic trajectories in T-X space and these two functions are multiplied together. This analysis allows an objective comparison of a large data set and an interpretation of the sediments' velocity structure. Events in the semblance x stack function can be correlated with normal incidence reflection data and are identified with the well known reflectors A^c, A*, β and acoustic basement. A low velocity event is observed after the basement arrival and is identified as shear reflections from acoustic basement. Compressional to shear conversion occurs at β , implying a Poisson's ratio of 0.40 for sediments beneath β . The stacking velocity and two-way travel time values are easily picked from the semblance x stack functions and can be interpreted for a velocity-depth function by an iterative modeling scheme or inversion. The results for the four layers agree closely. The upper sediments are 1.72 km/s and 0.27 km thick, sediments below A^c are 2.00 km/s and 0.2 km thick, below A*, 2.52 km/s and 0.25 km thick and below β , 3.1 km/s and 0.13 km thick. These velocities agree with the velocimeter measurements made nearby at

DSDP Site 387 and interpretations of interval velocity taken on thicker sections of these units on older crust.

In press: Journal of Geophysical Research.

Supported by: NSF Grant OCE79-09464.

WHOI Contribution No. 5230.

THE APPLICATION OF TAU-P MAPPING TO LOW SIGNAL TO NOISE RATIO WIDE-ANGLE REFLECTION DATA

K. Rohr

A tau-p analysis (Stoffa et al., 1980) was performed on wide-angle reflection data from upper oceanic crust in order to measure the initial velocity and velocity gradient in the upper basaltic layer (Layer 2). The large uncertainties in tau caused by a low signal to noise ratio and the topography of the basement reflector combined with the properties of tau-sum inversion resulted in a poorly constrained velocity-depth curve. Stacking the same data along hyperbolae gives a more detailed picture of the sedimentary velocity structure.

Supported by: NSF Grant OCE79-9464.

WHOI Contribution No. 5290.

VARIATION OF THE AMPLITUDES OF REFLECTORS FROM OCEANIC BASEMENT

Kristin Rohr

A study of the amplitudes of wide-angle reflections from the sediment-basement interface shows that the uppermost portion of Layer 2 is homogeneous on the 0.15-1.00 km scale, but not on the 3.0-8.0 km scale. The amplitude vs range patterns of the basement reflections recorded on 30 seismic lines were examined. Most patterns are affected by the rough topography of the sediment-basement interface. Three lines have smooth travel-time curves and appear to be unaffected by topography. The amplitude patterns of these lines are different from each other; they can be modelled by synthetic seismograms computed from velocity structures which have different thicknesses of a transition zone between the limestone sediments and the basaltic crust.

Supported by: NSF Grant OCE79-09464.

WHOI Contribution No. 5337.

DECONVOLUTION OF SEAFLOOR TOPOGRAPHY

Hans Schouten and Charles R. Denham

We develop a stochastic model of topography of normal sea floor formed by listric normal faulting of a thin brittle lithosphere that overlies the spreading center. We assume a Poisson density λ of fault locations and a Poisson density μ of horizontal extensions of the lithosphere (thickness T) along exponential faults with a decay parameter α . It is shown that this stochastic model is a second-order process. Deconvolution of a FAMOUS topography sample of normal sea floor (near 37°N, 33°W) is achieved by convolution of this sample with a two-term filter $(1, -3^{-\alpha})$ resulting in an alternating renewal process with dimensionless amplitude equal to αT . The renewal process displays the fault scarps that face inward toward the spreading center as well as the horizontal extensions along the faults in the sample topography. Reconstitution of the topography by convolution of a stylized version of the renewal process with the exponential fault solution (the inverse of the filter $(1, -3^{-\alpha})$) establishes an empirical relationship $\alpha T = 0.65$ and shows that a system of only inward-facing normal faults is a perfectly adequate model for this FAMOUS topography sample of normal sea floor. The listric normal faults reached a subhorizontal attitude at an estimated depth of approximately 1 km beneath the inner rift valley floor.

In press: Geophys. J.R. astr. Soc.

Supported by: ONR Contract N00014-82-C-0019.

WHOI Contribution No. 5391.

ESTIMATING THE ROUGHNESS OF SEAFLOOR

Hans Schouten and Charles R. Denham

The topography of normal seafloor is modeled as a stochastic process involving responses to listric-normal faulting in the young oceanic lithosphere. Fundamental wavelet decompositions, based on the prediction-error filter of high-resolution deep-tow data in the FAMOUS area, indicate that the basement topography there is statistically uniform. Ninety-eight percent of the topography variance can be represented by just three statistical parameters. The estimated response function accounts for both the steep fault-scarps and the decaying backward tilts that dominate the morphology of normal sea floor in the FAMOUS area. The response function characterizes the local thinning of the brittle, young oceanic crust that results from extension along listric

normal faults with exponential curvature. For FAMOUS topography, the faults are estimated to reach a sub-horizontal attitude at 1.0 ± 0.5 km depth.

Supported by: ONR Contract N00014-79-C-0071 and N00014-82-C-0019.

WHOI Contribution No. 5392.

FINITE DIFFERENCE SEISMOGRAMS FOR LATERALLY VARYING MARINE MODELS

Ralph A. Stephen

The method of finite differences is applied to the elastic wave equation to generate synthetic seismograms for laterally varying seafloor structures. The results are compared with borehole seismic data from the Gulf of California (DSDP Site 485) in which lines were shot over flat and rough topography. The significant new phenomena observed in both the synthetic seismograms and the field data is the generation of a 'double head wave' due to the interaction of the incident wavefront with the side of a hill and the flat sea floor adjacent to the hill.

In these models, the hills are on the order of a seismic wavelength in height and steep velocity gradients occur over distances comparable to wavelengths. Ray theoretical methods would not be suitable for studying such structures. True amplitude record sections are obtained by the finite difference method, which show for these models that the head wave generated at the flat sea floor adjacent to the hill is lower in amplitude than if the hill were not present and is lower in amplitude than the head wave generated at the hill.

A second feature which is important for borehole receivers is the existence of the 'direct wave root' in the upper basement. This energy occurs below the sharp interface when the direct wave impinges on the interface from above. There is no corresponding Snell's Law ray path for this energy and the energy is evanescent with depth in the lower medium.

The properties of both the double head wave and the direct wave root are clearly demonstrated in the finite difference 'snapshot' displays.

In press: Geophys. J.R. astr. Soc.

Supported by: ONR Contract N00014-79-C-0071.

WHOI Contribution No. 5551.

SYNTHETIC VERTICAL SEISMIC PROFILES BY THE METHOD OF FINITE DIFFERENCES

Ralph A. Stephen

Vertical seismic profiling is a particularly valuable tool in the investigation of lateral velocity and density variations around a borehole, as for example in the monitoring of enhanced oil recovery projects. In this application, both variable offset and variable azimuth shooting programs are of interest. As with any new experimental procedure, some theoretical analysis is warranted in order to best understand the acquired data. In the large offset VSP case, the interesting phenomena are wide angles of incidence, lateral variability, and seismic structure on the same scale as seismic wavelengths.

Unfortunately, all analytical attempts to simplify the elastic wave equation make limiting assumptions regarding one or more of these phenomena. Only finite element and finite difference methods are applicable for all three cases. This paper will discuss the finite difference method and give some examples of synthetic vertical seismic profiles for offshore wells.

In press: Vertical Seismic Profiling
Review Volume (Society of Exploration Geophysicists).

Supported by: ONR Contract N00014-79-C-0071.

WHOI Contribution No. 5487.

BOREHOLE SEISMIC EXPERIMENTS AND THE STRUCTURE OF UPPER OCEANIC CRUST

R.A. Stephen

This paper reviews the results of three borehole seismic experiments in oceanic crust and their implications for oceanic crustal structure. The experiments were carried out in the Western Atlantic (crustal age ~110 my), the Gulf of California (crustal age ~1 my) and the Costa Rica Rift Area (crustal age ~6 my).

The advantages of borehole data for resolving upper crustal velocity structure are summarized. These include the ability to obtain vertical velocity and attenuation information directly, improved resolution of upper crustal velocities, reduced sensitivity to effects of basement topography, and improved location and coupling.

Synthetic seismogram analysis by the reflectivity method resolves structure within the velocity-depth bounds determined from travel-times. Velocity gradients satisfy the amplitude criteria better than thick iso-velocity layers.

Particle motion analysis of three component data obtained in the Western Atlantic Ocean gives the first direct observation of seismic anisotropy in upper oceanic crust (Layer 2A).

All three experiments show relatively high upper crustal velocities (>4.0 km/sec) suggesting that layer 2A is not present even in the very young crust. All sites had over 100 m of sediment thickness, and it is postulated that sediment thickness and permeability, not merely age, governs the velocity of the upper crust by accelerating the cementation of fractures and cracks.

In press: Vertical Seismic Profiling Review Volume (Society of Exploration Geophysicists).

Supported by: NSF Grants OCE79-09351, OCE78-19801 and OCE81-17571.

WHOI Contribution No. 5193.

TRAVEL-TIME ANALYSIS OF BOREHOLE SEISMIC DATA

R.A. Stephen and A.J. Harding

A method is presented for reducing travel-time data from borehole seismic experiments to velocity-depth structure. The technique, which treats simultaneously data from any number of depths in the borehole in addition to surface data is based on the linear inversion scheme of Dorman and Jacobson (1981). Given the parameters $(p) = T(p) + pX(p)$ and $(p) = T(p) - pX(p)$ the method solves for depth to specified slownesses, assuming linear gradients (in velocity) between slowness values.

A practical limitation on the use of surface travel-time measurements to resolve velocity-depth structure is the necessity of assuming a surface velocity. This is not necessary for the case of borehole data. For a borehole receiver the velocity at the depth of the receiver can be obtained from the slope of the inflection point of the travel-time curve. Thus a direct measure of the uppermost velocity of a section can be obtained from the travel-time data of a shallow borehole receiver. Estimation of velocities from the inflection points of deeper receivers improves the resolution of the velocity-depth function which would be obtained from surface data alone.

The technique is applied to data from three borehole seismic experiments in oceanic crust. The experiments were carried out in the Western Atlantic (crustal age ~ 110 my), the Gulf of California (crustal age ~ 1 my) and the Costa Rica Rift Area (crustal age ~ 6 my). All three experiments show relatively high upper crustal velocities (>4.0 km/sec) suggesting that layer 2A is

not present even in the very young crust. All sites had over 100m of sediment thickness and it is postulated that sediment thickness and sediment permeability, not merely age, governs the velocity of the upper crust by accelerating the cementation of fractures and cracks.

Published in: Journal of Geophysical Research, 88, 8289-8298.

Supported by: NSF Grant OCE81-17571.

WHOI Contribution No. 5332.

CRUSTAL STRUCTURE IN THE OROZCO TRANSFORM ZONE

A.M. Trehu and G.M. Purdy

Seismic refraction data from lines of small shots and from individual large shots recorded during project ROSE are used to determine the crust and upper mantle velocity structure in the Orozco transform zone. The objective of this study is to provide constraints on the origin of a deep north-south trending trough flanked by two shallow ridges. These topographic features are incompatible with the present plate tectonic framework as defined by the regional spreading direction (approximately east-west) and did not display any seismic activity during project ROSE (Trehu and Solomon, 1982). Travel times and amplitudes from a reversed profile along one of the ridges reveal an anomalous crustal structure: a gradient of 0.5 sec^{-1} extending through most of the crust and decreasing to 0.1 sec^{-1} at the base, and a total crustal thickness of about 5.4 km. In contrast, three profiles from topographically normal oceanic crust indicate a crustal structure typical of oceanic crust throughout the oceans: a 1.5 km thick layer in which velocity increases from about 3 km/sec to 6.7 km/sec overlying a 4.5 km thick layer with a gradient of about 0.1 sec^{-1} . The upper mantle velocity beneath both structures, determined from travel times from large shots at ranges of 30 to 100 km, is about 8.1 km/sec. These results are interpreted to indicate that the trough and flanking ridges represent an extinct spreading center that developed an axial valley as the spreading center progressively shifted to the present axis of the East Pacific Rise.

In press: Journal of Geophysical Research.

Supported by: ONR Contract N00014-82-C-0019.

WHOI Contribution No. 5363.

CRUSTAL VELOCITY STRUCTURE ON THE FLANKS
OF THE MID-ATLANTIC RIDGE AT 24°N

Robert S. White and G. M. Purdy

A detailed seismic refraction survey was made over a 70 x 35 km area of young (9Ma) crust at 24°N 45°W to investigate the detailed velocity structure of a portion of 'typical' oceanic crust. An array of eight ocean bottom hydrophones was used to monitor 358 explosive charges and over 750 airgun shots. The crust exhibits normal oceanic structure, with a steep vertical velocity gradient through layer 2 and almost 3 km of layer 3 material with a mean velocity of 6.65 km/s overlying mantle. Ray tracing shows that the major amplitude variations observed in the upper crustal returns are governed more strongly by the seafloor topography than by the seismic velocity structure within the crust. A delay time function method was used to invert the 1613 well determined first arrival travel times from the explosives. We find that the delay time varies by only 0.05 to 0.10 s over most of the survey area, but that there are rather larger delay times in the north-west corner which may be associated with crust formed in the vicinity of a ridge jump. We also find a systematic correlation between variations in delay time and in the seafloor depth at the ray entry point, with smallest delay times where the seafloor is deepest. This is consistent with a model of the crust in which the topography of the base of layer 2 and the Moho is a subdued representation of the upper basement surface.

Published in: Geophys. J. Roy astr.
Soc, 75, 361-385, 1983.

Supported by: ONR Contract N00014-74-
00262 and N00014-82-C-0019.

WHOI Contribution No. 5406.

M A R I N E C H E M I S T R Y

REDUCED ADVECTION INTO ATLANTIC OCEAN
DEEP EASTERN BASINS DURING LAST
GLACIATION MAXIMUM

Bill Curry and G. P. Lohmann

Causes of change in deep water $\delta^{13}\text{C}$ can be either global or local in extent. Global causes include (1) climatically-induced changes in the amount of terrestrial biomass which alter the average carbon isotopic composition of the oceanic reservoir, and (2) erosion and deposition of organic-rich, continental shelf sediments during sea level fluctuations which change the mean organic

carbon:phosphorus ratio. Regional gradients of $\delta^{13}\text{C}$ are created by remineralization of organic detritus within the deep ocean itself thus reflecting the distribution of water masses and modern thermohaline flow. Changes in a single geological record of benthic foraminiferal $\delta^{13}\text{C}$ can result from any combination of these global and abyssal circulation effects. By sampling a large number of cores collected over a wide bathymetric range yet confined to a small geographical region, we have minimized the ambiguity. We can assume that each $\delta^{13}\text{C}$ record was equally affected by global causes of $\delta^{13}\text{C}$ variation. The differences seen between the $\delta^{13}\text{C}$ records must, therefore, reflect changes in the distribution of $\delta^{13}\text{C}$ in the deep ocean. We interpret these differences in distribution in terms of changes in the ocean's abyssal circulation. Benthic foraminiferal carbon isotopic evidence from a suite of Sierra Leone Rise cores indicates that the deeper parts of the eastern Atlantic basins underwent a reduction in $[\text{O}_2]$ during the maximum of the last glaciation. Reduced advection of O_2 -rich deep water through low-latitude fracture zones, associated with increased delivery of organic matter to the deep ocean, lowered the $\delta^{13}\text{C}$ of deep water CO_2 at all depths below the sill separating the eastern and western Atlantic basins. This decreased advection into the eastern Atlantic Ocean coincides with the overall decrease in deep water production in the North Atlantic during the last glacial maximum.

In press: Nature, 306, 577-580.

Supported by: NSF Grants OCE 78-19768,
OCE81-10935 and OCE82-00151.

WHOI Contribution No. 5424.

P L A T E T E C T O N I C S

OVERLAPPING SPREADING CENTERS
ON EAST PACIFIC RISE

Hans Schouten and Kim Klitgord

A correlation between overlapping spreading centers (OSE, Macdonald and Fox, 1983) on the East Pacific Rise and the pattern of fracture zones predicted by Klitgord and Mammerickx (1982) from bathymetry and magnetic anomalies, suggests that OSC's represent zero-offset transform zones (Schouten and White, 1980; Schouten and Klitgord, 1982) that have been stable features on the East Pacific accreting boundary for at least 5 Myr.

In press: Nature, 303, 549-550, 1983.

Supported by: ONR Contract N00014-74-C-0262 and NSF Grant OCE76-22512 A01.

WHOI Contribution No. 5312.

P A L E O N T O L O G Y

CENOZOIC GEOCHRONOLOGY

William A. Berggren, Dennis V. Kent,
John J. Flynn and John A. Van Courvering

We present a revised Cenozoic geochronology based upon a best fit to selected high temperature radiometric dates on a number of identified magnetic polarity chrons (within the late Cretaceous, Paleogene, and Neogene) which minimizes apparent accelerations in sea floor spreading. An assessment of over 200 first order correlations of calcareous plankton biostratigraphic datum events to magnetic polarity stratigraphy yields an improved correlation of the standard magnetostratigraphic, standard biostratigraphic (zonal) and chronostratigraphic boundaries as well as improved resolution in marine-continental stratigraphic correlations. The time scale presented here has been accepted by the Committee on Geochronology as the standard time scale for the Cenozoic for the Decade of North American Geology (DNAG).

In press: Geological Society of America Bulletin.

Supported by: NSF Grant OCE80-19052 and the Senior Studies Award.

WHOI Contribution No. 5478.

LATE TERTIARY/QUATERNARY MAGNETOSTRATIGRAPHY AND BIOSTRATIGRAPHY OF THE VEMA CHANNEL SEDIMENTS

David A. Johnson, Michael T. Ledbetter,
Eric Tappa and Robert Thunell

Twenty-one piston cores in a transect of the Vema Channel near 31°S were analyzed magnetostratigraphically and biostratigraphically in order to ascertain the depositional continuity in the late Tertiary and Quaternary. The cores span a depth interval of 2941 to 4235 meters and represent three distinct physiographic provinces, each of which shows a relatively consistent accumulation pattern. The deepest cores are from the broad terraces which flank the Vema Channel axis on both

the west (~3650 m) and east 4150-4250 m); these cores show the highest accumulation rates in the region (>20 m/m.y.) and no evidence of unconformities. Twelve cores from the flanks of the Rio Grande Rise (2941-3950 m) show depositional continuity in the late Quaternary and normal pelagic accumulation rates (5-10 m/m.y.), with evidence in two of the cores for minor erosional hiatuses on a local scale, perhaps due to slumping. A relatively narrow transition zone between the Rio Grande Rise flanks and the eastern terrace (3950-4150 m) shows buried manganese pavements and near-outcrops of Miocene/Pliocene strata, indicating intermittent erosion during the late Tertiary and Quaternary, perhaps due to episodes of intensified AABW flow. Modern flow within this transition zone and over the eastern terrace is relatively weak ($v \approx 2$ cm/sec) and southward. The presence of well-developed sediment waves on the terrace and buried erosional unconformities upslope to the east suggests substantially higher current velocities during the late Tertiary than those present today.

In press: Marine Geology.

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5477.

HIGH-RESOLUTION STRATIGRAPHY IN LATE PLEISTOCENE-HOLOCENE SEDIMENTS OF THE VEMA CHANNEL

Glenn A. Jones, David A. Johnson
and William B. Curry

Twenty-five giant gravity cores (GGC) were obtained from the southwest flank of the Rio Grande Rise in the summer of 1980 and were analyzed for time-slice interpretations of the late Pleistocene paleo-oceanography of the southwestern Atlantic. The cores span a depth interval of 1430 to 4148 meters, encompassing each of the principal deep-water masses. The suite of cores contains a continuous depositional record of the past 150,000 to 200,000 years.

From calcium carbonate, oxygen isotope, carbon-14, and biostratigraphic analyses, it is possible to determine the precision in core-to-core correlation which is obtainable within this region of relatively low accumulation rates (0.5 to 2.0 cm 10^{-3} years). Calcium carbonate analyses were performed on each of the 25 cores at 1-cm sampling intervals, using a technique that yields an analytical precision of $\pm 0.25\%$. The carbonate curves show glacial/interglacial amplitudes of typically 10% to 30% CaCO_3 . Higher-frequency amplitudes of 2% to 5% are also present. Both the glacial/interglacial and the higher-frequency car-

bonate signals can be correlated from core to core over the entire depth interval of the transect and are essentially synchronous based on independent stratigraphic control provided in selected cores by $\delta^{18}\text{O}$, ^{14}C , and biostratigraphic analyses. Despite the low sedimentation rates within this particular region, we obtain a precision in core-to-core correlation of ± 1000 to 2000 years over the late Pleistocene/Holocene ($< 30,000$ yBP) and ± 4000 to 10,000 years over the late Pleistocene from 30,000 yBP to 150,000 yBP. Comparison of the carbonate stratigraphy with the planktonic foraminiferal $\delta^{18}\text{O}$ record shows that there is an apparent lag of approximately 4000 years between the $\delta^{18}\text{O}$ and CaCO_3 curves at Terminations I and II but no lag at full glacial or interglacial time.

These high-resolution studies from the Vema Channel have important implications for interpreting the effects of bioturbation on a given stratigraphic sequence. Modeling the effects of bioturbation on the abundance and isotopic records of the benthonic foraminifera *Planulina wuellerstorfi* suggests that, for the region of the Vema Channel/Rio Grande Rise, invalid stratigraphic and paleoceanographic interpretations may result if abundance variations in this species are ignored.

The results of these analyses show that it is possible to generate detailed stratigraphies in regions that have not previously been selected for high-resolution studies solely because of relatively low accumulation rates of sediments. Our results indicate that carbonate analyses in some regions may provide a distinct advantage over $\delta^{18}\text{O}$ analyses because of the ease of analysis and high precision for identifying low-amplitude, high-frequency fluctuations which can be correlated from core to core. At the same time, independent stratigraphic control (via $\delta^{18}\text{O}$ or ^{14}C) is essential to verify whether this technique of carbonate curve-matching is valid.

In press: Marine Geology.

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5511.

EVIDENCE OF PUNCTUATED GRADUALISM
IN THE *Globorotalia tumida*
LINEAGE (PLANKTONIC FORAMINIFERA)

Bjorn A. Malmgren, W.A. Berggren
and G.P. Lohmann

The warm-water planktonic foraminiferal *Globorotalia tumida* lineage has been studied in a 10 Myr long sequence (Late Miocene through Recent) from Deep Sea Drilling Project Site 214 in the southern Indian Ocean

to determine long-term evolutionary patterns through the lineage's history and the tempo and mode of the evolutionary transition from *G. plesiotumida* to *G. tumida* across the Miocene-Pliocene boundary. Sampling resolution was about $1.5 \cdot 10^4$ yrs across this boundary and about $2 \cdot 10^5$ yrs otherwise. The shape was analyzed using Eigenshape analysis. This method permits complete descriptions of shape by a single shape function. The *G. plesiotumida* - *G. tumida* transition involved a change in ventral-dorsal height proportions of the test and a size increase. The complete modification of shape took about 0.6 Myr (or $3 \cdot 10^6$ to $6 \cdot 10^6$ generations) and was not rapid as predicted by the punctuated equilibrium model. The modification was not unidirectional, but accomplished by shift in a general direction interrupted by several reversals. The Late Miocene and Pliocene-Pleistocene records show minor fluctuations in shape that did not lead to major phyletic changes. The pattern of shape variation through the sequence does not conform to either of the punctualism or gradualism models. Size began to increase in the latest Miocene, following considerable change in the Late Miocene, increased in a step-like fashion at the Miocene-Pliocene boundary, and showed a zig-zag pattern of change through the remainder of the sequence. This record is, therefore, not an example of punctualism despite the stepwise change at the Miocene-Pliocene boundary. Migration probably did not have any major influence on the observed patterns. If it had, we would have expected obvious relationships to known paleoclimatic oscillations, which could have reflected climatically induced migrations of distinct morphotypes from different climatic regimes across the site.

In press: Paleobiology.

Supported by: NSF Grant OCE80-19052.

WHOI Contribution No. 5335.

RADIOLARIA

Kozo Takahashi

Radiolarians settling through the oceanic water column were recovered from three stations (western Tropical Atlantic-Station E, central Tropical Pacific- P_1 and Panama Basin-PB) using PARFLUX sediment traps in moored arrays at several depths. The taxonomic diversity of the radiolarian assemblages in the sediment traps was very high. A total of 420 taxa (including 23 new taxa) were found at the three stations; of these, 208 taxa were found at station E. The polycystine radiolarians generally reach the sea floor with little change in abun-

dance or species composition, although slight skeletal dissolution occurs throughout their descent. The phaeodarian radiolarians, on the other hand, are largely dissolved within the water column; only a few species reach the sea-floor and these dissolve rapidly at the sediment water interface. Most radiolarian skeletons sink as individuals through deep water columns without being incorporated into large biogenic aggregates. Because significant numbers of nassellarian and phaeodarian species are deep-water dwelling forms, the diversity index of radiolarians increases with increasing depth in the mesopelagic zone.

The vertical flux of the total radiolarians arriving at the trap depths (in $\times 10^3$ individuals/m²/day ranged from 16-24 (E), 0.6-17 (P₁), and 29-53 (PB)). Of these, on the average, 25 percent and 69 percent of the total radiolarian flux is transported by Spumellaria and Nassellaria, respectively, while 5 percent is carried by Phaeodaria. The microscopic count, weight, and SiO₂ content data are combined in order to estimate SiO₂ supply by radiolarians. The supply of radiolarian silica (mg/m²/day) to each trap depth ranged from 2.5-4.0 (E), 0.9-3.2 (P₁), and 5.7-10.4 (PB). The Radiolaria appear to be a significantly large portion of the SiO₂ flux in 63 μ m size fraction and thus play an important role in the silica cycle. When the radiolarian fluxes at the three stations are compared with Holocene radiolarian accumulation rates in the same areas, it became apparent that several percent or less of the fluxes are preserved in the sediments in all cases and the rest must be dissolved on the sea floor.

Supported by: NSF Grant OCE 80-19386.

WHOI Contribution No. 5156.

P E T R O L O G Y

SYSTEMATICS OF MODAL PHENOCRUST ASSEMBLAGES IN SUBMARINE BASALTS: PETROLOGIC IMPLICATIONS

W.B. Bryan

Phenocryst assemblages in ocean-ridge basalts generally show an increasing proportion of plagioclase as the total amount of phenocrysts increases. The variations in phase assemblages, as well as most crystal liquid K_d's, are similar to variations (equimodal trends) predicted by low-pressure laboratory experiments, suggesting that many of these basalts have experienced varying degrees of low-pressure crystallization prior to quenching, with little sorting of

crystals and liquid. Important exceptions include moderately to highly phyric basalts enriched either in plagioclase or olivine which lie well off the experimental trends. In these basalts, megacrysts and xenocrysts usually cited as evidence for magma mixing commonly represent a small proportion of the total crystalline phase assemblage. However, phase proportions for many of these basalts lie well outside the range that could be produced by simple mixing; selective gravitative sorting either prior or subsequent to mixing appears to be the likely explanation for these phyric basalts. A relation between spreading rate and phase proportions is neither supported nor refuted by the data, which as yet do not adequately represent fast-spreading ridges.

Pyroxene-phyric varieties are especially common among LIL-element enriched (Group 2) basalts, and these basalts also show the greatest abundance of olivine-enriched (picritic) samples. Selective enrichment in plagioclase is more common among LIL-element depleted (Group 1) basalts, and pyroxene appears in Group 1 basalts only at relatively high degrees of crystallinity. These differences are consistent with expected compositional effects (including volatiles) on phase boundaries, as well as likely differences in depth (pressure) of mantle melting and magma fractionation.

Sparsely to moderately phyric basalts tend to contain only olivine ($\pm$ spinel) as phenocrysts and lie in the olivine field in the CMAS projection, consistent with the concept that these magmas approach low-pressure equilibrium by olivine fractionation from a picritic parent. The origin and relationships between these basalts remains an important fundamental problem.

Phenocryst phase assemblages provide a reliable guide to the low-pressure phase saturation of the associated liquid (glass) phase, as indicated by the projected positions of these glasses in CMAS. It is suggested that, in contrast to the classical practice of classifying basalts according to phase proportions, a classification based on first appearance of a crystalline phase would be petrogenetically more informative.

Published in: Cont. Mineral. Petrol.
83, 62-74, 1983.

Supported by: NSF Grant OCE81-10932.

WHOI Contribution No. 5342.

RECONSTRUCTION OF THE GEOMETRY OF
ACCRETION DURING FORMATION OF THE
BAY OF ISLANDS OPHIOLITE COMPLEX

J.F. Casey, J.A. Karson, D. Elthon,
Elric Rosencrantz and Marchall Titus

Two types of highly allochthonous ophiolitic terrains crop-out adjacent to one another in the Bay of Islands region of southwestern Newfoundland, Canada. The first, called the Coastal Complex (CC), is exposed as a long linear belt along the west coast. It consists of a highly deformed ophiolitic assemblage interpreted as oceanic crust and upper mantle with a previous transform fault tectonic history. The second, called the Bay of Islands Complex (BOIC), crops-out to the east and consists of a series of four relatively undeformed ophiolitic massifs interpreted as oceanic crust and upper mantle with a more normal seafloor spreading tectonic history. An autochthonous relationship can be demonstrated between the two types of ophiolitic terrains in the southernmost of these massifs indicating that each evolved along a single ridge-transform system. The internal structure of both the BOIC and the CC is consistent with the above interpretations. The NE-SW structural trends in the CC transform are oriented approximately parallel to the seafloor spreading direction inferred from the average orientation of the sheeted diabase dykes within the BOIC. One-way chilling statistics from the sheeted diabase dyke unit, growth directions of dendritic crystals along inclined comb layers in the plutonic section, and the sense of shear along the CC transform all indicate that the BOIC was accreted on the NE side of a NW-SE trending spreading center (present geographic reference frame). Juxtaposition of the BOIC with the NE-SW trending CC occurred at a ridge-transform intersection. The transform exhibited a left-lateral offset. Magma chamber profiles have been identified in the layered gabbroic sections of the BOIC using the patterns outlined by the attitudes of fine-scale igneous layering. These patterns depict arcuate structures which appear to roughly define a cross-sectional view of a systematically oriented half-cylinder the long axis of which is oriented parallel with the plane of the sheeted dykes and major lithologic unit contacts. Thus, we believe that a large cylindrical shaped crustal-level subaxial magma chamber extended in a direction parallel with the ridge axial plane at the time of formation of the BOIC. The CC marks the abrupt truncation of this long linear magma chamber. Petrologic constraints demonstrate that the ultramafic and transition zone cumulates, which form the base of the plutonic section, crystallized at a range of high to intermediate pres-

ures possibly along a long narrow conduit extending to >30km depth beneath the crustal level magma chamber. After formation, these cumulates were deformed at high temperatures in the zone of mantle upwelling and divergence, then tectonically transported to crustal or near crustal levels to form the basement upon which the remainder of the ophiolite stratigraphy was accreted. A geometric and tectonic model is presented for the accretion process responsible for formation of the Bay of Islands Ophiolite.

Published in: Tectonics, 2, 509-528.

Supported by: NSF Grant EAR80-26445.

WHOI Contribution No. 5382.

MINERALOGIC STUDIES OF THE RESIDUES
OF MANTLE MELTING: ABYSSAL AND
ALPINE-TYPE PERIDOTITES

H.J.B. Dick and R.L. Fisher

Abyssal peridotites dredged from the ocean ridges range from diopside-poor harzburgite to lherzolite, but all contain enstatite saturated with diopside, indicating that melting of the abyssal mantle was constrained by the pseudo-invariant point $Ol+En+Di+Sp+Melt$. We find systematic regional differences, which suggest that a range of primary melt compositions and large variations in the apparent degree of melting exist in the mantle beneath ocean ridges. We note in particular that North Atlantic peridotites are highly depleted relative to the average abyssal peridotite.

Alpine-type peridotites overlap the range for abyssal peridotites, but extend to far more depleted and enriched compositions. Many contain enstatite undersaturated with respect to diopside. Frequently, alpine-type peridotites contain highly magnesian Al-poor and Cr-rich minerals lying outside the abyssal range. Melting of many alpine peridotites, therefore, has occurred well into the three phase field $Ol+En+Sp+Melt$ under different conditions than for abyssal peridotites. Alpine-type peridotites evidently represent parageneses extending from relatively undepleted sub-continental upper mantle to peridotites melted in a volcanic-arc or near-arc oceanic environment.

Supported by: NSF Grants DPP77-21208
and DPP80-19769.

WHOI Contribution No. 5466.

FRACTIONATION OF PYROXENE-PHYRIC MORB AT
LOW PRESSURE: AN EXPERIMENTAL STUDY

T.L. Grove and W.B. Bryan

One-atmosphere melting experiments are used to assess the role of clinopyroxene in producing the compositional variations observed in mid-ocean-ridge basalts (MORBs) from the North Atlantic. Analog models of natural glasses and associated phenocrysts show that several possible parental magmas may undergo low pressure fractional crystallization involving olivine and spinel, followed by plagioclase, and then by augite. The phenocryst phase assemblages in natural deep-sea basalts are closely correlated with the major element compositions of their associated quenched glasses, and the projections of these glasses on the Oliv-Cpx-Qtz pseudoternary correspond to the 1-atmosphere phase boundaries and reaction points defined by laboratory experiments. Comparison of natural phenocrysts with experimental phases indicates that the augites preserved in moderately fractionated MORB from the FAMOUS area may have formed at or near the ocean floor and need not be relics of high pressure processes.

In press: Contributions to Mineralogy and Petrology.

Supported by: NSF Grant OCE81-10932.

WHOI Contribution No. 5491.

VARIATIONS IN STRUCTURE AND PETROLOGY IN
THE COASTAL COMPLEX, NEWFOUNDLAND:
THE ANATOMY OF AN OCEANIC FRACTURE ZONE

J.A. Karson

The Coastal Complex of western Newfoundland is perhaps the world's best example of an ancient oceanic fracture zone preserved within an ophiolite complex. The structural style, metamorphic zonation, and magmatic history of the Coastal Complex contrasts sharply with that of the contiguous, stratiform Bay of Islands Complex. The various exposures of the Coastal Complex provide cross sections of the fracture zone at various vertical and lateral positions. A reconstruction of these sections reveals structural and petrological variations that are a function of depth as well as position along strike. For example, deep crustal sections show a history of strike-slip deformation that is fairly homogeneously distributed across a 4 km wide high-strain zone. High-temperature metamorphic rocks, partial melting relationships, and peridotite intrusive bodies are also present. Diabase dykes cut the entire assemblage. At shallow crustal levels, deformation is concentrated along

relatively narrow, well-defined faults and shear zones that overall have an anastomosing aspect. High-grade metamorphic rocks occur only locally and have often suffered retrograde effects. Felsic intrusive bodies occur locally and serpentinite diapirism is widespread. Volcanic rocks unconformably overlie a brecciated horizon that transects the entire assemblage. The most striking lateral variations are the discontinuous nature of lithologic units and the apparent bifurcation of major high-strain zones along strike.

Oceanic fracture zones and their ophiolite analogs probably display a wide range of geological relationships that vary with depth, with position along strike, and in scale. Thus, the Coastal Complex should probably be viewed only as a single example of how exceedingly complex and variable the oceanic lithosphere may be near transform faults and their nontransform extensions. Clearly, similar ophiolitic assemblages should not be considered when attempting to model processes along purely extensional plate boundaries.

In press: Geological Society of London.
Special Publication on Ophiolites.

Supported by: NSF Grant EAR80-26445.

WHOI Contribution No. 5350.

GEOCHEMISTRY, MINERALOGY AND PETROGENESIS
OF LAVAS ERUPTED ALONG THE SOUTHWEST
INDIAN RIDGE BETWEEN THE BOUVET TRIPLE
JUNCTION AND 11 DEGREES EAST

A.P. leRoex, H.J.B. Dick, A.J. Erlank,
F.A. Frey and S.R. Hart

Basaltic lavas from fracture zones and ridge segments have been systematically sampled along the Southwest Indian Ridge from the Bouvet triple junction to 11°E. Detailed studies of the major and trace element and isotopic variations in these lavas serve to distinguish three compositional groupings which occur throughout this region. The one type comprises basalt with typical 'depleted' (N-type MORB) characteristics with high Zr/Nb (>17), Y/Nb (>8) and $^{143}\text{Nd}/^{144}\text{Nd}$ ($0.51312 \pm 2 - 0.51302 \pm 2$) ratios, LREE depletion ($\text{La}/\text{Yb}_N = 0.35-1.1$) and with relatively low $^{87}\text{Sr}/^{86}\text{Sr}$ ($0.70267 \pm 4 - 0.70297 \pm 4$) ratios. The second, least abundant, type has typical 'plume' characteristics (P-type MORB) with low Zr/Nb (5.8-6.8), Y/Nb (0.9-1.2) and $^{143}\text{Nd}/^{144}\text{Nd}$ ($0.51295 \pm 2 - 0.51286 \pm 2$) ratios, strong LREE enrichment ($\text{La}/\text{Yb}_N = 4.8-6.9$) and relatively high $^{87}\text{Sr}/^{86}\text{Sr}$ ($0.70356 \pm 4 - 0.70364 \pm 4$) ratios. These compositional features are similar to those found in the nearby Bouvet Island lavas. The third type has a transitional

chemistry (T-type MORB) with relatively low Zr/Nb (7.7-11.8), Y/Nb (1.3-3.0) and $^{143}\text{Nd}/^{144}\text{Nd}$ (0.51301 $\pm$ 2 - 0.51284 $\pm$ 2) ratios, slight LREE enrichment ($\text{La}/\text{Yb}_N = 1.7-4.3$) and with relatively high $^{87}\text{Sr}/^{86}\text{Sr}$ (0.70291 $\pm$ 4 - 0.70370 $\pm$ 4) ratios. All three lava types show similar variations in terms of major element composition (e.g., $\text{Mg}\alpha$).

N- and T-type lavas are volumetrically the most abundant, and lavas from both groups occur at most localities that have been sampled. P-type lavas have been found at approximately half the localities sampled and crossing REE patterns are a common phenomenon of these data sets. There is no evidence for a 'geochemical gradient' away from the present location of the Bouvet mantle 'plume'; rather, a juxtapositioning of all three lava types occurs along this entire length of the Southwest Indian Ridge system.

A comparison of lavas dredged from fracture zones and ridge segments shows no inherent geochemical differences which can be related to the source regions giving rise to these lavas. The fracture zone basalts do however tend to more evolved compositions, a feature attributed to the cooling effect of the adjacent, cold passive lithospheric plate.

Consideration of trace element and isotopic systematics supports the concept that the lavas have been derived from a mantle that is chemically heterogeneous on the scale of tens of kms. The available data are generally consistent with a model in which the lavas with P-type characteristics have been derived from an 'enriched' source region related to the upwelling associated with the Bouvet mantle 'plume' and similar to that giving rise to the Bouvet Island lavas, while the N-type lavas have been derived from normal 'depleted' sub-oceanic mantle. It is suggested that the T-type lavas represent mixing products derived from an N-type source region that has experienced less enrichment (by veining by small degree partial melts and/or metasomatic fluids) than that giving rise to the P-type magmas.

Supported by: NSF Grants DPP77-21208, DPP79-00036 and DPP 80-19769.

WHOI Contribution No. 5373.

SEDIMENTOLOGY

BEACH CHANGES ON COASTS WITH DIFFERENT WAVE CLIMATES

David G. Aubrey

Seasonal and longer-term beach variability is quantified for seven U.S. beaches exposed to widely varying wave climates. One U.S. west coast location (southern California) and six U.S. east coast locations (from North Carolina to Massachusetts) form the basis of this study. Wave exposure varies from complete exposure to open ocean waves, to partly sheltered locations, and finally to nearly complete sheltering where locally-generated waves dominate. Beach response was documented with beach profiles distributed along each of the seven coastal locations, spanning a minimum of five years of observation. Frequency of measurement was at least once per month, with periods of more intense weekly sampling lasting for up to two years (southern California location). Wave climate was either measured directly or estimated from hindcast and/or compilations of ship observations. Consequently, wave information varies in detail from joint statistics of wave height, frequency, and direction, to compilations of local storm history (and hence inferred wave behavior). Magnitude of annual beach variability ranged from 3.3 m³ per meter of beach to 0.2 m³ per meter of beach, with the greatest variability in regions exposed to open ocean waves (most severe wave climate) and the lowest variability along protected coasts (least severe wave climate). All open coast locations studied had a seasonal variability which accounted for at least 50 percent of the beach variability. Protected coastal locations had less pronounced seasonal signatures. These seasonal and aseasonal beach responses mirror corresponding seasonality (or lack thereof) in wave and storm climates. The study re-emphasizes the need for careful measurements of coastal wave climate to enable predictive modeling of shoreline behaviour. It also discusses different analysis techniques for analyzing changes in beach profiles through time.

Published in: Beach Changes on Coasts
With Different Wave Climates.
McLachlan and Erasmus, eds. D.W.
Junk Publishers, The Hague, 63-85.

Supported by: NOAA Sea Grant (NSTS
project) NA80-AA-D-0047.

WHOI Contribution No. 5324.

UPDRIFT MIGRATION OF TIDAL INLETS: A
NATURAL RESPONSE TO WAVE, TIDE AND
STORM FORCING

David G. Aubrey, Paul E. Speer
and Eric Ruder

Three mechanisms are responsible for tidal inlet migration in an updrift direction (counter to net longshore transport): 1) attachment of distal ebb tide delta bars to the downdrift barrier spit; 2) storm-induced breaching and subsequent stabilization to form a new inlet; and 3) ebb tide discharge around a sharp channel bend creating a three-dimensional flow pattern which erodes the outer channel bank and extends the inner channel bank. The last two mechanisms can result in either updrift or downdrift inlet migration, depending on channel geometry in the bay and barrier beach configuration. Analysis of historical charts and aerial photographs, combined with a historical storm synthesis, shows that all three mechanisms are active at a natural tidal inlet along a sandy coast (Nauset Inlet, Cape Cod, MA). On a time scale of ten years, these mechanisms were effective in producing an updrift migration of more than two kilometers.

Supported by: NOAA Sea Grant No.
NA80-AA-D-00077 (R/B-21) and U.S.
Army Research Office No.
DAAG29-81-K-0004.

WHOI Contribution No. 5205.

SEDIMENTATION OF PARTICULATE
MANGANESE IN THE DEEP OCEAN

Susumu Honjo, Jonathan J. Cole,
Alan P. Fleer and Cynthia H. Pilskaln

The vertical flux of particulate Mn was measured at four open ocean locations using PARFLUX sediment traps with 1.5 m² openings. Several traps were deployed for a few months at each station in mooring arrays along bottom-tethered taut lines in the mesopelagic, bathypelagic, and near bottom abyssal layers. The bottom depth of the trap stations ranged from 3,850 m to 5,792 m. Additionally, at the Panama Basin station, the seasonal variability in Mn flux was measured by PARFLUX time series sediment traps at 890 m, 2,590 m, and 3,560 m.

At all stations and at all times of the year the flux of particulate Mn increased with increasing depth, strongly suggesting a deep-water source for settling Mn. The flux of Mn in abyssal water ranged from very low values in East Hawaii (4 $\mu\text{g}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$ at 5,582 m) to extremely high values at the Panama Basin (up to 3,000 $\mu\text{g}\cdot\text{m}^{-2}\cdot\text{day}^{-1}$ at 3,560 m). At the Panama Basin, the flux

of Mn varied seasonally and highest rates appear to coincide with periods of high biological productivity in surface waters. The largest flux of Mn occurred when a coccolithophorid bloom rapidly sank to the bottom.

The particulate Mn collected by our sediment traps is neither of biogenic nor lithogenic origin. Most of the Mn (often above 90 percent in deep water) was in the form of "non-structural" Mn, absorbed or loosely bound to other particles such as large amorphous aggregates (LAA). It is possible that Mn was removed from solution by the carbonate in coccolithophorids which, in turn, were scavenged from the water column by LAA. Although fecal pellets contributed less than 10 percent to the Mn flux, the concentration of Mn in the pellets was higher than in the bulk trap material.

Our data indicate that Mn in surface waters is a minor contribution to Mn flux in deep water. Rather, most of the Mn is supplied by independent advective deep water processes. These data also indicate that Mn has a short residence time in the deep water column.

In press: Earth and Planetary
Science Letters.

Supported by: NSF Grant OCE82-08736.

WHOI Contribution No. 5552.

SEASONALITY IN THE FLUXES OF SUGARS, AMINO
ACIDS AND AMINO SUGARS TO DEEP OCEAN
PART II. PANAMA BASIN

Venugopalan Ittekkot, Egon T. Degens
and Susumu Honjo

Time series sediment traps were deployed for an entire year at depths of 890, 2590 and 3560 m at a station in the Panama Basin during 1980. Fluxes of sugars, amino acids and amino sugars varied seasonally at each depth. Two peak fluxes were observed: one in February-March, and the other in June-July. These peaks were associated with a high productivity period caused by regional upwelling and an unusual coccolithophorid bloom. There were significant differences in the distribution pattern of sugars and amino acids associated with these fluxes. The peak flux of June-July was characterized by high amounts of arabinose and ribose within the sugar and high amounts of aspartic acid in the amino acid fractions. These differences were recorded at all three depths simultaneously, indicating rapid vertical transport without significant dissolution or decomposition. The observed pattern indicates the utility of specific compounds like sugars and amino acids as tracers of particle flux in the marine environment.

In press: Deep-Sea Research.

Supported by: NSF Grant OCE81-25429.

WHOI Contribution No. 5441.

LATE CENOZOIC TURBIDITE AND CONTOURITE
DEPOSITION IN THE SOUTHERN BRAZIL BASIN

David A. Johnson and Kenneth A. Rasmussen

Piston and giant gravity cores, together with 3.5 kHz and seismic profiles, reveal a complex pattern of dispersal of bioclastic calcareous turbidites in the southern Brazil Basin. Extensive canyons, up to 500 meters deep, dissect the steep north flank of the Rio Grande Rise and serve as conduits for turbidity currents which have eroded modern sediments as well as outcrops of Tertiary and late Cretaceous strata. The resulting turbidites are up to several meters thick in channel axes and commonly several cm thick at distances up to hundreds of km from the canyon source region. The lithology of both proximal and distal turbidites, their presence well below the modern CCD, and the lack of correspondence between their presence and that of displaced Antarctic diatoms, verifies that these calcareous deposits are not products of episodic Antarctic Bottom Water flow, but originated from a shallow-water source area. Individual turbidity currents were sufficiently strong to transport fine-grained components several hundred meters upslope and several hundred km laterally across the Brazil Basin.

The late Pleistocene and Holocene witnessed major turbidity current episodes at intervals on the order of 50,000 years, with at least two episodes coinciding approximately with interglacials following Terminations I and II. Sediments older than isotopic stage 8 (~275,000 yBP) include thinly-laminated "contourites." We interpret these units as intervals of normal pelagic sedimentation which were subsequently reworked under the influence of AABW pulses. Episodes of intensified AABW flow may have immediately preceded the onset of major glacial episodes, extending corrosive eastward-flowing AABW to relatively shallow depths (4200-4300m) along the north flank of the Rio Grande Rise. The undercutting of outcropping strata by chemical dissolution at the base of the faulted north flank has probably played a critical role in the initiation of canyon-cutting. Episodic slumping due to oversteepening and the generation of debris flows progressed upslope in a dendritic pattern, gradually extending the canyons to shallower elevations on the Rise. Vertical crustal displacements associated with the Rio Grande Fracture Zone have controlled the pathway of AABW in the southern Brazil Basin, and accentuated the

steep relief on the north flank of the Rise. These two processes have been essential conditions for the extensive canyon cutting and subsequent dispersal of carbonate turbidites over the Brazil Basin.

In press: Marine Geology.

Supported by: NSF Grant OCE80-25208.

WHOI Contribution No. 5440.

ANALYSIS OF A LONGITUDINAL RIPPLE
FROM THE NOVA SCOTIAN CONTINENTAL RISE

I.N. McCave, C.D. Hollister, D.J.
DeMaster, C.A. Nittrouer, A.J. Silva
and J.Y. Yingst

A longitudinal ripple was obtained in a box core taken from the Nova Scotian Continental Rise in July 1982. Soft brown mud comprising 1.5 to 10 percent sand, ~60 percent silt and ~35 percent of <2 μ m clay forms the 5 cm high and ~40 cm wide ripple. A maximum thickness of 8 cm of this mud under the crest, thinning to 2 cm on the ripple's flank, overlies stiff muddy foraminiferal ooze. The vane shear strength of the brown mud is ~0.4 kPa whereas that of the ooze is ~4 kPa. X-radiographs show the mud to contain many fine burrows and a few larger ones as well as winnowed horizons rich in foraminifera of both benthonic and planktonic origin. Gentle wet sieving of the sand fraction on board ship showed the sand fraction to contain very few fecal pellets. Thus, it appears unlikely that bedload movement played a large part in formation of the ripple. Rapid initial deposition from a concentrated suspension is suggested to have formed the structure, but this was followed by periods of erosion to yield winnowed horizons and further rebuilding with material deposited from suspension. Radiochemical data (^{234}Th and ^{210}Pb) and X-radiographs suggest rapid particle mixing ($D_B \sim 90 \text{ cm}^2 \text{ yr}^{-1}$) and rapid sediment deposition ($\sim 1.5 \text{ cm month}^{-1}$). Thus it is possible that the bedform (and associated structures) are destroyed and recreated on a time-scale of a few months.

In press: Marine Geology.

Supported by: ONR Contract N00014-82C-0019.

WHOI Contribution No. 5460.

SEDIMENT DISCHARGE FROM THE INDUS RIVER
TO THE OCEAN: PAST, PRESENT AND FUTURE

John D. Milliman, G.S. Quraishie
and M.A.A. Beg

Until the late 1940's, the Indus River apparently discharged more than 600 mt of suspended sediment annually from the Himalaya Mountains. Assuming that most of the coarse fraction settled from suspension as the river flowed over the broad Indus Plain, somewhat less than 250 mt reached the Indus delta annually. Since then, various man-made structures have greatly decreased the sediment load of the Indus. Today, less than 50 mt reaches the estuary, and by the end of this decade it may be practically nothing. To date, downstream barrages have accounted for most of the sediment decrease; dams have apparently only affected upper portions of the river basin. Such a dramatic drop in sediment and water flux can have serious impact upon the coastal environment.

In press: Marine Geology and
Oceanography of the Arabian Sea and
Coastal Pakistan (Book).

Supported by: ONR Contract N00014-82C-
0009.

WHOI Contribution No. 5486.

TIDAL PHASE CONTROL OF SEDIMENT
DISCHARGE FROM THE YANGTZE RIVER

J.D. Milliman, Ya Hsueh, Dun-Xin Hu,
David J. Pashinski, Huang-ting Shen,
Zuo-sheng Yang and Peter Hacker

Tidal phase plays a major role in controlling sediment discharge from the Yangtze River estuary in eastern China. Direct measurements indicate that during spring tide in mid-November 1981 approximately 3 times the sediment passed the main channel of the river as during the next neap tide, 3 days later. The estuary presumably acts as a conduit for river sediment during spring tide but as a sink during neap tide.

In press: Estuarine Coastal and Shelf
Science.

Supported by: NOAA, ONR Contract
N00014-82-C-0009 and the Andrew
Mellon Foundation.

WHOI Contribution No. 5357.

TRANSFER OF PARTICULATE ORGANIC CARBON
AND NITROGEN FROM THE YANGTZE RIVER TO
THE OCEAN

John D. Milliman, Xie Qinchon and
Yang Zuosheng

The Yangtze River transports 12 million tons of particulate organic carbon (POC) to the oceans annually, with a computed yield of 6 t km² of drainage basin area. Concentrations of POC can exceed 80 mg/L in near bottom waters but generally range from 4 to 40 mg/L throughout the water column. Most POC percentages vary from 1.7 to 4.0 percent, the percentage decreasing with increasing suspended matter concentration although at lesser gradient than that predicted by Meybeck (1981, 1982). The divergence from Meybeck's world curve may be related to the high degree of human activity in the Yangtze drainage basin, or it may reflect lack of data from large rivers with high suspended loads. If the Yangtze values are representative of other turbid Asian rivers, present global estimates of fluvially-derived POC may be as much as 50 percent too low.

In press: American Journal of Science.

Supported by: NOAA, and ONR Contract
N00014-81C-0009.

WHOI Contribution No. 5433.

FLUX OF SUSPENDED SEDIMENT IN THE
CHANG JIANG (YANGTZE) ESTUARY

J.D. Milliman, Z.-s. Yang and R.H. Meade

The suspended matter regime within the Chang Jiang estuary is strongly dependent upon river flow, tidal stage and tidal phase as well as location within the estuary itself. Averaged over a complete tidal cycle, net transport of sediment is mostly offshore, although during low river runoff, near-bottom transport may be landward in controlling sediment flux, the estuary tending to act as a conduit during spring tide and trap during neap tide. As a result, sediment flux at spring tide during low river stages can approximate the flux at neap tide during high river runoff.

Further studies of sediment and water fluxes within the Chang Jiang estuary should involve synoptic observations. Two or more ships should take similar measurements at a variety of locations within the estuary. Observations should be taken for at least a fortnight in order to take into account the strong changes in hydrographic and sedimentary conditions that are related to tidal phase.

Published in: Proceedings to the
International Symposium on
Sedimentation on the Continental
Shelf. China Ocean Press.

Supported by: NOAA Cooperative
Agreement No. NA81AA-H-00008 and
ONR Contract N00014-81-C-0009.

WHOI Contribution No. 5355.

PHAEODARIAN SKELETONS: ROLE OF SILICA
TRANSPORT TO THE DEEP SEA AND
DISSOLUTION

Kozo Takahashi, David C. Hurd and
Susumu Honjo

The skeletons of phaeodarian Radiolaria play an important role in transporting and redistributing silica to the tropical deep oceans by dissolving in the water column and in the sea-floor. The skeletons are initially solid but within a few days to months become progressively porous while settling through the water column. Phaeodarian Radiolaria are rarely preserved in the bottom sediments, in contrast to polycystine Radiolaria which are the dominant Radiolaria preserved in oceanic sediments. This preservational difference is suggested to be due to differences in skeleton constitution between the two groups.

Published in: Science, 222, 616-618,
1983.

Supported by: NSF Grant OCE80-19386.

WHOI Contribution No. 5280.

LATE QUATERNARY SEDIMENTATION ON THE
SOUTHERN SOHM ABYSSAL PLAIN

Gustavs Vilks, D.E. Buckley
and L.D. Keigwin, Jr.

Six closely spaced sediment cores taken below the carbonate compensation depth penetrated fine to silty muds and entered sandy sediment at 10-12 meters below sea floor. Foraminiferal assemblages and $\delta^{18}O$ analyses on planktonic foraminifera indicated that the surface muds down to 2 meters are Holocene and derived from local promontories above the CCD. Below these sediments are about 6 meters of clays deposited during the late Wisconsin. These are unfossiliferous and have a possible northern source suggested by the higher chlorite content. Sandy sediments below 8 meters in the cores contain well preserved benthic foraminifera from the Scotian Shelf. Glacial $\delta^{18}O$ values on planktonic tests indicate the sandy sediments are most likely of latest Wisconsin

age. Thus during interglacials sand fraction of the southern Sohm Abyssal Plain sediments are mostly locally derived, but during glacial maxima the sediments have a distal northern source containing quartz sand that was initially deposited on the Scotian Shelf 1,500 km to the north.

Supported by: Sandia Laboratories
Contract 74-3133 and 13-9944.

WHOI Contribution No. 5503.

TECHNICAL REPORTS

COASTAL SEDIMENT TRANSPORT
POPPONESSET BEACH, MA

David G. Aubrey and Margaret R. Goud

Pathways and rates of near-bed sediment transport near Popponneset Beach, MA, were calculated using several distinct techniques. For the nearshore platform, sand transport in the form of sand waves was determined from vertical aerial photography spanning periods of four decades. In addition, calculations based on theoretical and empirical equations for near-bed sediment transport were made using field measurements of wind waves and tidal currents. Net sediment transport to the southwest inferred from these two techniques differed by about a factor of five. The higher net transport rate predicted in the aerial photographic method is a result of lack of wave measurements during storm conditions. Storm waves increase the net transport through a local increase in bed shear stress. Net transport to the southwest across the platform is between 700 and 3300 m³/yr.

Littoral sand transport along Popponneset Beach was calculated from one month of directional wave measurements, extrapolated to a yearly value using long-term meteorological observations. Littoral transport from these calculations is 10,000 m³/yr to the northeast, opposite the sense of alongshore transport in the shallow nearshore.

Patterns of shoreline change are discussed from a historical perspective, and using the transport calculations discussed above. Several management alternatives for coping with predicted shoreline change are presented for consideration by the Town of Mashpee.

W.H.O.I. Technical Report 83-26.

Supported by: Town of Mashpee;
Coastal Zone Management Program;
Sea Grant NA80AA-D-00077(RB-40); and
the ALCOA Foundation.

SEDIMENT TRANSPORT IN A TIDAL INLET

David G. Aubrey and Paul E. Speer

Various aspects of sediment transport in and around natural, unstructured tidal inlets were investigated over the two year period of study. Concentrating on two tidal inlets (Nauset Inlet and Popponesset Inlet, Cape Cod, MA) and combining detailed field observations with numerical model studies of tidal flows in inlet/estuarine environments, several aspects of tidal inlet behavior have been clarified. In addition, field work has resulted in a number of technical publications of general utility to a wide spectrum of coastal research interest. Primary scientific items addressed in this study include: 1) diagnostic numerical model of generation and propagation of tidal nonlinearities in shallow estuarine channels; 2) effects of flow curvature on tidal inlet sediment transport; 3) definition of mechanisms by which tidal inlets migrate in a direction opposite to the net littoral drift direction; 4) hypothesis of a mechanism for rapid barrier spit growth in locations with low rates of littoral transport; 5) clarification of long-term patterns of sea-level rise in the United States to assess its role in tidal inlet/estuarine evolution; 6) historical descriptions of massive inlet migration at two study inlets as supporting evidence for the inlet modeling studies. Technical information generated by the study includes a description of a low-cost, reliable method to join nearshore electrical cables; description and intercomparison of instrumentation and analysis routines for estimating directional spectral parameters from wave gage data; and development of a field system and laboratory analysis package for preparing accurate bathymetric charts in shallow, nearshore regions, using microwave navigation and precision echo-sounding.

W.H.O.I. Technical Report 83-20.

Supported by: U.S. Army Research Office Grant DAAG 29-81-K-0004 and the Department of Commerce, NOAA Grants NA79AA-D-00102 and NA80AA-D-00077.

W A V E M E C H A N I C S

FIELD INTERCOMPARISON OF NEARSHORE DIRECTIONAL WAVE SENSORS

W.G. Grosskopf, D.G. Aubrey
and M.G. Mattie

Five measurement strategies (four in situ, one remote) for estimating direc-

tional wave spectra were intercompared in a 1980 experiment at the Coastal Engineering Research Center's Field Research Facility in Duck, NC. The systems included two pressure sensor/biaxial current meter combinations (different manufacturers), a triaxial acoustic current meter, an SXY guage (square array of four pressure sensors), and a shore-based imaging radar. A detailed error analysis suggests sources for differences in estimated wave spectra from the different instruments; in general, they intercompare favorably. The major deviation among in situ gauges was associated with the triaxial acoustic current meter. Reliance on a vertical velocity measurement (instead of a direct pressure or sea surface elevation measurement) can contribute additional uncertainty in the directional spectral estimates. The imaging radar was successful in distinguishing multiple wave trains at the same frequency, which was not possible with the simple spectral estimation analysis applied to in situ data. However, the radar is not useful in providing accurate estimates of spectral density nor in distinguishing multiple wave trains of different frequencies coming from the same direction. Selection of a measurement strategy for a particular need depends on the precise data requirements for that application. Although the five tested intercompared well, in practice not all are equally suitable for every application.

Published in: Journal of Oceanic Engineering, IEEE, OE-8, 254-271, 1983.

Supported by: Exxon Production Research Co. under Technical Services Agreement PR-6520; NOAA Sea Grant NA79AA-D-00102, NA80AA-D-00077; and U.S. Army Contract No. DAAG29-81-K-0004.

WHOI Contribution No. 5222.

DEPARTMENT OF OCEAN ENGINEERING

Robert C. Spindel, Department Chairman

OCEAN ENGINEERING

ACOUSTIC SIGNAL
PROCESSING

CONVERGENCE ANALYSIS OF LMS FILTERS
WITH UNCORRELATED GAUSSIAN DATA

A. Feuer and E. Weinstein

Statistical analysis of the Least Mean Squares (LMS) adaptive algorithm with uncorrelated Gaussian data is presented. Exact analytical expressions for the steady state mean square error (m.s.e.) and the performance degradation due to weight vector misadjustment are derived. Necessary and sufficient conditions for the convergence of the algorithm to the optimal (Wiener) solution within a finite variance are derived. It is found that the adaptive coefficient, which controls the rate of convergence of the algorithm must be restricted to an interval significantly smaller than the domain commonly stated in the literature. The outcome of this paper, therefore, places fundamental limitations on the m.s.e. performance and rate of convergence of the LMS adaptive scheme.

Accepted: IEEE Transactions on
Acoustics Speech and Signal
Processing.

WHOI Contribution No. 5483.

ANGLE-INDEPENDENT DOPPLER MEASUREMENT

Nadav Levanon and Ehud Weinstein

The well-known relation between the radar Doppler shift and the target velocity requires knowledge of the angle between the target heading and the radar beam. A new explicit relation is presented which is angle-independent, however, it requires the first and second time derivatives of the Doppler shift. An optimal measuring scheme to obtain those derivatives is suggested, based on phase observations of the received signal and a least square filter. Error analysis of the angle-independent velocity measurement is provided and compared with the angle-dependent one.

Accepted: IEEE Transactions on
Aerospace and Electronic Systems.

Supported by: ONR Contract
N00014-77-C-0196.

WHOI Contribution No. 4935.

DELAY AND DOPPLER ESTIMATION BY
TIME-SPACE PARTITION OF THE ARRAY DATA

Ehud Weinstein and Doron Kletter

An instrumentationally attractive technique for estimating the Doppler time-compression and its time derivatives is proposed and analyzed in this paper. The basic scheme is illustrated in Figure 5. It consists of $(M - 1)$ Generalized Cross-Correlators (GCC) in parallel (where M is the number of sensors in the receiving array), each of which is implemented successively to generate a sequence of differential delay measurements. The resulting spatial-temporal differential delay data is then compressed into the estimates of the Doppler and its derivatives using a Least Squares (LS) filter. It is shown that for signal-to-noise ratio (SNR) conditions above the threshold level, the indicated two-step estimation procedure is very nearly optimal. When operating near the threshold point, however, the proposed scheme is only distinctly suboptimal.

Accepted: IEEE Transactions on
Acoustics, Speech and Signal
Processing.

Supported by: ONR Contract
N00140-82-M-MK39.

WHOI Contribution No. 5339.

FUNDAMENTAL LIMITATIONS IN PASSIVE
TIME-DELAY ESTIMATION
PART I: NARROW-BAND SYSTEMS

Anthony J. Weiss and Ehud Weinstein

Time delay estimation of a noise-like random signal observed at two or more spatially separated receivers is a problem of considerable practical interest in passive radar/sonar applications. A new method is presented to analyze the mean square error performance of delay estimation schemes based on a modified (improved) version of the Ziv-Zakai lower bound (ZZLB). This technique is shown to yield the tightest results on the attainable system performance for a wide range of signal-to-noise ratio (SNR) conditions.

For delay estimation using narrowband (ambiguity-prone) signals, the fundamental result of this study is illustrated in Figure 3. The entire domain of SNR is divided into several disjoint segments indicating several distinct modes of operation. If the available SNR does not exceed SNR_1 , signal observations from the receiver outputs are completely dominated by noise thus essentially useless for the delay estimation. As a result, the attainable mean

square error, e^2 , is bounded only by the a-priori parameter domain. If SNR_1 is less than SNR and SNR is less than SNR_2 , the modified ZZLB coincides with the Barankin bound. In this regime, differential delay observations are subject to ambiguities. If SNR is greater than SNR_3 , the modified ZZLB coincides with the Cramer-Rao lower bound indicating that the ambiguity in the differential delay estimation can essentially be resolved. The transition from the ambiguity-dominated mode of operation to the ambiguity-free mode of operation starts at SNR_2 and ends at SNR_3 . This is the threshold phenomenon in time delay estimation.

The various deflection points, SNR_1 and the various segments of the bound (Figure 3), are given as functions of such important system parameters as time-bandwidth product (WT), signal bandwidth to center frequency ratio (W/l_0) and the number of half wavelengths of the signal center frequency contained in the spacing between receivers. With this information, the composite bound illustrated in Figure 3 provides the most complete characterization of the attainable system performance under any prespecified SNR conditions.

Published in: IEEE Transactions on
Acoustics, Speech and Signal
Processing, ASSP-31, 472-485, 1983.

Supported by: ONR Contract
N00014-82-C-0152.

WHOI Contribution No. 5326.

FUNDAMENTAL LIMITATIONS IN PASSIVE TIME DELAY ESTIMATION PART II: WIDEBAND SYSTEMS

Ehud Weinstein and Anthony J. Weiss

This is the second part of the study which deals with the problem of time delay estimation between noise-like Gaussian signals radiated from a common point source and observed at two spatially separated receivers, each of which also receives uncorrelated Gaussian noise. A modified (improved) version of the Ziv-Zakai lower bound (ZZLB) is used to analyze the attainable mean square estimation errors. In Part I we concentrated on narrowband signals. The outstanding observed feature was a distinct threshold phenomenon whereby below a certain signal-to-noise ratio (SNR), the lower bound increases suddenly and precipitously. In Part II we concentrate on wideband signals. If we relax the assumption concerning signal bandwidth, we discover a second threshold phenomenon occurring at a lower SNR . The two threshold effects separate the entire SNR domain into

three disjointed segments indicating three distinct modes of operation. At high SNR , the attainable mean square error (m.s.e.) is characterized by the Cramer-Rao lower bound (CRLB). At moderate SNR (between the two threshold points), the modified ZZLB coincides with the Barankin lower bound, which exceeds the CRLB by a factor of the order of $(l_0/W)^2$ where W and l_0 are, respectively, signal bandwidth and center frequency. At low SNR , the attainable m.s.e. is bounded only by the variance of the a-priori search domain of the delay parameter.

Accepted: IEEE Transactions on
Acoustics, Speech and Signal
Processing.

Supported by: ONR Contract
N00014-82-C-0152.

WHOI Contribution No. 5472.

C O A S T A L D Y N A M I C S

RIPPLED SCOUR DEPRESSIONS ON THE CONTINENTAL SHELF OFF CENTRAL CALIFORNIA

D.A. Cacchione, D.E. Drake,
W.D. Grant and G.B. Tate

Side-scan sonar records taken during the recent Coastal Ocean Dynamics Experiment (CODE) show elongate, shore-normal rippled depressions of low relief on the inner continental shelf off central California between Bodega Bay and Point Arena. These features extend up to 2 kilometers from the coast into water depths to 65 meters. The proposed mechanism for their generation is storm-generated bottom currents associated with coastal downwelling during the late fall and winter which scour the surficial fine sand sediment and expose the coarser sand substrate in the depressions. The zones of most intense erosion and the irregular spacing of the features may be controlled by submerged rock ledges and other prominent coastal features. The large straight-crested ripples within the depressions (heights to 40 cm; wave lengths to 1.7 m) are probably formed by large-amplitude, long-period surface waves generated by winter storms.

In press: Journal of Sedimentary
Petrology.

Supported by: NSF Grant
OCE80-14938 and USGS.

WHOI Contribution No. 5469.

VARIABILITY OF SEA-FLOOR ROUGHNESS
WITHIN THE COASTAL OCEAN DYNAMICS
EXPERIMENT (CODE) REGION

David A. Cacchione, David E. Drake,
William D. Grant, Albert J. Williams III
and George B. Tate

This report summarizes the geological and biological data taken on the northern California Continental Shelf before and during the Coastal Ocean Dynamics Experiment (CODE) by the principal investigators of the bottom stress/bottom boundary layer component of CODE (D. Cacchione, D. Drake, USGS and W. Grant, A. Williams, WHOI). The report concentrates on a description of the seafloor relief that likely influences the near bottom flow field at the CODE site, and that significantly affects bottom stress estimates. The general physical characteristics of the CODE site, the macro- and micro-physiography, sediment distribution and discharge of the larger coastal rivers are described. On a large scale, the shelf is relatively narrow, diminishing in width from SE to NW with an average cross-shelf gradient of 0.004. The outer shelf is flat and floored by relict sand to the south and sandy-silt to the north. The central shelf is also flat and is floored by silt. The inner shelf is sandy with irregularly spaced submerged outcrops. On the small scale, the central shelf surface is highly bioturbated, consisting of animal tracks, mounds and furrows. The inner shelf is covered with oscillatory ripple marks.

Supported by: NSF Grant
OCE80-14938; OCE80-14941.

WHOI Technical Report No. 83-25.

BOTTOM STRESS ESTIMATES AND THEIR
PREDICTION ON THE NORTHERN
CALIFORNIA CONTINENTAL SHELF DURING
CODE-1: THE IMPORTANCE OF WAVE-CURRENT
INTERACTION

W.D. Grant, A.J. Williams
and S.M. Glenn

High quality near-bottom boundary layer measurements obtained at a mid-shelf location (90 m water depths) in the CODE region off Northern California are described. Bottom tripod velocity measurements and supporting data obtained during typical spring and early summer conditions (June, 1981, during Code-1) are analyzed to obtain both velocity profiles and mean bottom stress and bottom roughness estimates. During the time period described, the mean near-bottom (< 2 m) velocity profiles are highly logarithmic ($R > 0.997$) approximately 30 percent of the time. Effects

induced by unsteadiness from internal waves result in some degradation of the profiles ($0.96 \leq R \leq 0.997$) the rest of the time. Mean stress profiles indicate the logarithmic layer is approximately a constant stress layer. The near bottom flow field is composed of mean currents and oscillatory currents due to swell. Typical mean u_* values estimated from measurements greater than 30 cm above the bottom have magnitudes of 0.5-1.0 cm/sec; mean stress values are three to seven times larger than expected from predictions using a typical smooth bottom drag coefficient and one and one half to three and one half times larger than expected for predictions using a drag coefficient based on the observed rough bottom. Corresponding z_0 values have magnitudes of approximately 1 cm; an order of magnitude larger than the observed physical bottom roughness. These values are demonstrated to be consistent with those expected from theoretical models for combined wave and current flows. The u_* values estimated from the CODE-1 data and predicted by the Grant and Madsen (1979) model typically agree within 10-15 percent. The waves influencing the mid-shelf bottom stress estimates are 12-20 second swell associated with distant Pacific storms. These waves are present over most of the year. The results demonstrate that waves must be taken into account in predicting bottom stress over the Northern California Shelf and that these predictions can be made using existing theory.

Accepted: Journal of Physical
Oceanography.

Supported by: NSF Grant
OCE80-14938.

WHOI Contribution No. 5432.

CONTINENTAL SHELF BOTTOM BOUNDARY
LAYER MODEL
VOLUME I: THEORETICAL MODEL DEVELOPMENT

William D. Grant and Scott M. Glenn

A continental shelf bottom boundary layer model is presented which includes the effects of (1) combined wave and current interaction, (2) moveable bed effects (ripples, near bed transport, bioturbation), (3) stratification due to temperature and salinity, (4) stratification due to suspended sediment, and (5) planetary rotation. The theoretical background for each component included in the model is developed and the limitations of the approach taken are carefully discussed. Special attention is given to treating the problem of unsteady oscillatory flow and its effects on the mean flow field and

sediment transport characteristics important to predicting boundary layer velocity structure.

The model was developed for the purpose of predicting the near bottom velocity field necessary to calculate forces on submarine pipelines, but its usefulness is much more general. It predicts the vertical velocity profile, steady and unsteady components of the boundary shear stress, the sediment concentration field and near bed sediment transport.

Published in: Final Report to
American Gas Association, 1, 1983.

Supported by: American Gas Association
Project No. PR-153-126.

CONTINENTAL SHELF BOTTOM BOUNDARY
LAYER MODEL
VOLUME II: MODEL/DATA COMPARISON

W.D. Grant and S.M. Glenn

High quality bottom boundary layer measurements from the CODE site are described. Bottom tripod velocity measurements and supporting data typical of everyday spring and summer conditions and during a winter storm are analyzed to obtain both velocity profiles and mean bottom stress and bottom roughness estimates. These values are compared to the AGA model developed in Volume I. The everyday measurements were taken during CODE-1 at C3 (90 m) by Grant and Williams, WHOI; the winter storm data were taken prior to CODE at the R2 (80 m) site by Cacchione and Drake, USGS.

The mean-near bottom ($< 2m$) velocity profiles are logarithmic ($r^2 > 0.993$) much of the time for everyday flows; deviations are primarily due to kinematical effects induced by unsteadiness from internal waves. Stress profiles show the logarithmic layer corresponds to a constant stress layer as expected for the inertial region of a boundary layer. Stress estimates made from dissipation and profile techniques agree at 95 percent confidence level. Typical z_0 values are an order of magnitude larger than the physical bottom roughness. Typical u_* values are more than twice as large as expected from a usual drag law prediction (this corresponds to over a factor of four in stress). These values are demonstrated to be consistent with the effects of combined waves and currents (Volume I, Grant and Madsen, 1979, and Smith, 1977). The u_* values estimated from the CODE-1 data and predicted by the Grant and Madsen (1979) model are typically within 10-15 percent. Similar results are demonstrated for the winter storm conditions during which large sed-

iment transport occurs.

The waves influencing the mid-shelf bottom stress estimates are 14-20 second swell associated with Southern Pacific storms. These waves are present over most of the year. The results clearly demonstrate that waves must be taken into account in predicting near bottom velocity profiles at the CODE site.

Published in: Final Report to Pipeline
Research Committee, 2, 1983.

Supported by: American Gas Association
Project No. PR-153-126.

HIGH FREQUENCY BOTTOM STRESS
VARIABILITY AND ITS PREDICTION
IN THE CODE REGION

W.D. Grant, A.J. Williams, 3rd,
S.M. Glenn, D.A. Cacchione
and D.E. Drake

High quality bottom boundary layer measurements obtained in the CODE region off Northern California are described. Bottom tripod velocity measurements and supporting data obtained during typical spring and early summer conditions and during a winter storm are analyzed to obtain both velocity profiles and mean bottom stress and bottom roughness estimates. The spring/summer measurements were taken in June 1981 during Code-1 at C3 (90 m) by Grant and Williams, WHOI; the winter storm data were taken in November, 1980 prior to CODE-1 at the R2 (80 m) site by Cacchione and Drake, USGS.

The mean near-bottom ($< 2m$) velocity profiles are logarithmic ($r^2 > 0.993$) much of the time for everyday flows; deviations are primarily due to kinematical effects induced by unsteadiness from internal waves. Stress profiles show the logarithmic layer corresponds to a constant stress layer as expected for the inertial region of a boundary layer. Stress estimates made from dissipation and profile techniques agree at the 95 percent confidence level. Typical z_0 values estimated from measurements greater than 30 cm above the bottom have magnitudes of approximately 1 cm; an order of magnitude larger than the physical bottom roughness. Corresponding u_* values have typical magnitudes of 0.5-1.0 cm/sec; more than twice as large as expected from a usual drag law prediction (corresponding to over a factor of four in mean stress). These values are demonstrated to be consistent with those expected for combined wave and current flows predicted theoretically by Grant and Madsen (1979) and Smith (1977). The u_* values estimated from the CODE-1 data and predicted by the Grant and Madsen (1979)

model typically agree within 10-15 percent. Similar results are demonstrated for the winter storm conditions during which large sediment transport occurs. (Typical z_0 values are 4-6 cm; typical u_* values are 3-6 cm/sec.)

The waves influencing the mid-shelf bottom stress estimates are 14-20 second swell associated with Southern and Western Pacific storms. These waves are present over most of the year. The results clearly demonstrate that waves must be taken into account in predicting bottom stress over the Northern California Shelf.

Supported by: NSF Grant
OCE80-14938.

WHOI Technical Report No. 83-19.

C O M P U T E R S C I E N C E

FINDIF: A SOFTWARE PACKAGE TO CREATE
SYNTHETIC SEISMOGRAMS BY FINITE
DIFFERENCES

Mary M. Hunt, Lee Gove
and Ralph A. Stephen

In order to study seismic wave propagation through laterally varying sea floor structures, a software package has been created to generate synthetic seismograms by finite differences. The elastic wave equation can be solved in two dimensions either for point sources in cylindrical coordinates or for line sources in rectangular coordinates. Vertical and radial variations of the elastic parameters are allowed.

The package includes four programs. Input to the system consists of a short file containing parameter values to describe the model. The first program is used to initialize the system for the particular model being used. The source arrays and velocity matrices are each computed by a separate program. The final program, which actually carries out the finite difference calculations, includes six subroutines to implement different options based on alternative finite difference formulations. Two different kinds of output files are created by this program: one or more snap-shot files, and a one time series file, which will usually include more than one series.

Supported by: ONR Contract
N00014-82-C-0019.

WHOI Technical Report No. 83-42.

G E O L O G Y

THE EAST PACIFIC RISE NEAR 21°N, 13°N
AND 20°S: INFERENCES FOR
ALONG-STRIKE VARIABILITY OF AXIAL PROCESSES
OF THE MID-OCEAN RIDGE

Jean Francheteau and
Robert D. Ballard

Data obtained along various segments of the Mid-Ocean Ridge (MOR) are used to construct an idealized model for crustal accretion. The model seeks to predict the topographic, volcanic, tectonic, and hydrothermal characteristics of any given spreading segment of the MOR as a function of distance away from the bounding transform faults. This model is based on a series of detailed mapping efforts carried out on segments of the MOR having a broad range of spreading rates between 2 and 16 cm/yr. This paper includes the results of the French SEARISE program carried out during the summer of 1980 aboard the N/O JEAN CHARCOT and two American cruises conducted in 1981 aboard the R/V MELVILLE.

Published in: Earth and Planetary
Science Letters, 64, 93-116, 1983.

Supported by: ONR Contract
N00014-82-C-0019; NSF Grant
OCE80-20425.

WHOI Contribution No. 5218.

GEOLOGIC PROCESSES OF THE MID-OCEAN
RIDGE AND THEIR RELATIONSHIP
TO SULFIDE DEPOSITION

Robert D. Ballard and
Jean Francheteau

Enough detailed data has been obtained along the axis of the Mid-Ocean Ridge to construct a kinematic model which attempts to describe observed variations in time, space, and spreading rate of volcanic, tectonic, and hydrothermal processes and their relationship to the deposition of massive sulfide deposits. The model draws upon submersible and towed camera mapping programs in the FAMOUS area of the Mid-Atlantic Ridge where the spreading rate is approximately 2 cm/yr. In addition, recent efforts along various segments of the East Pacific Rise and Galapagos Rift have the most profound impact on the model as they involve spreading rates ranging from 6.0 to 10.2 cm/yr where hydrothermal circulation and associated massive sulfide deposition is well developed.

Published in: Marine Polymetallic
Sulfides: A National Overview and
Future Needs, Workshop Proceedings,
47-62, January 19-20, 1983.

Supported by: NSF Grant OCE80-20425.

WHOI Contribution No. 5300.

INTENSE HYDROTHERMAL ACTIVITY AT THE
AXIS OF THE EAST PACIFIC RISE NEAR 13°N:
SUBMERSIBLE WITNESSES THE GROWTH OF
SULFIDE CHIMNEY

R. Hekinian, J. Francheteau, V. Renard,
R.D. Ballard, P. Choukroune, J.L. Cheminee,
F. Albarede, J.F. Minster, J.C. Marty,
J. Boulegue and J.L. Charlou

A submersible study of a fast spreading (12 cm/yr) Mid-Ocean Ridge segment led to the discovery of intense hydrothermal activity (East Pacific Rise near 12°50'N). Twenty-four sites with active vents and sixty inactive hydrothermal deposits were found within a narrow graben less than 300 m wide along a 20 km long segment of the ridge crest. The graben is floored with fresh basaltic sheet flows (including collapsed pits or lava lake structures). From both deep towed camera stations and manned submersible observations, it is estimated that the average spacing of the hydrothermal deposits along the rise axis (10-50 m in diameter) lies between 100 to 200 meters.

The hydrothermal deposits found in the central graben are believed to have formed rapidly (on the order of a few years). Detailed investigations of one active site have enabled us to witness the growth of an active chimney which increased its size by 40 cm in five days.

The most extensive hydrothermal deposits were found on an off-axis seamount located 4 km east of the ridge axis. The hydrothermal deposits found on both the ridge axis and on the seamount are similar in composition and consist essentially of zinc, copper and iron mineral phases.

In press: Earth and Planetary Science Letters.

Supported by: NSF Grant OCE80-20425.

WHOI Contribution No. 5271.

PROJECT CYATHERM: DETAILED
INVESTIGATION OF HYDROTHERMAL ACTIVITY
AT 12°48'N ON THE EAST PACIFIC RISE

R.D. Ballard

In 1980, the French Government began a long-term investigation of the East Pacific Rise (EPR) between the Orozco (15°N) and Clipperton (10°N) fracture zones (Figure 1). The objective of this investigation is to better understand the axial processes of a fast spreading ridge (12 cm/yr) and compare those findings with similar efforts already carried out at moderate and slow spreading ridge segments. More specifically, the work between 10° and 15°N seeks to determine the nature of hydrothermal circulation along the axis of the Mid-Ocean Ridge (MOR) and its relationship to the occurrence and distribution of massive sulfide accumulations.

In press: Earth and Planetary Science Letters.

Supported by: NSF Grant
OCE80-20425; ONR Contract
N00014-82-C-0019.

WHOI Contribution No. 5431.

I N S T R U M E N T A T I O N

A CCD CHIRP-Z FFT DOPPLER SIGNAL
PROCESSOR FOR LASER VELOCIMETRY

Y.C. Agrawal

A charge-coupled Chirp-Z fast Fourier transform device which has been configured to obtain the frequency of peak spectral amplitude in signals from laser velocimeters is described for use in oceanographic and other low laser power field applications where poor optical signals and frequent drop out are encountered. The resulting signal processor gives 8 bit resolution in the velocity word, with a cycle time of approximately 4 msec while scanning a spectral range of 0-125 kHz; higher frequency ranges are possible at proportionately faster spectral scan rates, up to a maximum of 2 MHz. Error estimates are presented which indicate 2 bit accuracy at signal-to-noise power ratio of -2 db. The combination of this processor together with a photodiode array detector described by the author (Agrawal and McCullough, 1981) radically simplifies the practice of laser velocimetry.

Accepted: Journal of Physics E:
Scientific Instruments.

Supported by: ONR Contract
N00014-82-C-0019; Sea Grant
NA-80-AA-00077.

WHOI Contribution No. 5314.

DIRECTIONAL LASER VELOCIMETRY WITHOUT FREQUENCY BIASING - PART II

Y.C. Agrawal and J.B. Riley

In an earlier paper, Agrawal and McCullough (1981) demonstrated that a single laser beam, focused at the point of measurement, can be used to obtain both the magnitude and direction of local fluid velocity without the use of beam splitting optics or frequency biasing devices. Additionally, the photoelectric signal was shown to be a zero-mean sine wave suitable for existing counter type laser Doppler velocimeter signal processors. The method described in the above paper indicated the use of two offset photo-detector arrays; the resulting photocurrents from the two arrays were correlated to give direction. In this paper, we describe the implementation of the method using a self scanning diode array. Detector sensitivity in the array is discussed and microchannel plate detectors are suggested for increased sensitivity.

In press: Applied Optics.

Supported by: ONR Contract
N00014-82-C-0019; NOAA Sea Grant
NA80AA-D-00077.

WHOI Contribution No. 5314.

CTD LOWERING MECHANICS

H.O. Berteaux and R.G. Walden

The many difficulties encountered in the deployment of CTD instrument packages, including the loss of an appreciable number of them, has prompted a comprehensive study of CTD instrument lowering mechanics. Conditions causing large cyclic cable tensions, shock loads or cable slack had to be investigated, modeled, and, hopefully, measured in situ. The dynamic behavior of the instrument package needed to be assessed. Methods to improve its flight pattern as it travels through the water column and recommendations for safer and more efficient lowering procedures had to be formulated. This paper describes how the study was conducted and presents its salient findings and recommendations. In its conclusion, the paper points out the limits of present CTD lowering systems and advocates that servo-controlled motion compensating winches be used in the future.

In press: Deep Sea Research.

Supported by: ONR Contract
N00014-79-C-0071.

WHOI Contribution No. 5311.

EQUIPMENT LOWERING MECHANICS

H.O. Berteaux

Improper design considerations while lowering or retrieving equipment at sea can result in substantial cable damage including rupture followed by the total loss of valuable payload.

The hydrodynamic behavior of the payload at the end of the lowering cable is often unknown or ignored. Yet the shape and weight of the payload has considerable bearing on the cable performance. For example, a payload with a large drag area can substantially add to cable static tension when the cable is hauled in. Conversely, while paying out the payload may fall slower than the cable itself, thereby creating a slack cable condition. This slack condition may result in a subsequent snap load when the lowering is stopped and the payload impacts the cable, or in a kink in the wire, or both. Payload spinning, kiting and tumbling can also obviously impair an orderly lowering.

Recurrent causes of cable damage include loading the cable beyond its yield point (or breaking strength), fatigue failure, and kinking. In this section, the mechanisms which create these detrimental conditions are first reviewed. These potential failure mechanisms include quasi-static tensile loads, wave induced dynamic loads, zero load (slack cable), and impact (snap) loads.

The mathematical concepts which are used to predict and quantify these causes of cable failure are also reviewed and their use illustrated with a few typical examples.

In the last part of this chapter specific design recommendations are made for improving the payload hydrodynamic behavior. Finally, the operational limits, such as maximum length of cable paid out or allowable payout rates, are discussed.

Published in: Handbook of
Oceanographic Winch, Wire and Cable
Technology. A.H. Driscoll, ed.
February, 1983.

Sponsored by: ONR Contract
N00014-82-C-0116 (at U.R.I.).

EXPERIMENTAL EVALUATION OF CTD PACKAGE
HYDRODYNAMIC BEHAVIOR AND RECOMMENDATIONS
FOR IMPROVED LOWERING TECHNIQUES

H.O. Berteaux, R.G. Walden,
P.R. Clay and R.E. McDevitt

This report is the last of a series of three reports on a comprehensive study of CTD instrument lowering mechanics. The first report, WHOI 79-81, "A Study of CTD Cables and Lowering Systems," examines the causes and modes of lowering cable failures, both mechanical and electrical, and makes recommendations to improve existing instrument packages and lowering procedures. The second report, WHOI 81-76, "Hydrodynamics of CTD Instrument Packages," is a theoretical study of instrument package stability when cable-lowered or free-falling. The model is used to predict the hydrodynamic response of CTD packages in their present or improved configuration. This report, WHOI 83-21, is more factual. It describes the tests performed on scale models and actual CTD packages to actually observe and/or measure their hydrodynamic behavior. Analytical results and experimental data obtained in this study are used to draw recommendations for CTD package improvement and future lowering procedures.

Supported by: ONR Contract
N00014-72-C-0019.

WHOI Technical Report No. 83-21.

AIR DEPLOYED OCEANOGRAPHIC
MOORING - AB1034

L.W. Bonde, D.B. Dillon, E.J. Softley,
R. Walden, H. Berteaux and T. Popp

In response to the need for low cost, rapidly deployed, deep ocean, instrumented data platforms, a program was sponsored by the Office of Naval Research to develop an Air Deployed Oceanographic Mooring (ADOM). This mooring consists of a subsurface buoy, taut moored by a cable with in line sensors, and contains a computer for data acquisition and processing. A surface float housing a satellite transmitter and antenna is tethered to the subsurface buoy by a compliant data line (Figure 1). The design evolved from a feasibility study, theoretical analyses, and testing in the laboratory, dockside, and at sea.

ADOM is bolted together for launch as a single cylinder (Figure 2). The anchor shell contains the bottom finder assembly, mooring line, mooring lock-up mechanism, and sensor array. The computer controller is housed in the subsurface buoy. The parachute deploys from one end.

The parachute is deployed in stages by explosive line cutters. It is released and the mooring deployed by explosive bolts fired in sequence using batteries activated by seawater.

A one-year active life is attained by using low power "CMOS" for the processor, memory, and sensor circuits. A unique scheme was devised to address the sensors. An array of temperature sensors has been integrated into a single conductor armored cable.

The program included a series of system deployments at sea. Temperature data were acquired by ADOM and telemetered to the shore station via the USAF LES-9 satellite according to a pre-set schedule. The test program concluded with two air drops from a U.S. Marine Corps C-130 aircraft. These tests demonstrated that the Air Deployed Oceanographic Mooring is a practical method for ocean data acquisition.

In press: Proceedings Marine
Technology Society and IEEE
Annual Convention, San Francisco,
CA, September 1983.

Supported by: ONR Contract
N00014-77-C-0230.

WHOI Contribution No. 5457.

A COHERENT APPROACH TO INSTRUMENT
INTERCOMMUNICATION AND TESTING USING SAIL

Albert M. Bradley
and William E. Terry, Jr.

The SAIL protocol for instrument intercommunication has become a standard in the oceanographic community during the past several years. Several autonomous instruments, not designed for shipboard data acquisition, have been developed which utilize the SAIL protocol for communication during development and predeployment testing. The ease with which instruments may be added to the loop allows one to test many different instruments in remote locations from one central point. A logical extension of this capability is the development of smart controllers to handle routine test procedures automatically.

This paper describes an addition to the SAIL protocol to allow many levels of smart controllers. Provisions are allowed for using a controller as a modem to communicate with any instrument in a lower level loop. Potential problems with loop errors, real time clocks and data formatting are discussed. The result is a highly flexible addition to the growing uses of SAIL.

Published in: Proceedings of Third
Working Symposium on Oceanographic
Data Systems, IEEE Computer
Society, 51-54, October, 1983.

Sponsored by: ONR Contract
N00014-82-C-0019; NASA Grant
NAG-5-281.

INTEGRATED COMMUNICATIONS IN BUOY SYSTEMS

Ed Mellinger and Al Bradley

Buoy systems designed for oceanographic research have traditionally employed underwater sensors distributed along a mooring line some kilometers in length. The sensor packages have evolved as autonomous data acquisition units, each equipped with local processors, data recorders, and power supplies. Emerging applications, including event driven sampling schemes and satellite data collection systems, will require a more general system architecture capable of bringing distributed sensor data to a central point for analysis and/or relay to shore. In this paper we propose to adapt the shipboard Serial ASCII Instrumentation Loop (SAIL) system architecture to the underwater environment. Necessary additions to the SAIL protocol are discussed, and an implementation of an underwater SAIL based system is described.

Published in: The Marine Technology
Society-NOAA Data Buoy Center-1983
Symposium on Buoy Technology,
April, 1983.

Supported by: NASA Grant NAGW-265;
NASA Grant NAG-5-281.

WHOI Contribution No. 5345.

THE LOTUS DISCUS MOORING

Peter R. Clay

The Woods Hole Oceanographic Institution has been involved in a Long Term Upper Ocean Study (LOTUS) for the past three years. The goal of the project is to obtain a continuous two year record of atmospheric data combined with current and temperature measurements in the upper water column. An engineering test stage of two years preceded the experiment during which time much was learned and incorporated in later operational moorings. Two of four moorings in the experiment were heavily instrumented. This paper discusses the initial surface mooring design, subsequent changes and finally some of the engineering data obtained.

Published in: Proceedings of Marine
Technology Society, 1, 108-113,
1983.

Sponsored by: ONR Contract
N00014-76-C-0197.

WHOI Contribution No. 5354.

SEISMIC REFLECTION PROFILING WITH DEEP TOWED HYDROPHONE ARRAY

J. Collins, D. Koelsch and G. Purdy

A seismic reflection profiling system consisting of a 264 m long, deep-towed, 15-element, end-fire, vertical array and a 40 cubic inch airgun was successfully used to profile a sediment pond in the trough of the inactive segment of the Kane Fracture Zone close to its intersection with the Mid-Atlantic Ridge at 24°N. The increased signal-to-noise ratio achieved with the array demonstrates that it is a useful tool for detailed seismic profiling in areas of rough topography in the deep ocean.

In press: Marine Geophysical
Researches.

Supported by: NSF Grant OCE80-25206.

WHOI Contribution No. 5443.

DIRECT OPTICAL ASSESSMENT OF MACROSCOPIC AGGREGATES IN THE DEEP OCEAN

Susumu Honjo, Kenneth W. Doherty,
Yogesh C. Agrawal and Vernon L. Asper

Large amorphous aggregates (LAA), frequently called "marine snow," may accelerate the flux of fine particles and dissolved matter in the ocean by adsorption. The aggregates are extremely fragile and their sampling with conventional towing and casting is impossible. To assess the size and spatial distribution of such aggregates, we have developed a photographic system that can be lowered to full ocean depth. The system, based on light scattering in water, uses a well-collimated beam pointed downward from the verticle at an angle of 45°. The beam is produced by placing high-powered stroboscopic lights at the back focal plane of a Fresnal lens system. Illuminated aggregates in the light are photographed from a direction perpendicular to the beam axis. The photographed volume, 0.66 m³, is 106 x 156 x 40 cm. The resolution of the instrument is about 0.2 mm. Aggregates were distributed throughout the water column at all stations along a 200-km transect off Monterey Bay. The abundance of macroscopic aggregates (less than 0.5

mm) ranged from several per liter near the surface to 1 or less at depth.

In press: Deep Sea Research, 31(1), 1984.

Supported by: NSF Grant OCE80-25209.

WHOI Contribution No. 5036.

AN EVENT RECORDING OCEAN BOTTOM SEISMOGRAPH

K.R. Peal and R.E. Kirk

The paper describes an all-digital ocean bottom seismograph which can record data only when an event is detected. The system is in fact an intelligent data logger which can record data on a predetermined time programmed schedule or when an event of a desired character occurs. Considerable flexibility in defining events is provided by software algorithms which are applied by a dedicated microprocessor.

Major portions of the system are described in detail. The sensor complement at present consists of a three-axis seismometer, a hydrophone, and a compass. The signal conditioning consists of preamplifiers, anti-aliasing filters, a signal multiplexer, and a gain-ranging amplifier. Dynamic range is over 100 dB.

A large buffer store is used so that data are collected when the tape recorder is not active thus eliminating a significant source of system noise. Use of a large buffer also allows large blocks of data to be examined for events.

A second microprocessor is used to control the system and to provide extra capabilities such as in situ calibration, accurate data time tagging, and computer-assisted preparation and checkout.

Published in: IEEE Proceedings of Third Working Symposium on Oceanographic Data Systems, IEEE CH 1926-5/83/0000/0114.

Supported by: Institute of Oceanographic Sciences, Wormley, Godalming, Surrey, UK.

A DEEP OCEAN AMBIENT NOISE SPECTRA RECORDING INSTRUMENT

K. Prada, D. Koelsch, K. Von der Heydt, E. Mellinger, S. Liberatore and F. Schuler

An instrument has been developed for measuring and recording ambient noise spectra in the deep ocean. Two microprocessors are used, one to perform general instrument control and a second for arithmetic opera-

tions. Noise data in the 0-200 Hz range are sampled continuously, transformed into the frequency domain, averaged, filtered and recorded.

Among the unique design features are 2048 point complex FFT at faster than real-time data rates, bubble memory recording for 512k bytes of processed data, tenth order elliptic anti-alias filter with 77db attenuation over a 25 Hz range, low power 14-bit A/D converter, interactive monitor, and external communications using the SAIL protocol.

The instrument is designed for a minimum 30-day life. Bubble memory storage and batteries in a separate case mean that the instrument need only be opened for repair.

Submitted to: Proceedings of the Third Working Symposium on Oceanographic Data Systems, IEEE Computer Society, 129-134, October, 1983.

Supported by: Western Electric Purchase Order #DNV-326708.

FREE DRIFTING RELAYS BUOY SYSTEM

R.G. Walden and H.O. Berteaux

A description follows of a free drifting buoy system (RELAYS) capable of receiving acoustic signals from SOFAR floats and of measuring oceanic parameters from the surface down to the depth of the SOFAR sound channel (1100 m). Data acquired are processed on board the buoy and transmitted for distribution via the ARGOS system using two NOAA polar orbiting satellites. System analysis, component design, and system evaluation are presented with emphasis on the mechanical aspects of the system.

Published in: Proceedings 1983 Symposium on Buoy Technology, New Orleans, 118-124, April, 1983.

Supported by: NASA Grant NAGW-265; NASA Grant NAG-5-281.

WHOI Contribution No. 5353.

HEBBLE SEA DUCT ANNUAL PROJECT REPORT #3 YEAR TWO, 1982-1983

Clifford L. Winget

Design and fabrication in several critical areas, namely the flume duct work, test section, raw power, and recirculating pump, have been given "first priority" and are now part of a high acceleration design and fabrication schedule. This portion of

the program was originally the responsibility of the NASA/JPL team, unfortunately, budget difficulties prevented the group from completing their portion of the project.

Supported by: ONR Contract
N00014-82-C-0019.

OCEAN ACOUSTICS

FLUCTUATIONS OF SOUND PROPAGATING VERTICALLY THROUGH THE OCEAN

Yves Desaubies

Fluctuations of sound propagating vertically through the ocean is considered. Expressions are derived for the acoustic phase structure function and spectra. As an application, the errors incurred by systems based on travel time measurements (depth sounding and tracking systems) are estimated. They are found to be generally negligible.

Published in: Journal of the
Acoustical Society of America, 74,
1245-1249, 1983.

Supported by: ONR Contract
N00014-82-C-0019.

WHOI Contribution No. 5306.

SHALLOW WATER WAVEGUIDE CHARACTERIZATION USING THE HANKEL TRANSFORM

George V. Frisk and James F. Lynch

A technique for acoustically characterizing shallow water waveguides is presented. For a horizontally stratified ocean and bottom, the method consists of measuring the magnitude and phase versus range of the pressure field due to a cw point source and numerically Hankel transforming this data to obtain the depth-dependent Green's function versus horizontal wavenumber. It is shown that, in the context of normal mode theory, the Green's function contains information about the nature of the discrete and continuous modal spectra as well as the plane-wave reflection coefficients of the waveguide boundaries. Inversions are performed using pressure field data generated synthetically over reasonable experimental apertures (1-5 km) to obtain Green's functions for the cases of an isovelocity water column overlying both hard and fast isovelocity bottoms (Pekeris waveguide). The Green's function results show excellent agreement

with theory, although the subsequently calculated reflection coefficient of the bottom is of substantially lower quality. It is shown that features of the Green's function itself can be used to extract modal properties and characteristics of the bottom. The effects of sediment attenuation and shear in the Pekeris case are discussed, and a comparison of this method with conventional phased array mode resolution techniques is made.

Accepted: Journal of the
Acoustical Society of America.

Supported by: ONR Contract
N00014-82-C-0152; ONR Contract
N00014-83-K-0594.

WHOI Contribution No. 5534.

ON THE USE OF FOCUSED HORIZONTAL ARRAYS AS MODE SEPARATION AND SOURCE LOCATION DEVICES IN OCEAN ACOUSTICS PART I: THEORY

James F. Lynch

It is shown that one can perform mode separation and source location using a focused horizontal array by utilizing the fact that the system response is greatest to a source located at or near the focal point. This mode separation/source location scheme is discussed for four different physical situations: (1) a motionless point cw source in a range invariant environment, (2) a motionless point cw source in a range varying environment, (3) a motionless point broadband source in a range invariant environment, and (4) a moving cw source in a range invariant waveguide environment. Two types of (horizontal) arrays are treated, the line array and the convex circularly curved array.

Published in: The Journal of the
Acoustical Society of America,
74(5), November, 1983.

Supported by: ONR Contract
N00014-77-C-0196.

WHOI Contribution No. 5316.

SOME SIMPLE EXPRESSIONS FOR THE BEAMFORMING PROPERTIES OF FOCUSED HIGH RESOLUTION CIRCULAR ARRAYS, WITH APPLICATIONS TO REFOCUSING SYSTEMS

James F. Lynch

Some simple analytical expressions for two important optical characteristics of focused high resolution circular arrays are

derived, specifically those for beam patterns and depth of field. The expressions generated, in addition to providing simple estimational tools for array design purposes, also lend themselves to straightforward interpretation in terms of the Fourier (farfield) and Fresnel (nearfield) kernels. Applications to refocusing arrays are discussed, as well as similarities and differences between such (sonar oriented) systems and radar and optical systems.

Published in: Journal of the
Acoustical Society of America,
74(3), September, 1983.

Supported by: ONR Contract
N00014-77-C-0196.

WHOI Contribution No. 5281.

A HYBRID NUMERICAL/ANALYTIC TECHNIQUE
FOR THE COMPUTATION OF WAVE FIELDS
IN STRATIFIED MEDIA BASED ON THE
HANKEL TRANSFORM

Douglas R. Mook, George V. Frisk
and Alan V. Oppenheim

A hybrid numerical/analytical technique is presented for computing the field due to a monochromatic point source in a horizontal medium. It is based on the exact Hankel transform relationship between the field in the range domain and the associated depth-dependent Green's function in the horizontal stratified wavenumber domain. The method uses a numerical evaluation of the Hankel transform. It is shown that a major source of error in such an evaluation arises from undersampling of the Green's functions at points where it becomes infinite. This error is described in terms of aliasing, analogous to the aliasing that has been well described for the discrete Fourier transform. It is shown that the error can be substantially reduced by removing the infinities, calculating the Hankel transform of the remaining portion of the Green's function numerically, and adding to it the analytically computed Hankel transform of the infinities. The sum of the analytic terms and the remaining Hankel transform always exactly equals the true field with no errors introduced other than those associated with the numerical evaluation of the Hankel transform, and the method is accurate in both the near and far field regions. The technique is developed in detail for the acoustics problem of a monochromatic point source and receiver in an isovelocity fluid half-space overlying a horizontally stratified fluid medium. It is found that under circumstances of interest in ocean bottom acoustics, where the Green's function has only a few singularities along the real horizontal wavenumber

axis, the technique is efficient and extremely accurate.

Accepted: Journal of the
Acoustical Society of America.

Supported by: ONR Contract
N00014-82-C-0152 at WHOI; Advanced
Research Projects Agency, monitored
by ONR under Contract N00014-81-K-
0742-NR-049-506; NSF Grant ECS80-
07102, Sanders Associates, Inc. at
MIT

WHOI Contribution No. 5453.

MEASUREMENTS OF GULF STREAM
MEANDERING AND EVIDENCE OF SEASONAL
THERMOCLINE DEVELOPMENT USING LONG RANGE
ACOUSTIC TRANSMISSION

J.L. Spiesberger, T.G. Birdsall,
K. Metzger, R.R. Knox, C.W. Spofford
and R.C. Spindel

Phase-coded signals with 60 ms resolution were transmitted from acoustic sources at ≈ 2000 m depth in the Sargasso Sea to three bottom-mounted receivers designated as West, East, and North stations at ranges between ≈ 1000 and ≈ 2000 km. The transmission paths to West and East stations were entirely in the Sargasso Sea. The path to North station crossed the Gulf Stream and so traversed one of the most time and range-dependent environments found anywhere in the ocean. Arrivals at all three stations were stable and could be identified from range dependent ray traces. Travel times at West station clearly change in response to the warming of the seasonal thermocline from Spring to Summer. The travel time change agrees with predictions. Travel time changes at North station primarily respond to the north-south meandering of the Gulf Stream.

Published in: Journal of Physical
Oceanography, 13, 1836-1846, 1983.

Supported by: ONR Contract
N00014-79-C-0071; NSF Grant
OCE80-17309.

WHOI Contribution No. 5434.

PERTURBATIONS IN TRAVEL TIME
AND RAY GEOMETRY DUE TO MESOSCALE
DISTURBANCES: A COMPARISON OF EXACT AND
APPROXIMATE CALCULATIONS

John L. Spiesberger and Peter F. Worcester

Exact calculations of the changes in eigenray travel time and geometry, due to range-dependent mesoscale disturbances, are

compared to three approximations. In the first approximation, the sound-speed perturbation is integrated along the unperturbed path. In the second approximation, the range average of the sound-speed perturbation is integrated along the unperturbed path. The third approximation uses the range-average raytrace to estimate changes of travel time and geometry. Calculations made both for a measured section through a mesoscale eddy and for a simplified model of an eddy (with variable strength, position, and boundary width) suggest that (i) while none of the approximations tested is uniformly the best, the first approximation is to be preferred; (ii) the eddy position relative to the ray path is quite significant; and (iii) the exact travel-time perturbations are not sensitive to whether the eddy boundaries are assumed sharp or gradual. For an eddy perturbation of ~ 9 m/s ($\sim 1.9^\circ\text{C}$ or Rossby wave displacement of about ± 200 m) at 700 m depth, the travel-time approximations are typically in error by 10%-20%. The associated change in turning depth from the unperturbed state to either the perturbed or the range-average state is typically ~ 100 -200 m. These errors are significant, but not large, and suggest that it will be beneficial to iterate linear inverse schemes used to estimate sound-speed perturbations from travel-time perturbations.

Published in: Journal of the
Acoustical Society, 74(1), 219-225,
1983.

Supported by: Scripps Institution of
Oceanography, University of
California at San Diego, La Jolla,
CA 92093.

EDDIES AND ACOUSTICS

R.C. Spindel and Y.J.F. Desaubies

This paper summarizes analytic and experimental work regarding the effects of eddies on acoustic propagation.

The ocean is a fluid waveguide through which sound propagates according to the scalar wave equation

$$D^2 p = c^2 A^2 p \quad (1)$$

In this form p is the acoustic pressure perturbation, c is the speed of sound, and the operator

$$D = \frac{\partial}{\partial t} + \underline{u} \cdot \nabla$$

represents the material derivative in the presence of a large scale (eddy) velocity field $\underline{u}$. In general c and $\underline{u}$ are functions of three spatial coordinates and time which

determine, with appropriate boundary conditions, the propagating acoustic field. The ocean surface acts nearly as a reflector scattering almost all of the sound energy in space and dispersing it in frequency. The ocean bottom has highly variable acoustic properties that can span the range from an almost perfect reflector to an almost perfect attenuator. Sea floor roughness and volume variability of c and $\underline{u}$ can also cause significant scattering. Although the sound speed usually varies only weakly with range, in the case of propagation through ocean eddies it is the eddy associated horizontal sound speed variations that are most significant in altering the pattern of acoustic energy distribution. In the case of warm or cold core rings horizontal variations in c can be quite large.

Published in: Eddies in Marine
Science. A.R. Robinson, ed.
Springer-Verlag Berlin Heidelberg.

Supported by: ONR Contract
N00014-82-C-0019; ONR Contract
N00014-82-C-0152.

WHOI Contribution No. 4795.

SUBMERSIBLES

DSRV ALVIN WEIGHT AND STABILITY

Arnold G. Sharp

Weight and stability calculations are presented for the research submersible ALVIN for the year 1983. Computations were done for each major ALVIN subassembly as well as for the entire vehicle in the normal surface and normal submerged conditions. Also considered were several emergency conditions where major submersible components may be jettisoned. In late 1982, ALVIN began a period of refit in preparation for operations aboard the newly assigned mother ship ATLANTIS II. The refurbished ship will have a stern-mounted A-frame, and launch and recovery of ALVIN will be by nylon rope using a single point attachment. As a result, ALVIN's structural frame has required modification to provide the lift attachment and additional strength members. The present calculations, and the accompanying results from the tethered trim dive and inclining experiments, reflect these most recent changes to the submersible's structure.

WHOI Technical Report.

Supported by: ONR Contract
N00014-82-C-0019; NSF Grant
OCE80-21920.

DEPARTMENT OF PHYSICAL OCEANOGRAPHY
Nicholas P. Fofonoff, Department Chairman

STRUCTURE OF THE CHANGJIANG RIVER PLUME
IN THE EAST CHINA SEA DURING JUNE 1980

Robert C. Beardsley, Richard Limeburner,
Dun Xin Hu, Ken Tang Le, Glenn A. Cannon
and David J. Pashinski

Hydrographic and current meter data collected in the East China Sea during June, 1980, has been used to describe the spatial and temporal structure of the Changjiang River discharge over the inner and mid-continental shelf off Eastern China. The fresh water discharge near the mouth of the Changjiang River exhibited a bimodal distribution, with the freshest water extending in a band to the south along the coast, and a relatively shallow, low salinity plume-like structure extending offshore on average towards the northeast. T/S analysis of the hydrographic data collected in this study indicates that the water structure in the East China Sea during early summer is dependent on three major influences: advection of warm, saline Taiwan Warm Current water from the south, advection of cooler, less saline Yellow Sea Cold Water from the north, and the fresh-water discharge of the Changjiang River. Based on the numerical model results of Choi (Korean Ocean Research and Development Institute Report 80-02, 72 pp., 1980), we predict that tidal mixing should produce well mixed zones along the central Chinese coast between 29°30' and 34°30'N. However, the buoyancy flux of the Changjiang River is locally larger than the amount of mixing due to tidal dissipation so that a stratified surface exists over the inner shelf. CTD and current profile data indicate that local dissipation is sufficient to keep a lower layer beneath the pycnocline vertically well-mixed over much of the 1980 study area.

CTD: conductivity, temperature, depth
T/S: temperature/salinity

In press: The Proceedings of the International Symposium on the Continental Shelf with Special Reference to the East China Sea, April 12-16, 1983, Hangzhou, China.

Supported by: NOAA Cooperative Agreement NA81AA-H-00008 and NSF Grant OCE80-14941.

WHOI Contribution No. 5328.

Kenneth H. Brink, Bruce A. Magnell
and Marlene A. Noble

An attempt is made to summarize the present observational and theoretical understanding of low-frequency velocity and bottom-pressure fluctuations over and around Georges Bank. We define the low-frequency band as the range of periods from about two to thirty days. Generally, the low-frequency current ellipses are oriented such that the strongest flow component tends to parallel the isobaths. Current fluctuations tend to be weakest over the Bank top and stronger near the Bank edges. Although alongbank currents are often coherent with each other along a given flank of the Bank, bank-wide coherence in currents is generally lacking. In contrast, bottom-pressure fluctuations are extremely coherent over the entire Gulf of Maine-Georges Bank region, and are clearly driven by local meteorological forcing. Wind-driven currents over the Bank appear to be related to this regional pressure response. Current and pressure variability is much greater in winter than summer, qualitatively in accord with seasonal changes in wind variability.

The few available theoretical models of the region deal primarily with wind-driving of current and pressure variability. Apparently, the regional models, which treat Georges Bank in its larger geographical setting, will prove the most successful to accounting for forced motions.

In press: Georges Bank, R. H. Backus, ed. The MIT Press.

Supported by: NSF Grant OCE82-00126;
the U.S. Bureau of Land Management;
and the U.S. Geological Survey.

WHOI Contribution No. 5456.

OBSERVATIONS OF THE COASTAL UPWELLING
REGION NEAR 34°30'N OFF CALIFORNIA:
SPRING 1981.

Kenneth H. Brink, David W. Stuart
and John C. Van Leer

Coordinated meteorological and oceanographic (CTD) measurements were made near Point Conception, California, during March-April 1981. The goal of the observations was to study coastal upwelling and the local characteristics of the assumed wind driving. Results showed substantial topographically-induced spatial structure in the near-surface winds, with weaker winds generally occurring within the Santa Barbara Channel. The 1981 "Spring transi-

tion" event was monitored via hydrographic and sea level measurements. The details of the event suggest that it was not entirely driven by local wind stress. The mean sea surface temperature pattern suggests the existence of an upwelling center between Points Arguello and Conception. The individual sea surface temperature charts are all dominated by patchiness on a 5-15 km scale. The nature of these structures is not well understood, but on the one occasion where a patch was isolated by a CTD survey, its structure penetrated to at least 50 dbar.

CTD: conductivity-temperature-depth sensor

In press: Journal of Physical Oceanography.

Supported by: NSF Grants OCE80-21028 and OCE80-21029.

WHOI Contribution No. 5319.

LONG-TERM OBSERVATIONS ON THE SOUTHERN FLANK OF GEORGES BANK: SEASONAL CYCLE OF CURRENT, TEMPERATURE, STRATIFICATION AND WIND STRESS

Bradford Butman and Robert C. Beardsley

Nearly continuous current measurements at 45 and 75 m were made from May 1975 to March 1979 at 40°51'N, 67°24'W on the southern flank of Georges Bank in water 85 m deep. Measurements at 15 and 84 m were made less often. The mean flow at 45 and 75 m was southwestward at approximately 8.6 and 3.7 cm/s, respectively. At 45 m, an average seasonal change of approximately 6 cm/s was observed in the monthly mean flow; strongest southwestward flow was in early September and weakest flow was in early March. At 75 m there was no significant seasonal change in the monthly mean flow. Hydrographic observations suggest that all of the mean shear and all of the seasonal variation between 45 and 75 m can be attributed to the seasonal change in the cross-bank density field. The current observations on the southern flank and additional measurements made at other locations around the perimeter of Georges Bank suggest that although a monthly mean subsurface clockwise circulation around the bank exists throughout the year, the flow was strongest in late summer and early fall and that recirculation around Georges Bank may be most likely in late summer. The flow was weakest and most variable in winter.

Supported by: the U.S. Geological Survey; the U.S. Bureau of Land Management; and NSF Grant OCE80-14941.

WHOI Contribution No. 5411.

THE PHYSICAL OCEANOGRAPHY OF GEORGES BANK: INTRODUCTION AND SUMMARY

Bradford Butman and Robert C. Beardsley

Extensive new direct current and temperature observations have been made between 1975 and 1980 on Georges Bank and the adjacent continental shelf. Preliminary analysis of these new observations have focused on documenting the seasonal mean circulation, the low-frequency current variability, and the tidal currents. The new measurements confirm the presence of a clockwise circulation around the Bank, similar to that described previously by Bigelow and Bumpus for the surface and bottom currents and provide important new information on flow variability. The clockwise flow exists throughout the year, is strongest in late summer and early fall and weakest and most variable in winter. Preliminary analysis of the current observations shows that subtidal current fluctuations are primarily aligned parallel to the local isobaths and are weaker in the Georges Bank region than in the adjacent Middle Atlantic Bight. The semidiurnal tidal currents dominate the currents in the Georges Bank region, especially over the shallow crest of the bank, and partially control the hydrography through vertical mixing, and may also partially drive the observed mean around-bank flow. There remain, however, many unanswered questions concerning the physical oceanography of Georges Bank and how local and regional physical processes influence circulation, mixing and productivity.

In press: Georges Bank. R. H. Backus, ed. The MIT Press.

Supported by: U.S. Bureau of Land Management; U.S. Geological Survey; and NSF Grant OCE80-14941.

WHOI Contribution No. 5504.

THE SEASONAL MEAN CIRCULATION ON GEORGES BANK: OBSERVATION AND THEORY

Bradford Butman, John W. Loder and Robert C. Beardsley

An initial description of the seasonal mean circulation around Georges Bank is presented based upon recent observations. Simplified models are used to examine qualitatively the extent to which various forcing mechanisms can explain the observed summertime and wintertime mean circulation. Rectification of the M_2 tidal currents contributes to the clockwise flow around the Bank in all seasons and can account for most of the along-bank flow on the northern flank in winter and as much as 50% of that observed on the southern flank. Inside the

shelf/slope front on the southern flank the forcing mechanism of the dominant year-round component of the southwestward flow is unclear: the leading candidates are along-bank bottom density and surface elevation gradients. During the summer the formation of tidal fronts around the shallow central Bank can explain much of the increased flow on the northern and southern flanks and should reinforce the tendency due to tidal rectification for recirculation of flow on the eastern side of Great South Channel. Cross-isobath currents are difficult to resolve in the presence of energetic tidal and low frequency motions. This may be a partial explanation for the relatively poor agreement between theoretical predictions of cross-isobath flow and the observations.

In press: Georges Bank. R. H. Backus, ed. The MIT Press.

Supported by: U.S. Geological Survey; Bedford Institute of Oceanography; Coastal Research Center, WHOI; and NSF Grant OCE82-00126.

WHOI Contribution No. 5539.

AN UPPER BOUND FOR THE TIDALLY RECTIFIED CURRENT AT ONE LOCATION ON THE SOUTHERN FLANK OF GEORGES BANK

Bradford Butman, Marlene A. Noble, David C. Chapman and Robert C. Beardsley

Long-term current observations at 45 and 75 m at one location on the southern flank of Georges Bank in water 85 m deep were examined for evidence of tidal rectification. Loder has shown analytically that rectification of the strong semidiurnal tidal current can cause a mean along-bank flow and thus may partially drive the observed clockwise circulation around Georges Bank. The amplitude of the tidally rectified along-bank flow is proportional to the squared amplitude of the cross-bank tidal current. A simple extension of Loder's model to include the weaker N_2 and S_2 tidal components suggests that fortnightly (354 h) and monthly (661 h) variations of the square of the cross-bank tidal current should cause a modulation of the subtidal along-bank flow. The predicted ratio (R) of the fortnightly and monthly modulation of the along-bank flow to the mean along-bank flow on the southern flank was a function of position and ranged from ~ 0.1 – 0.5 . The amplitude of modulation of the along-bank flow at 360 and 648 h, estimated from the (weak) coherence between the observed along-bank flow and the subtidal envelope of a simulated surface tide, was less than ~ 1.1 and 0.9 cm s^{-1} , respectively, at 45 m. The amplitude of the mod-

ulation which can be attributed to tidal rectification may be in error by the astrophysically forced M_m and MS_f tidal currents, which are undescribed in this region. However, the magnitude of the mean along-bank tidally rectified current determined from the observed modulation and R predicted by the analytical model was $\sim 2.0 \text{ cm s}^{-1}$ at 45 m (36% of the observed mean current in winter) and less than 1.6 cm s^{-1} at 75 m (43% of the observed mean current). Although R may change in a more realistic model, this analysis suggests that only part of the seasonal-mean along-bank flow on the southern flank of Georges Bank may be caused by tidal rectification.

Published in: Journal of Physical Oceanography, 13, 1452–1460, 1983.

Supported by: NSF Grant OCE80-14941 and a WHOI postdoctoral fellowship.

WHOI Contribution No. 5366.

DISPERSAL BY RANDOMLY VARYING CURRENTS

Gabriel T. Csanady

The long-term oceanic dispersal of persistent contaminants is approached as a problem in turbulent diffusion, with tidal, wind-driven, and other variable currents relegated to turbulence. The mean advection velocity in this problem is typically small compared with the r.m.s. fluctuation. Therefore, close to a continuous concentrated source, puffs of contaminant of all ages are present and have significant effects. Old puffs, i.e. those released a long time previously, give rise to a background concentration field. Young puffs affect the local contaminant concentration p.d.f. according to the probability of their presence, quantified by the visitation frequency.

The behaviour of young puffs is governed by variable advection and may be described approximately in terms of probability distributions obtainable from current-meter data. The visitation frequency can be calculated from the distribution of escape probability density, a Lagrangian equivalent of flux. A long-term effect of variable advection is the distribution of the contaminant over an 'extended' source, which serves as a starting point for the random walk of old puffs. The conventional approach of using the diffusion equation to describe this random walk is therefore valid as a description of the near-source background concentration field, provided that the extended source is used in place of the physical source.

A sample calculation for a typical open coastal case shows that the background con-

centration plume becomes wide compared with source dimensions, of order K_H/U , with K_H horizontal (eddy and shear) diffusivity, U mean advection velocity. The near-source value of the background concentration is correspondingly low, of order m/hK_n , with m the mass-release rate and h the water depth. Visitation frequencies calculated with the aid of current statistics drop rapidly with distance from the source, especially in the cross-shore direction. The typical cross-shore diameter of the extended source region is a few kilometres.

r.m.s.: root mean square

p.d.f.: probability distribution function

Published in: Journal of Fluid Mechanics, 132, 375-394, 1983.

Supported by: DoE Contract DE-AC02-79 EV10005 and Sherman Fairchild Fund, California Institute of Technology.

WHOI Contribution No. 5310.

THE "INSULATING" EFFECT OF A STEEP CONTINENTAL SLOPE

Gabriel T. Csanady and Ping Tung Shaw

Geochemical evidence shows that the mid-continental slope off the Mid-Atlantic Bight constitutes an important sink for substances attached to fine particles. It is reasonable to infer that near-bottom currents in this region are generally much weaker than those over the adjacent continental shelf, allowing the settling out of fine particles and their incorporation into permanent sediment. A partial dynamical explanation of this phenomenon may be sought in the "insulating" effect of a steep slope. At levels below the main thermocline the bottom pressure perturbation associated with steady flow is governed by a parabolic equation which expresses vorticity tendency balance between bottom stress curl and vortex stretching. A heat conduction analogy is appropriate, in which the friction coefficient corresponds to conductivity and the bottom slope corresponds to heat capacity. Over the continental slope the latter is large, suppressing pressure gradients originating from the shelf. Thus steady, near-bottom currents over the middle slope are generally weak. Exceptions to this rule are locations subject to intense winter cooling and deep convection, which may generate strong thermohaline near-bottom flow. Oscillatory motions of importance (which may be taken to be coastally trapped waves and topographic waves propagating shoreward from deep water) are weak partly because an efficient local generation mechanism is

lacking. Disturbances originating from the Gulf Stream decay as the depth reduces below 3 km, by a mechanism the details of which are not clear at present. The findings in the Mid-Atlantic Bight are likely to apply to other continental slopes distant from western boundary currents and not subject to intense winter cooling.

Published in: Journal of Geophysical Research, 88, 7519-7524, 1983.

Supported by: DoE Contract DE-AC02-79 EV10005.

WHOI Contribution No. 5379.

THE HYDROGRAPHIC STRUCTURE AND VARIABILITY OF GEORGES BANK

Charles N. Flagg

A review of the hydrographic structure and variability of the Georges Bank region is presented based upon observations spanning almost a century. The Georges Bank/Gulf of Maine system is extremely complex with as many as nine distinct water masses found within a 300 km radius. The bank topography which prevents easy entry of slope water into the Gulf plus strong fronts and the seasonal thermocline are primarily responsible for the simultaneous existence of the various water masses. Interannual fluctuations are significant in the region with annual mean temperature variations of order $\pm 1^\circ\text{C}$. The annual means also exhibit changes of the same order which last a decade or longer. The important hydrographic structures of the Bank are: the shelf-water/slope-water front which is also found along the northeastern U.S. coast; a region of cold near-bottom water referred to as the cold band; a tidal mixing front surrounding the shallow well-mixed central Bank, especially during the spring and summer; and a subsurface front along the northern edge of the Bank associated with a seasonally fluctuating jet. The offshore sloping shelf-water/slope-water front is found along the 100 meter isobath ± 10 km, migrating onshore from winter through summer with an offshore adjustment in fall. The transport of mass and momentum through this front is unknown although crude estimates of water exchange exist for the effect of warm core rings and diffusive processes. The cold band is formed from remnant winter shelf water during spring when the seasonal thermocline isolates the near-bottom water from vernal warming. This remnant water is replaced after 30 to 60 days by water from intermediate depths in the Gulf of Maine, surface water from the Bank and Gulf, and deep water from the Northeast Channel. Temperature/salinity

correlations and seasonally-averaged cross-isobath transects of temperature, salinity, sigma-t, and dissolved oxygen based upon data taken during the intensively studied period 1975-1979 are presented to provide quantitative estimates of the physical property distributions.

In press: Georges Bank. R. H. Backus, ed. The MIT Press.

Supported by: U.S. Bureau of Land Management; DoE Contract DE-AC02-79 EV10005; and the Coastal Research Center, WHOI.

WHOI Contribution No. 5461.

WATER MASSES AND CIRCULATION IN THE EAST CHINA SEA

Richard Limeburner and Robert C. Beardsley

The Changjiang (Yangtze) River mixes with two main water masses in the East China Sea: the Yellow Sea Cold Water and the Taiwan Current Warm Water. In summer during high runoff, the river discharge flows to the south in a turbid coastal boundary region and to the northeast as a relatively shallow, intermediate salinity, low suspended sediment plume. In winter during low runoff, the river discharge flows only to the south along the coast and Yellow Sea Cold Water intrudes into the East China Sea. Remote sensing of the Yellow and East China Seas shows the large scale structure of the turbid coastal boundary region.

In press: The Proceedings of the International Symposium on the Continental Shelf with Special Reference to the East China Sea, April 12-16, 1983, Hangzhou, China.

Supported by: NOAA Cooperative Agreement NA81AA-H-00008 and NSF Grant OCE80-14941.

WHOI Contribution No. 5327.

I N S T R U M E N T A T I O N

POPUP: A PROTOTYPE BOTTOM-MOORED LONG-TERM CURRENT PROFILER

Arthur D. Voorhis and Albert M. Bradley

A self-contained, bottom mounted instrument for measuring high resolution vertical profiles of horizontal current in the deep ocean over long time periods (up to a year) has been designed, constructed, and

successfully tested. It employs an interferometric technique to track small, expendable, buoyant probes containing c.w. acoustic beacons which are released on a preset schedule. A carrying capacity of 60 probes is quite feasible. Tracking data is stored on cassette tape and is recovered with the bottom instrument. The design of the instrument is discussed and errors analyzed. The uncertainty in the horizontal current averaged over 10 m vertical intervals is approximately 2 cm s^{-1} and is due primarily to acoustic phase noise from reflection off the rough sea surface. Considerable improvement in future design can be expected.

Supported by: NSF Grants OCE79-21786 and OCE82-00024.

WHOI Contribution No. 5569.

L A B O R A T O R Y M O D E L S

LABORATORY MODELS OF BAY-TYPE CONTINENTAL SHELVES IN THE WINTER

Takashige Sugimoto and
John A. Whitehead, Jr.

A laboratory experiment consisting of a shallow sea of constant depth bounded by a deep ocean through a uniformly sloping continental rise was conducted. The experiment is cooled from above and there is a region that exhibits sinking convection cells, which form the coldest water. This water then spills off the right-hand side of the shallow sea as a density current (looking downstream for counterclockwise rotation) and forms bottom water in the deep experimental ocean. There are three regimes. For very slow (scaled) rotations there is one big gyre in the shallow sea and the sinking (convection) region is near the coast. For faster rotations there are two or three gyres, and for even faster rotation there are many gyres. In this latter case the sinking regions are very patchy and sporadic. Comparison with theoretical estimates of temperature difference between the coldest shelf water and the offshore water, based upon the assumption that the density current removes all the cold water, was good but yielded somewhat low predictions.

Published in: Journal of Physical Oceanography, 13, 1819-1828, 1983.

Supported by: Ministry of Education, Science and Culture, Japan and NSF Grant OCE80-18322.

WHOI Contribution No. 5336.

DISLOCATIONS IN CONVECTION
AND THE ONSET OF CHAOS

John A. Whitehead

High Prandtl number convection possesses a square flow pattern that is steady and is apparently stable to infinitesimal disturbances. This pattern is unstable to finite-amplitude disturbances, however, because a more chaotic (in time and space) spoke pattern of convection eats its way into the squares from the lateral boundaries. Experiments are described in which the breakup of the squares is initiated by dislocating one square in the middle of the apparatus with the use of a small, heated resistor. Once a critical heating rate and time is exceeded, the dislocation initiates a spoke cell which then systematically destroys neighboring square cells, resulting in the more chaotic spoke pattern. If the critical rate is not exceeded, the cell becomes severely deformed during the heating, but will relax back to a square convection cell after heating has ceased.

Published in: Physics of Fluids, 26,
2899-2904, 1983.

Supported by: NSF Grant OCE80-18322.

WHOI Contribution No. 5429.

A LABORATORY STUDY OF GYRES AND UPLIFT
NEAR THE STRAIT OF GIBRALTAR

John A. Whitehead, Jr.

The Stommel, Bryden and Mangelsdorf idea -- that water from great depth in the Mediterranean may be drawn up and out through the Strait of Gibraltar through selective withdrawal -- is tested in the laboratory. The gyre corresponding to the Alboran Sea gyre is found to play an important role in the selective withdrawal process. A gyre is also observed in the Atlantic side, produced by turbulent suction of the plume of Mediterranean water plunging down into the Atlantic.

Supported by: ONR Contract N00014-82-C-
C-0019, NR 083-004.

WHOI Contribution No. 5596.

LARGE SCALE OCEANIC
HEAT AND WATER
PROPERTIES

WARM WATER TO COLD WATER CONVERSION
IN THE NORTHERN NORTH ATLANTIC OCEAN

Michael S. McCartney and Lynne D. Talley

A three box model of warm water to cold water conversion in the northern North Atlantic is developed and used to estimate conversion rates, given specified water mass temperatures, conversion paths and rate of air-sea heat exchange. Each box is required to satisfy heat and mass balance statements. In the Norwegian Sea box, warm water enters from the south, is cooled in the cyclonic gyre of the Norwegian-Greenland Sea. Cold water returns southwards to the open subpolar North Atlantic. Some exchange also is included with the North Polar Sea. The open subpolar North Atlantic has two boxes. In the abyssal box, the dense overflows from the Norwegian Sea flow south, entraining warm water from the upper ocean box. In the upper ocean box, warm water enters from the south, and circulates cyclonically, culminating in the production of Labrador Sea Water. Branches of warm water feed the Norwegian Sea box, and downward entrainment feeds the abyssal box. Our estimates are that $14 \times 10^6 \text{ m}^3/\text{sec}$ of warm (11.5°C) water flows north and to the west of Ireland, with about a third of this branching into the Norwegian Sea. The production rate for Labrador Sea Water is $8 \frac{1}{2} \times 10^6 \text{ m}^3/\text{sec}$, and this combines with a flow of dense Norwegian Sea Overflow waters (with entrained warmer waters) at $2 \frac{1}{2} \times 10^6 \text{ m}^3/\text{sec}$ to give a Deep Western Boundary current of $11 \times 10^6 \text{ m}^3/\text{sec}$. The total southward flow east of Newfoundland is this plus $4 \times 10^6 \text{ m}^3/\text{sec}$ of cold light Labrador Current waters (there is a net southward flow between Newfoundland and Ireland of about $1 \times 10^6 \text{ m}^3/\text{sec}$ of ultimate origin in the North Pacific, that enters the Atlantic from the north via the North Polar Sea).

In press: Journal of Physical
Oceanography.

Supported by: ONR Contracts N00014-74-
C-0262, NR 083-004 and N00014-79-C-
0071, NR 083-004; and NSF Grant
OCE78-22223.

WHOI Contribution No. 5410.

THE STRUCTURE, DYNAMICS AND ORIGIN OF
A SMALL SCALE LENS OF WATER IN
THE WESTERN NORTH ATLANTIC THERMOCLINE

Stephen C. Riser, W. Brechner Owens,
H. Thomas Rossby and Curtis C. Ebbesmeyer

We report in detail on a small scale, isolated lens of water observed in the western North Atlantic thermocline during the POLYMODE Local Dynamics Experiment. The lens was first revealed by anomalous anticyclonic motion in the trajectory of a SOFAR float at 750 m. Using a combination of float, hydrographic, nutrient, and moored current and temperature data, we deduce that the lens was about 20 km across with a thickness not greater than 300 m, centered on 750 m; that its shape was not inconsistent with that of a vertical-radial Gaussian eddy, that at its center the lens had strong anomalies of salinity, dissolved oxygen, nitrate, and potential vorticity; and that these properties are consistent with an origin for the feature at least several thousand kilometers from where it was observed in the western North Atlantic, although an unambiguous origin cannot be discerned from the data. If such phenomena are not uncommon, they may be an important mechanism for large-scale mixing of water properties in the ocean.

SOFAR: sound fixing and ranging

Supported by: NSF Grants OCE82-14541,
OCE79-01087, OCE77-19403 and
OCE78-18662.

WHOI Contribution No. 5518.

O C E A N C I R C U L A T I O N

LIMITING PROCESSES THAT DETERMINE
BASIC FEATURES OF THE CIRCULATION
IN THE MEDITERRANEAN SEA

Harry L. Bryden and Henry M. Stommel

Large mixing in the Mediterranean Sea, driven in part by the large-scale buoyancy loss due to evaporation, forces the salinity difference between Atlantic and Mediterranean waters to a minimum and the magnitude of the two-layer flow through the Strait of Gibraltar to a maximum consistent with the internal hydraulic control at the Strait. Predicted values of the salinity difference and inflow and outflow through the Strait agree reasonably well with observed values and are independent of the mixing as long as it is large enough. This asymptotic limit has been called overmixing. It describes the basic features of

the circulation of Atlantic and Levantine intermediate waters in the upper 1000 m of the western Mediterranean. For the deeper circulation, the outflow velocity of the intermediate water over the Gibraltar sill provides a Bernoulli head capable of sucking deeper and denser water up and over the sill. Water which is too dense or too deep to be easily sucked over the sill is defined as deep water. A grazing streamline analysis for the interface between intermediate and deep waters and a model of deep water formation due to late-winter buoyancy loss events allow the depth of the interface and the density contrast across it to be computed. This asymptotic limit predicts characteristics of the deep water circulation in reasonable agreement with those observed.

Supported by: ONR Contract N00014-82-C-0019, NR 083-004 and NSF Grant OCE80-15789.

WHOI Contribution No. 5405.

CURRENT VARIABILITY
NEAR THE SOUTHEAST NEWFOUNDLAND RIDGE

Nick P. Fofonoff and Ross M. Hendry

A cooperative moored array experiment to measure currents and temperatures in the vicinity of the Southeast Newfoundland Ridge was carried out over a 14-month period starting September, 1979 and terminating in December, 1980. Measurements were obtained from an incoherent exploratory array of nine moorings instrumented at 500, 800, 1500 and 4000 m levels. The objectives were to explore the amplitude, spatial variability and time scales of the eddy field in the decay region of the Gulf Stream. An additional experiment and mooring to examine the influx of Norwegian Sea Overflow Water into the western North Atlantic Basin was imbedded in the array. Supplementary CTD and XBT profiles and a single current profile were taken during deployment and recovery cruises to aid in interpretation of the moored measurements and identification of water masses.

A description of the array experiment and preliminary results are reported. Mean velocities were weak compared with rms eddy velocities and were not determined with statistical significance at most locations. Maximum daily-averaged speeds ranged from 0.3 to 0.6 m/s at 500 m and 0.09 to 0.3 m/s at 4000 m. Eddy kinetic energies at 500 m ranged from 0.034 J/kg near the Ridge to 0.005 J/kg in the southeastern corner of the array. Deep energies were reduced by a factor of 4 to 12 with the 4000-m level showing bottom intensification at most locations. Topographic effects were noted in

the orientation of eddy velocities and reduction of low-frequency components near the Ridge. The eddy field showed no clear-cut evidence of wave-like propagation or advection in streamfunction plots. An increase in time scales is evident north and east of the Ridge towards the Newfoundland Basin and the Mid-Atlantic Ridge, where most of the eddy kinetic energy at 500 m is at time scales of 150-200 days compared with 100 days southwest of the Ridge and about 50-60 days in the central POLYMODE array at 55°W.

CTD: conductivity-temperature-depth sensor
rms: root mean square
XBT: expendable bathythermograph

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5593.

EVIDENCE FOR BAROCLINIC INSTABILITY IN THE GULF STREAM RECIRCULATION

Nelson G. Hogg

Long term current measurements from POLYMODE Array 2 are analyzed using the technique of empirical orthogonal functions applied in the frequency domain. The region of especially good instrument coverage and clearest decomposition into empirical modes is also in the Gulf Stream recirculation (~ 36°N, 55°W). On theoretical grounds, this is expected to be an area actively unstable to baroclinic disturbances.

Both the horizontal and vertical structure of motions with periods ranging from 10 to 120 days are studied. Spectra reveal that kinetic energy is peaked at 30 day periods with a weaker peak at 12 days. The most energetic empirical mode accounts for about 50 percent of the measured energy although somewhat less than this at higher subinertial frequencies. Consistent with the theoretical calculations of Gill, Green and Simmons it is found that the zonal phase propagation is nearly independent of frequency and averages about 12 cm/sec toward the west. The 30 and 12 day energy peaks correspond to wavenumbers (wavelengths) of $-.018 \text{ km}^{-1}$ (350 km) and $-.033 \text{ km}^{-1}$ (190 km) respectively. These are almost identical to the predicted scales for secondary and primary growth rate maxima for baroclinically unstable disturbances to a zonal current which is horizontally uniform but whose strength decreases exponentially with depth. Energy in these motions increases toward the west at a rate which is statistically indistinguishable from zero but also from the predicted rate.

The vertical structure of these motions is also qualitatively similar to the theo-

retical modes. Phase for velocity decreases with depth, whereas temperature phase increases leading to energy release concentrated in the thermocline near the steering level which is projected to be somewhere above the shallowest observational level (600 m). The 30 day wave is much less depth dependent in amplitude than the 12 day wave and is responsible for the intense deep eddy field observed in this area by Schmitz.

At this point in the recirculation the instability process is still active and the resulting motions have not yet reached a finite amplitude state.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5384.

THE DYNAMICS OF THE OCEANIC GENERAL CIRCULATION: THE POTENTIAL VORTICITY FIELD

William R. Holland, Thomas Keffer
and Peter B. Rhines

The general circulation of the oceans is determined by the field of potential vorticity, q , and the fields of temperature and salinity. q is conserved on fluid particles, in absence of external forcing or dissipation. A combination of memory of sea-surface boundary conditions and subsequent mixing act to determine the history of q , as fluid descends from the surface to ventilate the ocean interior. Predictions of q from theory and numerical simulations are here compared with oceanic observations. The North Atlantic and North Pacific fields strongly support theoretical models in which geostrophic contours deform to follow the circulation, homogenization occurs in gyres, and injected 'mode' waters visibly affect the q -field. The intense thermohaline circulation in the mid-depth North Atlantic is reflected in the maps of potential vorticity. By contrast, the North Pacific is weakly ventilated at depth, and the potential vorticity comes to a remarkably uniform value over its entire extent, north of the tropics.

In press: Nature.

Supported by: NSF Grant OCE83-19780;
Center for Analysis of Marine
Systems, WHOI; and NSF to the
National Center for Atmospheric
Research.

WHOI Contribution No. 5492.

THE VENTILATION OF THE WORLD'S OCEANS:
MAPS OF THE POTENTIAL VORTICITY FIELD

Thomas Keffer

Maps of potential vorticity (q) within four different density layers are presented for the North and South Atlantic, North and South Pacific, and Indian Oceans. Potential vorticity is evaluated as $(f/\rho)\partial\rho/\partial z$, a prescription that is valid for large slow scales and that allows its mapping from hydrographic data alone. Here, f is the Coriolis frequency, ρ the potential density, and z the vertical coordinate.

It is shown that the character of the subtropical thermocline will vary greatly depending on the extent to which it is "ventilated". That is, whether fluid particles tend to make many circuits around the gyre before having their potential vorticity reset (unventilated) or whether the flow is more "open-ended" or "ventilated" where particles have their value of q set by surface processes and then are subsequently injected into the thermocline (via Ekman pumping).

The North Atlantic thermocline is shown to be highly anomalous. The extensive outcropping of its density surfaces and the strong downward Ekman pumping south of the subpolar front allow the q contours to run up into the outcrop window, in the manner of Luyten, Pedlosky and Stommel (J. Phys. Oceanogr., 13, 292-309, 1983). However, the other oceans show a much greater tendency towards homogenization, even at great depth, suggesting that ventilation is relatively weak. Only in the very shallowest layers, where fluid particles are exposed to direct atmospheric forcing, is there a tendency for the outcrop properties to propagate into the interior. In these regions, the wintertime stratification is advected into the interior as tongues of high q .

Supported by: ONR Contract N00014-82-C-0019, NR 083-004 and the Center for Analysis of Marine Systems, WHOI.

WHOI Contribution No. 5538.

THE DENSITY JUMP ACROSS LITTLE BAHAMA BANK

James R. Luyten and Henry M. Stommel

There is a discontinuity in density across the Bahama Banks. On the western side of the Banks, beneath the mixed layer, isopycnals are about 100 meters shallower than on the eastern side. An explanation is offered in terms of a reduced gravity layered model of an inertial western boundary current forced to flow through a channel (the Florida Straits) which is narrower than its open ocean scale-width.

In press: Journal of Geophysical Research.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400 and NSF Grant OCE80-15789.

WHOI Contribution No. 5369.

UPSTREAM EFFECTS OF THE GULF STREAM ON THE
STRUCTURE OF THE MID-OCEAN THERMOCLINE

James R. Luyten and Henry M. Stommel

Attempts to resolve a paradoxical result by Blandford (Tellus, XVII, 69-76, 1965) led us to reexamine the degree of specification necessary to assure continuity of solution in a density stratified non-rotating channel through which two layers move over a deep resting one. We then proceeded to formulate and solve, numerically, a rudimentary beta-plane model of an inertial boundary current fed entirely from the equator: the zero potential vorticity case. Inspection of the numerical results indicates a simpler self similar form of the evolution of this problem, leading to same results but explicitly analytical. This model implies that the Gulf Stream exerts a control over mid-ocean thermocline structure.

In press: Progress in Oceanography.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400 and NSF Grant OCE80-15789.

WHOI Contribution No. 5341

A SYNOPTIC AND STATISTICAL DESCRIPTION
OF THE GULF STREAM AND SUBTROPICAL GYRE
USING SOFAR FLOATS

W. Brechner Owens

Trajectories of seventeen SOFAR floats ballasted to nominal depths of 700 and 2000 m and tracked for fifteen months have been analyzed to produce synoptic and statistical descriptions of the Gulf Stream and subtropical gyre. SOFAR floats launched in and near the Gulf Stream along with positions of the surface Gulf Stream and satellite tracked buoys demonstrate strong baroclinic shear. Nevertheless, the horizontal patterns of the Gulf Stream meanders and Gulf Stream rings are the same at the surface and at 700 and 2000 m depth. Above the thermocline, particle speeds greater than the propagation speed of the meanders cause parcels of water to move with the flow pattern. At greater depths the velocities are at most equal to the propagation speed, and the deep trajectories are free to pass through the Gulf Stream or ring.

Strong temporal variability is found near the tail of the Grand Banks. Instances of both flow southward into a narrow Gulf Stream recirculation and northward into the North Atlantic Current can be seen.

Statistical averages along 55°W are consistent with other measurements at that longitude. Eddy kinetic energy varies by over two orders of magnitude from the Gulf Stream to mid-gyre. A mean Gulf Stream, flowing eastward at all depths, can be seen. In the interior, the zonal velocity has a short meridional scale with bands of flow in either direction.

SO FAR: sound fixing and ranging

In press: Journal of Physical
Oceanography.

Supported by: NSF Grants OCE78-18007
and OCE81-09145.

WHOI Contribution No. 5394.

DRIFTING DERELICTS IN THE NORTH ATLANTIC 1883-1902

Philip L. Richardson

In December 1883 the U.S. Navy Hydrographic Office began to publish monthly Pilot Charts. These showed the positions and drifts of abandoned derelict sailing vessels and other dangers to navigation in the North Atlantic. During the years 1887-1893 a total of 1628 derelict sightings were made. The repeated sightings of 200 derelicts identified by name gave the first large-scale and long-term measurement of ocean trajectories. A new chart of these trajectories between 1883 and 1902 is presented showing features of the general ocean circulation and early evidence of the variability of ocean currents.

In press: Progress in Oceanography.

Supported by: NSF Grant OCE81-09145.

WHOI Contribution No. 5385.

AVERAGE SEASONAL VARIATION OF THE ATLANTIC NORTH EQUATORIAL COUNTERCURRENT FROM SHIP DRIFT DATA

Philip L. Richardson and Theresa K. McKee

Surface currents in the tropical Atlantic were studied using historical ship drift data. These data are the only available ones capable of resolving the long term seasonal fluctuations of currents over a broad geographical region. The North Equatorial Countercurrent was found to extend

as a continuous eastward current across the Atlantic in the band 5-8°N from July to December. The maximum average eastward velocity was about 30 cm/sec during July-September. From January to June the countercurrent disappears and westward currents are observed. Average westward velocity reaches 20 cm/sec in April and May centered near 40°W. The annual cycle in surface velocity agrees closely with that seen in geostrophic velocity inferred from variations in thermocline depth across the countercurrent as determined recently by Garzoli and Katz. A semi-annual variation of zonal currents along the equator was also observed.

Supported by: NSF Grant OCE82-08744
and the Centre National pour
l'Exploitation des Océans, France.

WHOI Contribution No. 5583.

OBSERVATIONS OF HORIZONTAL VELOCITIES AND VERTICAL DISPLACEMENTS IN THE EQUATORIAL PACIFIC OCEAN ASSOCIATED WITH THE EARLY STAGES OF THE 1982/83 EL NIÑO

John M. Toole and Mark D. Borges

Near equatorial absolute velocity data and CTD/O₂ observations collected from the R/V CONRAD in the fall of 1982 are presented. This cruise marked the first opportunity to sample the eastern third of the equatorial Pacific after the onset of the 1982/83 El Niño. The data may be characterized as exhibiting abnormal eastward flow in the near surface water and large depressions of the thermocline. These changes are quantified by comparing with previously collected data and discussed in terms of simple equatorially trapped wave dynamics.

CTD: conductivity-temperature-depth sensor

Supported by: NOAA Contract NA82RAC-
00149; NSF Grant OCE83-08415; and a
WHOI Summer Student Fellowship.

WHOI Contribution No. 5571.

WIND FORCED VARIABILITY IN THE CENTRAL EQUATORIAL PACIFIC DURING PEQUOD

Arthur D. Voorhis, James R. Luyten,
Gerald J. Needell and John H. Thomson

The variability of zonal current, dynamic height, and density in the central equatorial Pacific, computed from vertical profiler measurements, is discussed for three time periods: January 1981, February 1982, and April 1982. The increased

undercurrent in the last period, accompanied by a raising of sea level and a deepening of the upper ocean pycnocline, was due to a gravest mode Kelvin wave pulse propagating eastward at a speed of about 300 cm/sec. Evidence suggests that such pulses are forced by tropical storms in the western Pacific and have reoccurred every spring since 1979, at least.

Supported by: NSF Grant OCE79-21786.

WHOI Contribution No. 5352.

SOME PRELIMINARY RESULTS CONCERNING DEEP NORTHERN-BOUNDARY CURRENTS IN THE NORTH PACIFIC

Bruce A. Warren and W. Brechner Owens

Fourteen-month velocity records from five moorings along Long. 175°W, in combination with CTD and nutrient data, distinguish two pronounced, quite steady, deep zonal currents near the northern margin of the North Pacific. One flows westward at speeds of 1-3 cm s⁻¹ along the slope of the Aleutian Island Arc and appears to be the northern-boundary current required by deep circulation theory. The other flows eastward at about 1-2 cm s⁻¹ just to the south above the Aleutian Trench and upper Aleutian Rise and carries water that is slightly lower in temperature and dissolved silica and slightly higher in salinity and dissolved oxygen than that to the north or south. We believe that this eastward jet represents an extreme distortion of a level-bottom, interior flow pattern, as forced by the meridional variation in bottom slope on the southern flank of the Aleutian Rise, which imposes a varying effective "beta" on the system. A simple model of this effect is discussed.

In press: Progress in Oceanography.

Supported by: ONR Contracts N00014-79-C-0071, NR 083-004 and N00014-82-C-0019, NR 083-004.

WHOI Contribution No. 5484.

R I N G S, E D D I E S

THE ENERGY BALANCE DURING THE POLYMODE LOCAL DYNAMICS EXPERIMENT INTENSIVE CTD SURVEYS

Nancy A. Bray, W. Brechner Owens
and Bach-Lien Hua

Seven high resolution CTD surveys conducted during the POLYMODE Local Dynamics

Experiment, together with estimates of the velocity field from moored current meters, are used to calculate the thermocline energy balance. Energy is partitioned into that contained in the mean slope of isopycnals observed during the surveys and that contained in deviations of the density field from the mean. The conservation equations for the available potential energy (APE) and kinetic energy (KE) contained in both the mean and eddy fields are then examined.

It is found that eddy APE increases through conversion from mean APE and decreases at about the same rate through conversion to eddy KE. Those conversions are consistent with baroclinic instability of the large scale density field. Energy for the conversion of mean APE to eddy APE derives principally from changes in the local storage of mean APE, rather than from advection of mean APE into the region. Advective fluxes are generally found to be less important than conversion terms.

Supported by: NSF Grant OCE77-19403
and Mellon Foundation Fellowship.

WHOI Contribution No. 5580.

COMPARISON OF EDDIES OFF THE NORTH BRAZILIAN AND SOMALI COASTS

John G. Bruce

Repeated temperature sections using XBTs from ships of opportunity along the sea lane off the Brazilian coast indicate that a region of mesoscale eddies occurs in which the eddies are particularly well developed in northern summer and fall. These are located between approximately 3°N-10°N, are on the order of 200 to 400 km across, and may extend to 300 to 400 m in depth. Relatively fresh surface water, apparently from the region of the Amazon outflow, is often observed in the eddy region. Similar eddies, both in size and latitude, are found off the Somali coast. These are associated with the turning offshore of all or part of the strong north flowing Somali current. It is speculated that a somewhat similar circulation may also occur in northern summer in the Pacific Ocean.

XBT: expendable bathythermograph

In press: Journal of Physical
Oceanography.

Supported by: NSF Grant OCE79-25350.

WHOI Contribution No. 5329.

ENERGETIC EDDIES AT THE NORTHERN EDGE
OF THE ANTARCTIC CIRCUMPOLAR CURRENT
IN THE SOUTHWEST PACIFIC

Harry L. Bryden and Ron A. Heath

Two-year time series measurements of current and temperature were made at the northern edge of the Antarctic Circumpolar Current in the southwestern Pacific from 1978 to 1980 in order to investigate the role of eddy heat and momentum fluxes in the heat and momentum budgets of the circumpolar region. The measurement site near $49^{\circ}30'S$, $170^{\circ}W$ is a transition zone or front between subantarctic and subtropical waters in which meridional temperature and salinity gradients and, hence, vertical shear in eastward velocity extend down to 2000 m depth. Energetic eddies are found to have typical amplitudes of 20 cm s^{-1} at 1000 m depth, to be vertically coherent throughout the water column, to vary over temporal scales of 30 days and horizontal scales of 60 km and to propagate southeastward at about 12 cm s^{-1} . Observed eddy kinetic energy of $169\text{ cm}^2\text{s}^{-2}$ at 1000 m depth is larger than that at comparable depths in the Gulf Stream and Kuroshio extension currents and three times larger than that in central Drake Passage. Like Drake Passage eddies, these eddies transport heat poleward in the band of periods between 20 and 40 days, but this band does not dominate the spectrum the way it does in Drake Passage. Energetic lower frequency fluctuations effect an eddy heat flux whose sign is ambiguous so that the net meridional heat flux is not statistically significant. Observed eddy heat flux, however, is directed poleward and is large enough to balance the heat lost to the atmosphere by antarctic waters south of the polar front. Eddy momentum fluxes transport eastward momentum northward away from the Circumpolar Current and, hence, help to balance the eastward momentum put into the water column by the wind in the circumpolar region. Although the eddy momentum fluxes are statistically significant, they are only about one fourth of the size required to balance the wind input. Since this region was anticipated to be one of three zones in the circumpolar region where eddy momentum fluxes are especially large, it appears that eddy momentum fluxes due to temporal fluctuations are not a significant balancing mechanism in the zonal momentum budget for the circumpolar ocean. A preliminary estimate of the standing eddy momentum flux due to large spatial scale variations in the Circumpolar Current suggests that it may be large enough to balance the wind input.

Supported by: NSF Grant OCE77-22887 and the New Zealand Department of Scientific and Industrial Research.

WHOI Contribution No. 5455.

VELOCITY AND HYDROGRAPHIC STRUCTURE OF
A GULF STREAM WARM CORE RING

Terrence M. Joyce

The hydrographic and velocity structure of Gulf Stream Warm Core Ring 81D are investigated using a deep CTD- O_2 section through the ring together with horizontal currents at 100 m depth obtained with an acoustic-Doppler system. The pycnocline within the ring is depressed 500 m below that of the surrounding Slope Water and maximum currents at 100 m approach 2 m/s at a radius of 70-80 km from ring center. Water mass analysis reveals that the waters in the central core of the ring are similar to the Sargasso Sea and distinct from the Slope Water to depths of 1300 m. Near the surface, the central core is rotating as a solid body approximately once every 3.6 days. The azimuthal currents at 100 m were combined with the gradient current relationship to obtain absolute dynamic topography of the ring. Sea surface in ring center is 70 dynamic centimeters above that in the Slope Water, and ring currents decrease quickly with increasing depth to undetectable levels below 2000 m. The ring is less dense than the Slope Water below 100 m, but actually denser near the surface, resulting in a weak, subsurface velocity maximum. Oxygen utilization at 100 m is apparently higher in the ring than in the Sargasso Sea and there is also a suggestion of a weak, barotropic cyclone 120 km to the northeast of the center of 81D.

CTD: conductivity-temperature-depth sensor

In press: Journal of Physical
Oceanography.

Supported by: NSF Grant OCE80-16983
and NASA Grant NAGW-272.

WHOI Contribution No. 5530.

RAPID EVOLUTION OF
A GULF STREAM WARM-CORE RING

Terrence M. Joyce, Richard H. Backus,
Karen S. Baker, Patricia L. Blackwelder,
Otis B. Brown, Timothy J. Cowles,
Robert H. Evans, Greta A. Fryxell,
David G. Mountain, Donald B. Olson,
Ronald J. Schlitz, Raymond W. Schmitt,
Peter C. Smith, Raymond C. Smith
and Peter H. Wiebe

Gulf Stream warm-core rings form in the Slope Water between the North American continental shelf and the Gulf Stream by the separation of a north-extending meander. The initial physical, chemical and biological properties of the core of these 100-200

km eddies are often similar to those of its parent water mass, the Sargasso Sea. A clockwise rotating remnant of the Gulf Stream circulates around the core with surface current speeds of $50\text{--}200\text{ cm s}^{-1}$. While warm-core rings can slowly change over periods of months through interactions with the surrounding Slope Water, Gulf Stream meanders, the continental slope, other rings, and the atmosphere, we have discovered that major alterations to ring structure can occur during very short periods (two to five days) when an interaction with the Gulf Stream is particularly intense. Such short-period interactions between ring and Gulf Stream are a major factor governing a ring's evolution.

In press: Nature.

Supported by: NSF Grant OCE80-16983; NASA Grant NAGW-272; National Marine Fisheries Service; and Bedford Institute of Oceanography, Canada.

WHOI Contribution No. 5381.

AN UPPER OCEAN CURRENT JET AND INTERNAL WAVES IN A GULF STREAM WARM CORE RING

Terrence M. Joyce and Marvel C. Stalcup

On 22 June 1982, the R/V ENDEAVOR, while participating in a multi-ship study of a warm core ring 82B, encountered a strong front in the core of the ring. The vessel was headed on a radial section outward from ring center while a CTD was repeatedly raised and lowered between 10 and 300 meters. Current profiles in the upper 100 m were obtained continuously with a Doppler acoustic profiling system. Above the shallow 45 m seasonal thermocline, a current jet of 4 km width was encountered having a central core of relatively light water and a maximum current of 1.1 m/s. This jet was both highly non-linear and totally unexpected. A high frequency packet of directional internal waves was acoustically observed in the seasonal thermocline at the outer edge of the jet. Vertical velocities were large enough (6 cm/s) as to be directly observable in the Doppler returns. The waves were propagating from the northeast, parallel to the ship track, and orthogonal to the jet towards the center of the warm core ring. While a non-linear, centrifugal term was required for the force balance of the jet, the high-frequency internal wave packet could be explained with linear, gravest-mode wave dynamics.

CTD: conductivity-temperature-depth sensor

In press: Journal of Geophysical Research.

Supported by: NASA Grant NAGW-272 and NSF Grant OCE80-16983.

WHOI Contribution No. 5412.

THE SPATIAL AND TEMPORAL EVOLUTION OF A CLUSTER OF SOFAR FLOATS IN THE POLYMODE LOCAL DYNAMICS EXPERIMENT (LDE)

H. Thomas Rossby, James F. Price
and Douglas C. Webb

Forty neutrally buoyant floats (20 at 700 meters, 20 at 1300 meters) were used in the POLYMODE Local Dynamics Experiment (LDE) to provide a quasi-Lagrangian description of the structure and evolution of the mesoscale eddy field in a limited region characterized by higher kinetic energy levels than those obtained in MODE. This paper is an overview and summary of the data collected.

The temporal development of the two-level float array is presented in a sequence of maps, each spanning five days. In these one readily notices an "oscillation" of floats at 1300 meters along a line oriented northeast-southwestward before the cluster breaks apart. At 700 meters, the float cluster subdivides much more rapidly. The first group of 700 meter floats drifts to the west; the second group, launched two months later, goes far to the east.

Running cluster averages of the floats at 1300 meters reveal great sensitivity to the "oscillatory" velocity field while the array is "tight" or coherent; the corresponding 700 meter averages, although noisy, show clearly the westward and eastward motion of the first and second clusters, respectively. Grand averages for u, v (cm/s) and kinetic energy (ergs/gm) are $(-1.8, 0, 80)$ and $(-1.6, 0, 34)$ for the shallow and deep floats.

The spatial correlation functions show large scales of coherence. The zero crossing of the transverse velocity correlation function is about 100 km for both the 700 and 1300 meter floats compared to 55 km for the 1500 meter floats in MODE. The longitudinal correlation scale is also much larger.

MODE: Mid-Ocean Dynamics Experiment

Supported by: NSF Grants ID075-18930, OCE78-18662, OCE81-09145; and ONR Contract N00014-76-C-0226.

WHOI Contribution No. 5563.

ABYSSAL EDDY KINETIC ENERGY
IN THE NORTH ATLANTIC

William J. Schmitz, Jr.

The first map of the North Atlantic that contained estimates of the intensity of the abyssal eddy field is now several years old, and in the interim the relevant data base has increased roughly six-fold. Contemporary charts are presented, containing these more recent observations, along with some new styles of data presentation. The basic picture of maximum abyssal eddy kinetic energy near the fully developed Gulf Stream is consistently substantiated, along with a two order of magnitude latitudinal decay (from about 10^2 to roughly 1, in cm^2s^{-2}) into the interior of the subtropical gyre (west of the Mid-Atlantic Ridge). On the basis of data obtained in the last few years, eddy kinetic energy levels at abyssal depths near Cape Hatteras, in the vicinity of the Grand Banks, and east of the mid-Atlantic Ridge are down by 1-2 orders of magnitude from maxima near the Gulf Stream at intermediate longitudes in the western North Atlantic. These results lead to the first relatively clear-cut identification of the zonal scales of eddy kinetic energy variation near the Gulf Stream. The initial contour map of observed abyssal eddy kinetic energy is presented, albeit crude. A fundamental qualitative point is that both eddy and mean fields at abyssal depths have similar zonal and meridional scales of variation in the mid-latitude North Atlantic.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5407.

ABYSSAL EDDY KINETIC ENERGY LEVELS
IN THE WESTERN NORTH PACIFIC

William J. Schmitz, Jr.

An earlier observation of about $50 \text{ cm}^2\text{s}^{-2}$ for the maximum abyssal ($\sim 4000 \text{ m}$ depth) eddy kinetic energy (K_E) in the western North Pacific along 152°E , occurring in the vicinity of the Kuroshio Extension, is now supported by nearly two years of observation with moored instruments deployed from 28 to 41°N . The maximum occurred at the same site each year (nominal, from one array setting to the next), near 35°N , and the composite or two-year average is $45 \text{ cm}^2\text{s}^{-2}$. At analogous longitudes in the western North Atlantic one finds abyssal kinetic energies as high as $100 - 150 \text{ cm}^2\text{s}^{-2}$ in the vicinity of the Gulf Stream. However, the abyssal western North Pacific could be more energetic at longitudes other

than 152°E . Eddy intensities at abyssal depths near the Gulf Stream do drop off to smaller values approaching either Cape Hatteras or longitudes in the vicinity of the Grand Banks of Newfoundland.

Abyssal K_E values at specific sites along 152°E were observed to vary by as low as a few percent and as high as a factor of 2 from year to year. A prominent secondary maximum in K_E , primarily due to zonal variability, was observed to exist north of the Kuroshio Extension for one year, but not the next. However, the maximum K_E at 4000 m depth, its location, and the general shape of the latitudinal distribution of eddy intensity are relatively stable. Averages over all moorings where these data are available were 22.5 and $18 \text{ cm}^2\text{s}^{-2}$ for the two respective deployments. The mesoscale contribution (periods 30 to 150 days) to abyssal K_E along 152°E was typically 50 to 60 percent.

In press: Journal of Physical
Oceanography.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5409.

OBSERVATIONS OF THE VERTICAL STRUCTURE OF
THE EDDY FIELD IN THE KUROSHIO EXTENSION

William J. Schmitz, Jr.

Observations for 370 days at a site in the Kuroshio Extension along 152°E yield a unique eddy field. The lowest frequencies are strongly vertically coherent from 350 to 6000 m depths (nominal). Kinetic energy frequency distributions are independent of depth as a good first approximation. None of the other vertical sets of time series previously examined possess these characteristics to such a degree, coherence being typically lower across the thermocline, with spectral shape changing as well. The vertical distributions of eddy kinetic energy at this Kuroshio Extension site and for the Gulf Stream along 55°W are nearly identical, if a depth independent contribution of $100 \text{ cm}^2\text{s}^{-2}$ is added to the former.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5480.

THE GENERATION OF BAROTROPIC EDGE WAVES
BY DEEP-SEA INTERNAL WAVES

David C. Chapman

A simple two-layer, step-shelf model is used to demonstrate that barotropic (surface) edge waves of substantial amplitude can, in principle, be generated by deep-sea internal waves incident upon the coastal topography. Some qualitative features of the results suggest that this mechanism could account for the edge-wave "noise" observed by Munk, Snodgrass and Gilbert (J. Fluid Mech., 20, 529-554, 1964).

Supported by: NSF Grants OCE82-00126
and OCE80-14941.

WHOI Contribution No. 5579.

WARM WATER MASS FORMATION

Gabriel T. Csanady

Poleward heat transport by the ocean implies warm water mass formation, i.e. the retention by the tropical and subtropical ocean of some of its net radiant heat gain. Under what conditions net heat retention becomes comparable to latent heat transfer to the atmosphere depends on the relative efficiency of transfer processes across the air-sea interface, the top of the atmospheric mixed layer, and the floor of the oceanic mixed layer. A thermodynamic model of the interacting atmospheric and oceanic mixed layers, with the top of the atmospheric layer taken to be at cloud base, shows that net oceanic heat retention is significant under the following circumstances.

(1) Seasonal heat storage, amplitude of order 100 W m^{-2} . This is a fairly straightforward consequence of the large heat capacity of the oceanic mixed layer and leads the seasonal forcing by about a month.

(2) Massive upwelling (vertical velocity of order 10^{-5} m s^{-1}), mostly along equatorial cool tongues, with net heat retention of order 100 W m^{-2} . The upwelling cooler water is heated and transported away mainly by the divergence of surface layer flow (less by the increasing temperature in the direction of the flow).

(3) Cold water advection, mostly within the subtropical gyres, net heat retention of order 30 W m^{-2} . The latitudinal variation of radiant heating and generally equatorward surface flow in the northern portions of subtropical gyres leads to a moderate rate of warming of the water column as it

moves along; i.e., to net heat retention of the above order.

A comparison of model results with observation shows that, over the subtropical gyres, observed temperature and humidity relationships can only be simulated realistically if cold water advection is taken into account. In addition, it is necessary to suppose that the transfer coefficient at the top of the atmospheric mixed layer (at cloud base) is about as large as at the sea surface, while the transfer coefficient at the oceanic mixed layer floor is negligible, except in regions of massive upwelling. The general dominance of latent heat transfer arises from the large value of a non-dimensional latent heat coefficient (a material property) and from the rapid drop of saturation specific humidity with height in the atmosphere.

In press: Journal of Physical
Oceanography.

Supported by: NASA Grant NAG 5-185.

WHOI Contribution No. 5365.

THE NONLINEAR SPIN-UP OF A STRATIFIED OCEAN

William K. Dewar, Peter B. Rhines and
William R. Young

The adjustment of a nonlinear, quasi-geostrophic, stratified ocean to an impulsively applied wind stress is investigated. As in Rhines and Young (J. Fluid Mech., 122, 347-367, 1982) and Young and Rhines (J. Mar. Res., Suppl. to 40, 559-596, 1982; J. Mar. Res., 40, 849-972, 1982), barotropic advection of vortex tube length is the dominant modal interaction. The present study complements the steady state theories which have recently appeared, and extends earlier linear model results.

In terms of the Sverdrup transport, the equation for baroclinic evolution is a forced advection-diffusion equation. Solutions of this equation, subject to a "tilted disk" Ekman divergence, are obtained, analytically for the case of no diffusion and numerically otherwise. The similarity between the present equation and that of a forced barotropic fluid with bottom topography is shown.

Barotropic flow, which is assumed to mature instantly, can reverse the tendency for westward propagation and thus produce regions of closed geostrophic contours. Inside these regions, dissipation, or equivalently the eddy field, plays a central role. Here we assume that eddy mixing effects a lateral diffusion of potential vorticity; hence, within the closed characteristics our solution approaches a state of uniform potential vorticity. We demon-

strate that an acceptable steady state requires both nonlinearity and dissipation, but mention that the present steady state depends on our eddy parameterization.

The evolution of potential vorticity and the thermocline are critically examined, and it is shown that the adjustment time of the model is governed by dissipation, rather than baroclinic wave propagation as in linear theories. Therefore, spin-up of a nonlinear ocean may take several times that predicted by linear models, which agrees with analyses of eddy-resolving general circulation models.

Supported by: NSF Grant OCE80-23763.

WHOI Contribution No. 5421.

TRACER DISPERSAL BY MID-OCEAN MESOSCALE EDDIES. I: ENSEMBLE STATISTICS

Dale B. Haidvogel and Thomas Keffer

The rate at which, and the processes by which, a passive tracer is stirred and mixed in a turbulent mesoscale eddy field are examined for environmental parameters characteristic of a homogeneous mid-ocean region. The simulated, time-dependent eddy field is obtained by direct integration of the forced/damped barotropic vorticity equation. The dispersal of a spatially localized, instantaneous release of tracer (a "tracer spot") within the evolving velocity field is subsequently computed from the advective-diffusive equation. An ensemble of 10 independent releases is used to determine the average spreading properties of the tracer spot.

On an f -plane, the ensemble-averaged dispersal is approximately isotropic and is associated with an effective diffusion rate substantially greater than that supported in the absence of turbulent advection. Quantitatively, the effective diffusivity is shown to be $O(UL)$, where U and L are characteristic velocity and length scales of the turbulent flow. This estimate is consistent with the traditional mixing length hypothesis. With the addition of β , the simulated flow field has substantial zonal anisotropy. Ensemble-averaged dispersal of tracer spots is similarly anisotropic, and the overall rate of tracer dispersal is substantially reduced over its f -plane value.

Both with and without β , the initial rate at which maximum tracer concentration and total tracer variance (θ^2) decay are given by the approximate law $\exp[-\alpha\gamma t]$ where γ is the RMS rate of strain, and α is an $O(1)$ constant. The heightened rate of variance loss over that associated with pure (subgridscale) diffusion is shown to be accommodated by the rapid transfer of tracer

variance from the largest to the shortest scale tracer features, that is, by the rapid sharpening of tracer gradients by turbulent advection. A detailed examination of the dispersal of individual tracer realizations, and the associated question of tracer streakiness, will be addressed in part II of this paper.

RMS: root mean square

In press: Dynamics of Atmospheres and Oceans.

Supported by: NSF Grant OCE81-09486 and the Center for Analysis of Marine Systems, WHOI.

WHOI Contribution No. 5482.

A COMPARISON OF THE FORMULATION FOR EDDY DIFFUSION IN TWO ONE-DIMENSIONAL STRATIFICATION MODELS

Brian Henderson-Sellers, M. H. McCormick and Donald Scavia

Two models for thermal stratification based on turbulent diffusion concepts are analyzed and compared. The models by Henderson-Sellers and by McCormick and Scavia are shown to be equivalent at large values of the Richardson number, R_i . At small R_i , the simpler model reverts to specification of the turbulent diffusion as a constant value. This simplification is also demonstrated to be a realistic approximation only at low wind speeds and for deep lakes. By comparison of these model types, a (previously empirically defined by McCormick and Scavia) parameter β is related conceptually to the lake depth, H .

Published in: Applied Mathematical Modelling, 7, 212-215, 1983.

Supported by: DoE Contract DE-AC02-79 EV10005.

WHOI Contribution No. 5368.

EASTERN BOUNDARY VENTILATION AND THE STRUCTURE OF THE THERMOCLINE

Joseph Pedlosky

The effect of relaxing the zero geostrophic zonal flow condition at the eastern boundary on the structure of the ventilated thermocline is described. It is argued that the effect of dissipation can allow a vertical exchange of mass between layers at the eastern boundary. A model is studied analytically that assumes such an interchange while continuing to forbid a

net eastward mass flux at each latitude.

The consequences of the relaxation of this condition are striking. Each active layer in the ventilated thermocline can now have a non-zero thickness on the eastern wall. As a result, each such layer has a stagnant shadow zone near the eastern boundary. Each shadow zone extends across the gyre at low latitudes. Hence the isopycnal surfaces, while shallow, remain at finite depth as the equator is approached. The shadow zones are nested, i.e., the shadow zone of the deepest layer is laterally most extensive and the zone shrinks eastward and southward with decreasing depth. The vertical recirculation at the eastern boundary adds a new source of ventilation and a zone of motion to the structure of the ventilated thermocline described by Luyten, Pedlosky and Stommel (J. Phys. Oceanogr., 13, 292-309, 1983).

In press: Journal of Physical Oceanography.

Supported by: NSF Grant ATM79-21431.

WHOI Contribution No. 5430.

THE EQUATIONS FOR GEOSTROPHIC MOTION IN THE OCEAN

Joseph Pedlosky

A multiple-scale approach is used to develop the quasi-geostrophic dynamics for synoptic oceanic scales. This new approach allows the buoyancy frequency, N , and the Coriolis parameter f to be slowly varying functions of horizontal position. No series expansion of f about some arbitrary central altitude is required. Nor is there a limitation on the geographic extent of the synoptic scale domain.

Simultaneously, the equation for large scale geostrophic flow on gyre, or planetary scales is derived as a solvability or consistency condition on the synoptic scale dynamics. Thus a single derivation suffices to develop both sets of dynamical equations. In addition, the multiple-scale analysis explicitly describes the interaction terms between the synoptic scale and gyre scale fields of motion.

In press: Journal of Physical Oceanography.

Supported by: NSF Grant ATM79-21431.

WHOI Contribution No. 5442.

THE GROWTH AND DECAY OF FINITE-AMPLITUDE BAROCLINIC WAVES

Joseph Pedlosky

The finite-amplitude dynamics of an unstable baroclinic wave in an f -plane, two-layer model is examined in the case where the degree of Ekman friction is unequal in the two layers. In particular, the case is examined where one layer is free of frictional dissipation.

The method employed requires the calculations of the wave amplitude evolution on two long-time scales in order to describe the essential equilibration of the wave which at intermediate times is rendered unstable by asymmetric Ekman dissipation. The method of reconstitution is used to combine equations appropriate for each epoch of the wave's history to provide a single equation useful for calculation.

The main physical result is the emergence of a finite lifetime for the unstable wave. The wave grows, equilibrates and then decays leaving a wave-free state and an altered zonal flow with reduced potential energy. The speculation is offered that this process will be repetitive with the addition of a weak dissipative process to restore the wave-free state to its initial configuration.

Published in: Journal of the Atmospheric Sciences, 40, 1863-1876, 1983.

Supported by: NSF Grant ATM79-21431.

WHOI Contribution No. 5330.

ON THE RELATIVE IMPORTANCE OF VENTILATION AND MIXING OF POTENTIAL VORTICITY IN MID-OCEAN GYRES

Joseph Pedlosky

An a priori estimate is given of the effect of ventilation on the process of potential vorticity homogenization. Since the homogenization process depends on the presence of weak mixing, it is shown that only a small exposure to a zone of ventilation is required to lead to an arrest of the homogenization.

In press: Journal of Physical Oceanography.

Supported by: NSF Grant ATM79-21431.

WHOI Contribution No. 5428.

VENTILATION, POTENTIAL-VORTICITY
HOMOGENIZATION AND THE STRUCTURE
OF THE OCEAN CIRCULATION

Joseph Pedlosky and William R. Young

A model for the vertical structure of the oceanic circulation is presented which combines elements of the theory of the ventilated thermocline, given by Luyten, Pedlosky and Stommel (J. Phys. Oceanogr., 13, 292-309, 1983) with the theory of Rhines and Young (J. Mar. Res., Suppl. to 40, 559-596, 1982) for the wind driven circulation of an unventilated ocean. Our model consists of a ventilated thermocline region above an unventilated zone in which motion is limited to pools of constant potential vorticity. The model is non-linear and hence the presence of ventilation affects the dynamics of the unventilated motion and vice-versa. The planetary geostrophic equations are used and so the quasi-geostrophic assumption of Rhines and Young is relaxed, allowing large isopycnal excursions. It is shown that the presence of ventilation generally shrinks and weakens the size and vigor of the subsurface pools of homogenized potential vorticity. At the same time, within those domains, the strength of circulation in the ventilated zone is somewhat diminished as the subsurface layers carry a portion of the Sverdrup transport. We argue that the (mathematically) consistent circulation in the absence of sub-thermocline constant potential vorticity pools is unstable. The non-uniqueness of the non-dissipative Sverdrup dynamics is demonstrated by the ambiguity in the specification of potential vorticity in the deeper, unventilated layers. The study emphasizes the subtle importance of dissipation in selecting a unique solution.

In press: Journal of Physical
Oceanography.

Supported by: ONR Contract N00014-79-
C-0472 and NSF Grant ATM79-21431.

WHOI Contribution No. 5415.

T U R B U L E N C E A N D M I X I N G ,
F I N E S C A L E P R O C E S S E S

ESTIMATES OF THE RELATIVE ROLES OF
DIAPYCNAL, EPIPYCNAL AND DOUBLE-DIFFUSIVE
MIXING IN ANTARCTIC BOTTOM WATER
IN THE NORTH ATLANTIC

Trevor J. McDougall and John A. Whitehead

Whitehead and Worthington (J. Geophys.

Res., 87, 7903-7924, 1982) have measured the fluxes of heat and salt due to the northward flow of Antarctic Bottom Water through a passageway 300 km wide between the Ceara Rise and the Mid-Atlantic Ridge at about 4°N. Downstream of this "sill" the temperature and salinity of the underflowing water increase, and Whitehead and Worthington have described this change as being due to "downward" fluxes of heat and salt across isothermal surfaces.

We consider the relative roles of three separate mixing processes to these "downward" fluxes of heat and salt, and we use this information to decide between Whitehead and Worthington's two separate estimates of the volume transport of Antarctic Bottom Water, one based on current meter data, and the other based on geostrophic calculations.

The data derived using the geostrophic velocities implies very large diapycnal turbulent mixing in relation to both salt fingering and epipychnal (i.e. along isopycnal) mixing, and so we conclude that the current meter data are more realistic.

Supported by: ONR Contract N00014-82-C-
0019, NR 083-004.

WHOI Contribution No. 5581.

THE CHARACTERISTICS OF SALT FINGERS IN A
VARIETY OF FLUID SYSTEMS, INCLUDING STELLAR
INTERIORS, LIQUID METALS, OCEANS, AND MAGMAS

Raymond W. Schmitt

The growth rates, flux ratios, wavenumbers, and bandwidths of salt fingers are computed for Prandtl numbers from 10^{-7} to 10^4 and diffusivity ratios from 1 to 10^8 , using the exact similarity solutions of Schmitt (Deep-Sea Res., 26A, 23-40, 1979). This model successfully explains the variation in flux ratio in the heat/salt and salt/sugar systems and produces salt-finger spectra in agreement with ocean observations. The calculations presented here should be useful in the study of double diffusion in astrophysics, chemistry, geology, metallurgy, meteorology, and oceanography, and may be applicable to problems in the growth of semiconductor crystals.

Published in: Physics of Fluids, 26,
2373-2377, 1983.

Supported by: ONR Contract N00014-82-
C-0019, NR 083-004.

WHOI Contribution No. 5340.

FINESCALE VELOCITY-DENSITY CHARACTERISTICS
AND RICHARDSON NUMBER STATISTICS OF THE
EASTERN EQUATORIAL PACIFIC

John M. Toole and Stanley P. Hayes

An analysis of finescale horizontal velocity shear and density data collected along 110°W in the equatorial Pacific is presented. The measurements were made with the free-fall velocity-density profiler, TOPS. Twenty-five deployments are used to investigate the variability in the depth interval 150-900 m. In this interval the shear and strain fields are dominated by finescale structures. The extra equatorial latitude band 4-10°N exhibits shear and strain spectra as well as Richardson Number (Ri) statistics that are consistent with mid latitude internal wave model predictions. Approaching the equator an enhancement of shear and strain variance is found along with an accompanying increase in the occurrence of Ri less than 1/4. Consistent with previous studies, the present measurements suggest that the 13°C thermostat is turbulently mixed. A high occurrence of Ri less than 1/4 is also found below the thermostat and in the latitude range 2-4°S. The implications of these observations are discussed along with the representativeness of the present measurements.

TOPS: Total Ocean Profiler System

Supported by: NOAA Contract NA82RAC-00149 and ONR Contract N00014-82-C-0019, NR 083-004.

WHOI Contribution No. 5462.

U P P E R O C E A N , S U R F A C E
L A Y E R P R O C E S S E S
A N D M E T E O R O L O G Y

PRELIMINARY RESULTS FROM THE
LONG-TERM UPPER-OCEAN STUDY (LOTUS)

Melbourne G. Briscoe and Robert A. Weller

The Long-Term Upper-Ocean Study (LOTUS) is a two-year experiment at 34°N, 70°W, in deep water over the Hatteras Abyssal Plain, 180 nautical miles from major topography and the mean path of the Gulf Stream. The observations began in May 1983. The principal measurements are from an array of three subsurface moorings and one surface mooring carrying meteorological instrumentation as well as near-surface current meters. Some basic meteorological and engineering data are telemetered back via the

ARGOS system, but most of the data are self-recorded. Other observations in the program include XBT sections, CTD profiles, and cooperative efforts with other investigators to obtain satellite imagery and other in situ measurements.

The main scientific goals of the experiment are: to obtain a description of the response of the upper ocean to atmospheric forcing in a variety of situations, such as in the presence of Gulf Stream rings, near oceanic fronts, and under tropical storms, and even to try and describe the statistical response to the fluctuating forcing; to investigate the energy level of the internal wave field over several seasons and during the passage of various forcing events as mentioned above; and to obtain the complete profile of eddy kinetic energy.

We display here selected results from the first year of LOTUS, especially those that do not require the analysis of the complete array of measurements and supporting data. Of particular interest is the eddy kinetic energy profile showing a nearly constant energy in the top 500 m; the abyssal eddy energy is also about half the expected level for the site.

The high-frequency internal wave field at 300 to 500 m depths shows a seasonal variation (high in the winter, low in the early fall) that cannot be ascribed to changes in the hydrography; at greater depths the winter maximum is still visible. The overall variation at each depth is a factor of 2 or 3 smaller and larger than the mean energy at that depth.

Some of the energetic events in the near-inertial field are apparently unconnected to local wind events and may instead be related to the edge of strong mesoscale oceanic currents; perhaps they are being radiated during the process of geostrophic adjustment by the mesoscale features.

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5585.

INTERCOMPARISON OF AIRCRAFT AND SURFACE BUOY
METEOROLOGICAL DATA DURING CODE-1

Carl A. Friehe, Robert C. Beardsley,
Clinton D. Winant and Jerome P. Dean

Intercomparisons of meteorological data -- wind speed and direction, surface temperature and surface pressure -- were obtained for NCAR Queen Air overflights of four buoys during the CODE-1 experiment. The overflights were at a nominal altitude of 33 m. Wind and air temperature sensors were at 10 m on two National Data Buoy Office (NDBO) buoys and at 3.5 m on two Woods Hole Oceanographic Institution (WHOI) buoys.

The buoy wind speeds were adjusted to the aircraft altitude using diabatic flux-profile relations and bulk aerodynamic formulas to estimate the surface fluxes and stability. For the experimental period (22 April to 23 May, 1981) and location (northern coast of California), the atmospheric surface layer was generally stable, with the Monin-Obukhov length on average +500 m with large variability.

The results of the intercomparisons of the above variables were in general very good. Average differences (aircraft-buoy) and standard deviations were $+0.1 \text{ m s}^{-1}$ (± 1.8) for wind speed, 3.3 deg (± 11.2) for wind direction, $\pm 0.02^\circ\text{C}$ (± 1.7) for air temperature and $+0.8 \text{ mb}$ (± 1.0) for surface pressure. The aircraft downward-looking infrared radiometer indicated a surface temperature 1°C lower than the buoy hull (NDBO) and 1 m immersion (WHOI) sea temperature sensors.

NCAR: National Center for Atmospheric Research

Supported by: NSF Grants OCE80-14941, OCE80-14942 and OCE82-14949.

WHOI Contribution No. 5451.

DIURNAL CYCLING: OBSERVATIONS AND MODELS OF THE UPPER OCEAN RESPONSE TO DIURNAL HEATING, COOLING, AND WIND MIXING

James F. Price, Robert A. Weller and Robert Pinkel

Air/sea measurements made in the eastern North Pacific from R/P FLIP provide a new, detailed view of the upper ocean's diurnal cycle. The measurements were made during a 25 day period in May 1980 when the daily maximum heat flux was $320 \leq Q \leq 820 \text{ W m}^{-2}$, and the daily average wind stress, $0.02 \leq \bar{\tau} \leq 0.25 \text{ N m}^{-2}$. In response, the diurnal range of near surface temperature was observed to be $0.04 \leq \Delta T_s \leq 0.40^\circ\text{C}$, and the daily minimum trapping depth was $30 \leq D_T \leq 3 \text{ m}$. Scale analysis of a simple model gives that $\Delta T_s \sim Q^{3/2}/\bar{\tau}$, and $D_T \sim \bar{\tau}/Q^{1/2}$, are both roughly consistent with observations. In this data set, and probably generally, the day-to-day variation of the diurnal thermal cycle is attributable mainly to the variation of $\bar{\tau}$.

Heat and momentum appear to be mixed in a nearly similar way. Thus the diurnal thermal cycle is accompanied by a diurnal cycle in the wind stress-driven acceleration of the surface layer. The result is a diurnal jet in upper ocean velocity whose amplitude is typically $V \sim 0.1 \text{ m s}^{-1}$. Scale analysis gives that $V \sim Q^{1/2}$, roughly as observed. The diurnal jet reaches its maximum amplitude just before

sunset and then decays on account of vertical mixing and rotation of the jet into the wind.

As a consequence of diurnal cycling, the time-averaged velocity profile has a spiral shape reminiscent of the classical Ekman spiral. However, the parameter dependence of vertical shear is in some important ways just opposite that expected from the classical model.

FLIP: Floating Instrumented Platform

Supported by: ONR Contracts N00014-76-C-0197, NR 083-400 and N00014-75-C-0152.

WHOI Contribution No. 5427.

OBSERVATIONS OF THREE DIMENSIONAL FLOW IN THE SURFACE MIXED LAYER OF THE OCEAN

Robert A. Weller, Jerome P. Dean, James F. Price, Erika A. Francis, John Marra and David C. Boardman

For many years, somewhat regularly spaced, long, narrow surface slicks or rows of floating seaweed have been observed on the surface of the ocean and of lakes. Langmuir's drifter studies showed that the surface slicks formed in regions of convergent flow between counter rotating helical vortices in the near surface mixed layer. Recently, a study was begun to make measurements of and to assess the role of such organized three dimensional flows in the upper ocean. In a December 1982 pilot experiment, as the Research Platform FLIP drifted off the coast of California, instruments were lowered into the mixed layer to measure horizontal and vertical velocities, temperature, conductivity, and chlorophyll *a* concentration. On several occasions, as FLIP drifted relative to the surface layer, narrow regions of downward flow were encountered at nearly regular intervals; during those times, surface drift cards formed into long, narrow bands. Downwelling velocities were observed to be as high as 0.2 m s^{-1} , but such strong flow was not typical. Small temperature and chlorophyll anomalies were found in association with the downwelling regions.

FLIP: Floating Instrumented Platform

Supported by: ONR Contract N00014-76-C-0197, NR 083-400.

WHOI Contribution No. 5439.

TECHNICAL REPORTS

NANTUCKET SHOALS FLUX EXPERIMENT (NSFE79) PART 2: MOORED ARRAY DATA REPORT

Robert C. Beardsley, Carol A. Mills,
John A. Vermersch, Jr., Wendell S. Brown,
Neal R. Pettigrew, James D. Irish,
Steven R. Ramp, Ronald J. Schlitz
and Bradford Butman

The Nantucket Shoals Flux Experiment (NSFE79) was conducted across the continental shelf and upper slope south of Nantucket from March 1979 to April 1980 to measure the flow of shelf water from the Georges Bank/Gulf of Maine region into the Middle Atlantic Bight. Conceived as a cooperative field experiment involving the Northeast Fisheries Center, U.S. Geological Survey (Woods Hole), University of New Hampshire, and the Woods Hole Oceanographic Institution, the experiment contained two principal components: a moored array of current meter and bottom instrumentation deployed at six locations across the shelf and upper slope spanning a depth range from 46 m to 810 m, and a series of 27 hydrographic surveys made along or near the moored array line during the experiment. A basic description of the NSFE79 hydrographic data has been given in Part 1 by W. R. Wright (NOAA Technical Memorandum, NMFS-F/NEC-23, 1983). A description of the moored array components and the basic moored array data sets is presented here in Part 2.

Prepared for: NSF under Grants OCE78-19513, OCE80-14941 and OCE82-00126.

WHOI Technical Report 83-13.

MEAN EULERIAN SUBSURFACE CURRENTS MEASURED IN THE GULF OF MAINE AND ADJACENT SCOTIAN AND NEW ENGLAND SHELF AND SLOPE REGIONS, 1974-1980

Robert C. Beardsley, Carol A. Mills,
Peter C. Smith and Bradford Butman

The Bedford Institute of Oceanography, E.G.&G., National Marine Fisheries Service, U.S. Geological Survey and Woods Hole Oceanographic Institution have conducted separate moored array experiments during 1974-1980 to study various aspects of the regional circulation in the Gulf of Maine and adjacent Scotian and New England shelf and slope regions.

The mean currents and current variances measured in these experiments are summarized here in tabular and graphical form, together with other information about each experiment. While there have been few mea-

surements made in the interior of the Gulf of Maine, the map of mean subsurface currents demonstrates (a) a net flow of Scotian shelf water past Cape Sable into the Gulf, (b) a net inflow of slope water through the Northeast Channel into the Gulf, (c) a partially closed anticyclonic circulation around Georges Bank, and (d) a net outflow of shelf water south of Nantucket from the Gulf of Maine into the New England shelf.

Prepared for: U.S. Bureau of Land Management under Contract AA 551-MUO-18 and NSF under Grant OCE80-14941.

WHOI Technical Report 83-44.

HYDROGRAPHIC STATION DATA OF FIVE SURVEYS OF THE BETA-TRIANGLE IN THE EASTERN NORTH ATLANTIC, 1978-1981

David W. Behringer, George P. Knapp,
Robert J. Stanley and Henry M. Stommel

This is a data report, giving temperature, salinity and dissolved oxygen data obtained during five surveys of the beta-triangle area of the eastern North Atlantic in the years 1979-81. The report is divided into two parts. Part I consists of a listing of station data obtained on all stations occupied -- on the periphery of the triangle, inside it, and some nearby stations outside it. Part II is a set of profiles of this data drawn on the three sides of the triangle, for easy reference and visual comparison. Detailed scientific study of the data is published elsewhere.

Prepared for: NSF under Grant OCE80-15789 and NOAA.

WHOI Technical Report 83-24.

LONG-TERM UPPER-OCEAN STUDY (LOTUS) AT 34°N, 70°W: METEOROLOGICAL SENSORS, DATA, AND HEAT FLUXES FOR MAY-OCTOBER 1982 (LOTUS-3 AND LOTUS-4)

Clara Deser, Robert A. Weller
and Melbourne G. Briscoe

Meteorological data have been gathered from a moored surface buoy at 34°N, 70°W in the Long-Term Upper-Ocean Study (LOTUS) experiment. The meteorological results from the first year of LOTUS are encouraging; the data returned from redundant sensors agree closely. Surface heat fluxes calculated from the observations show the annual cycle of heat transfer to the mixed layer.

This report documents the meteorological sensors on the LOTUS-3 (May 1982-October 1982) and LOTUS-4 (November 1982-March 1983) surface buoys. It describes in detail the

telemetry of the meteorological data via the ARGOS satellite system. The measurements returned from LOTUS-3 are presented and evaluated. Monthly heat fluxes at the sea surface are computed using the bulk formulas and compared with the long-term means. The errors in the heat fluxes have been estimated.

Prepared for: ONR under Contract
N00014-76-C-0197, NR 083-400.

WHOI Technical Report 83-32.

A COMPILATION OF MOORED INSTRUMENT DATA
AND ASSOCIATED OCEANOGRAPHIC OBSERVATIONS
FROM THE VEMA CHANNEL, 1979 - 1981,
VOLUME XXXII

Ellen Levy

Summaries of long-term current and temperature measurements from moored instruments in the Vema Channel and a sample of hydrographic stations located in the area of these moorings are presented. Vema Channel is a deep trough in the Rio Grande Rise connecting the Argentine and Brazil Basins, in the vicinity of 30°S and 40°W. There were two settings in this experiment and the purpose was to study the flow of bottom water through the Vema Channel.

Tables, plots and statistics are presented for filtered and unfiltered time series.

Prepared for: NSF under Grant OCE78-
25405.

WHOI Technical Report 83-46.

A COMPILATION OF MOORED CURRENT METER DATA
FROM THE WESTERN NORTH PACIFIC,
VOLUME XXXI, 1980-1982

Ellen Levy and Susan A. Tarbell

Summaries of long-term current, temperature and pressure measurements from moored instruments in the western Pacific Ocean (along 152°E, between 28°N and 41°N) are presented. Tables, plots and statistics are presented for filtered and unfiltered time series. There were two consecutive settings of instruments, referred to as WesPac I and II, each spanning a one year interval (nominal). The objective of the experiment was to motivate exploratory descriptions of the low frequency variability in the western North Pacific.

Prepared for: ONR under Contract
N00014-76-C-0197, NR 083-400.

WHOI Technical Report 83-40.

CODE-1:
MOORED ARRAY AND LARGE-SCALE DATA REPORT

Leslie K. Rosenfeld, editor

The Coastal Ocean Dynamics Experiment (CODE) was undertaken to identify and study the important dynamical processes which govern the wind-driven motion of coastal water over the continental shelf. The initial effort in this multi-year, multi-institutional research program was to obtain high-quality data sets of all the relevant physical variables needed to construct accurate kinematic and dynamic descriptions of the response of shelf water to strong wind forcing in the 2 to 10 day band. A series of two small-scale, densely instrumented field experiments of approximately four months duration (called CODE-1 and CODE-2) were designed to explore and to determine the kinematics and momentum and heat balances of the local wind-driven flow over a region of the northern California shelf which is characterized by both relatively simple bottom topography and large wind stress events in both winter and summer. A more lightly instrumented, long-term, large-scale component was designed to help separate the local wind-driven response in the region of the small-scale experiments from motions generated either offshore by the California Current system or in some distant region along the coast, and also to help determine the seasonal cycles of the atmospheric forcing, water structure, and coastal currents over the northern California shelf.

The first small-scale experiment (CODE-1) was conducted between April and August 1981 as a pilot study in which primary emphasis was placed on characterizing the wind-driven "signal" and the "noise" from which this signal must be extracted. In particular, CODE-1 was designed to identify the key features of the circulation and its variability over the northern California shelf and to determine the important time and length scales of the wind-driven response. This report presents a basic description of the moored array data and some other Eulerian data collected during CODE-1. A brief description of the CODE-1 field program is presented first, followed by a description of the common data analysis procedures used to produce the various data sets presented here. Then basic descriptions of the following data sets are presented: (a) the coastal and moored meteorological measurements, (b) the moored current measurements, (c) the moored temperature and conductivity observations, (d) the bottom pressure measurements, and (e) the wind and adjusted coastal sea level observations obtained as part of the CODE-1 large-scale component.

Prepared for: NSF under Grant OCE80-14941.

WHOI Technical Report 83-23.

THE CAMS INTERACTIVE ATLAS PACKAGE:
A COMPUTERIZED PROGRAM FOR THE ARCHIVING
AND PRESENTATION OF OCEANOGRAPHIC DATA

Thomas A. Sgouros and Thomas Keffer

The CAMS Interactive Atlas Package allows a user to computer contour any hydrographic quantity on any isosurface in any global projection. It consists of two parts: a large, and growing, data base which can be easily supplemented with user-supplied data, and a set of computer software routines that manage the data, interpolate onto the isosurface, and draw the maps.

The data base presently includes the Levitus Climatological Atlas, IGY data, GEOSECS, TTO, and many NODC stations, all in a common format. In addition, the WHOI CTD data set can be easily accessed. This data base will be supplemented as new data becomes available.

The software routines allow the user to interpolate the data vertically onto a selected isopleth, such as an isobaric or isopycnal surface, and then horizontally interpolate the station data onto a regular grid for final contouring. Land mass areas can be automatically blanked out. The continental boundaries are drawn in and the contouring done in any one of fifteen projections. Station locations, cruise tracks, and bottom topography can all be included. The contour map can be plotted on either a vector plotter or a raster device, allowing the use of color in the final product.

CTD: conductivity-temperature-depth sensor
GEOSECS: the program to study the geochemical properties of the ocean with respect to large-scale circulations problems

IGY: International Geophysical Year (technically 1957-58, actually 1954-1962)

NODC: National Oceanographic Data Center

TTO: Transient Tracers in the Ocean program

Prepared for: The Center for the Analysis of Marine Systems (CAMS), WHOI, under funds provided by the Exxon Foundation, Mobil Foundation and an anonymous donor.

WHOI Technical Report 83-39.

USNS BARTLETT CRUISE 40-B, DATA REPORT

Marvel C. Stalcup, Terrence M. Joyce
and R. Lorraine Barbour

A joint cruise with Dr. Michael Gregg of the Applied Physics Laboratory at the University of Washington was conducted from 8-24 January, 1983, aboard the USNS BARTLETT to study the effects of wintertime cooling in a warm core ring. At the beginning of the cruise an XBT survey of ring 82I (found at 40°40'N, 66°W, east of the New England Seamounts) showed a rather confused pattern of surface temperature and salinity with the average depth of the mixed layer about 30 m. On January 16-17, a storm passed near the ring with winds to 45 knots and temperatures below 0°C. An XBT survey at the end of the cruise showed that vertical mixing and cooling during the outbreak of cold air resulted in a more coherent pattern in the surface temperature and salinity of the ring and an increase in the thickness of the mixed layer to 180 m.

XBT: expendable bathythermograph

Prepared for: ONR under Contract
N00014-82-C-0019, NR 083-004.

WHOI Technical Report 83-16.

THE LONG-TERM UPPER-OCEAN STUDY (LOTUS)
CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT
OCEANUS 119 - MAY 1982

Richard P. Trask and Melbourne G. Briscoe

OCEANUS cruise number 119 (6-14 May, 1982) was the seventh in a series of cruises to the Long-Term Upper-Ocean Study (LOTUS) area centered at 34°N, 70°W. During this cruise, five moorings were set at the LOTUS site; four for the LOTUS experiment and one, a profiling current meter mooring, for C. S. Draper Laboratory - MIT. In addition, an engineering mooring was set at 39°30'N, 70°W. Two XBT sections were made along 70°W between 40°N and 33°N during the trip to and from the LOTUS site. Five CTD stations were also completed in the LOTUS area.

Part I of this report is a summary of the major cruise activities, and part II presents the hydrographic data (CTD and XBT) collected during the cruise.

CTD: conductivity-temperature-depth sensor
XBT: expendable bathythermograph

Prepared for: ONR under Contract
N00014-76-C-0197, NR 083-400.

WHOI Technical Report 83-7.

THE LONG-TERM UPPER-OCEAN STUDY (LOTUS)
CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT
OCEANUS 129 - OCTOBER 1982

Richard P. Trask and Melbourne G. Briscoe

OCEANUS cruise number 129 (28 October - 4 November, 1982) was the eighth in a series of cruises to the Long-Term Upper-Ocean Study (LOTUS) area centered at 34°N, 70°W. In the LOTUS area, seven SOFAR floats were launched, two moorings were recovered (a LOTUS surface mooring and a C. S. Draper Laboratory profiling current meter mooring), and a surface mooring which replaced the one recovered was set. Seven CTD stations were also completed in the LOTUS area. Outside the LOTUS area, a subsurface mooring was set in the Gulf Stream in cooperation with H. Bryden's (WHOI) Gulf Stream Observations project and a WHOI engineering mooring at Site D was recovered, examined and redeployed. In addition, several XBT sections were made, one along 70°W between 40°N and 34°N, a second surveyed a cold core ring discovered during the trip to the LOTUS area and a third was made in the vicinity of the Gulf Stream Observations mooring.

Part I of this report is a summary of the major cruise activities and part II presents the hydrographic data (CTD and XBT) collected during the cruise.

SOFAR: sound fixing and ranging

Prepared for: ONR under Contract
N00014-76-C-0197, NR 083-400.

WHOI Technical Report 83-29.

THE LONG-TERM UPPER-OCEAN STUDY (LOTUS)
CRUISE SUMMARY AND HYDROGRAPHIC DATA REPORT
ENDEAVOR 97 - April 1983

Richard P. Trask and Melbourne G. Briscoe

ENDEAVOR cruise number 97 (8-19 April, 1983) was the ninth scheduled cruise to the Long-Term Upper-Ocean Study (LOTUS) area centered at 34°N, 70°W. During the cruise three LOTUS moorings (a near-surface and two sub-surface moorings) deployed eleven months earlier were recovered and replaced by a nearly identical set of moorings. The new array will remain in the water during the final year of LOTUS field work. The LOTUS surface mooring, scheduled to be recovered during ENDEAVOR 97, had been partially recovered one month earlier after the mooring parted and drifted off station. The lower portion of the surface mooring which went to the bottom when the mooring failed was successfully recovered during ENDEAVOR 97. A new surface mooring replacing the one that parted and a C. S. Draper Laboratory profiling current meter mooring

were also set during the cruise.

Non-mooring work included deploying three satellite tracked drifter buoys and completing five CTD stations in the LOTUS area. Several intercomparisons between shipborne meteorological sensors and similar sensors on the LOTUS surface buoy and the drifter buoys were made. An XBT section was also completed along 70°W between 40°N and 34°N.

Part I of this report is a summary of the major cruise activities and part II presents the hydrographic data (CTD and XBT) collected during the cruise.

Prepared for: ONR under Contract
N00014-76-C-0197, NR 083-400.

WHOI Technical Report 83-33.

FINAL TECHNICAL REPORT
CONTRACT N00014-81-C-0010
1 OCTOBER 1980 - 30 SEPTEMBER 1982

John A. Whitehead, Jr.

Laboratory, theoretical and numerical research was conducted into the structure and stability of baroclinic non-linear currents in a rotating fluid. A rotating version of the dam-break problem in which a density current is generated after a barrier has been removed was studied.

Theoretical studies of the stability of a free isolated baroclinic jet whose free surface in cross-section intersects the water surface at two points by Griffiths, Killworth and Stern (J. Fluid Mech., 117, 343-377, 1982) was undertaken. Killworth and Stern (Geophys. Astrophys. Fluid Dyn., 22, 1-28, 1982) showed that a coastal density current in a rotating system is unstable to downstream wave disturbances when the mean potential vorticity increases towards the (vertically-walled) coast and when the mean current vanishes there.

Paldor, in his Ph.D. thesis (Univ. Rhode Island, 1981), used Rayleigh integral to prove that an unbounded geostrophic front of uniform potential vorticity is stable with respect to small perturbations of arbitrary wavelength. Stern and Paldor (Phys. Fluids, 26, 906-919, 1983) used extremum concepts to analyze large amplitude disturbances in a boundary layer shear flow with an inviscid and longwave theory.

Prepared for: ONR under Contract
N00014-81-C-0010 and for NSF under
Grant OCE80-18322.

WHOI Technical Report 83-11.

MARINE POLICY
AND OCEAN MANAGEMENT PROGRAM

David A. Ross, Director

THE SOCIOLOGY OF PRODUCTION IN RURAL
MALAY SOCIETY

Conner Bailey

The basic question addressed in this book is how differing modes of production and the social relations of production which they engender influence broader patterns of social behavior. Three rural Malay communities in Peninsular Malaysia, each with distinctly different economic adaptations (rice farming, rubber tapping, and fishing), provide empirical reference for this comparative case study.

Following detailed descriptions of these three communities and the manner in which they are organized economically, separate analytical chapters compare differences in labor organization and capital formation.

It was found that differences in how communities are organized for production purposes systematically influence non-economic social organization of these communities. It is argued that these differences are relevant to the design of workable rural development programs, which often fail to consider variability among rural communities.

Published by: Oxford University Press,
Kuala Lumpur. 1983. 225 pp.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

SMALL SCALE FISHERIES OF SAN MIGUEL BAY,
PHILIPPINES: OCCUPATIONAL AND
GEOGRAPHIC MOBILITY

Conner Bailey

Existing patterns and future potentials for occupational and geographic mobility among small-scale fishermen of San Miguel Bay, Philippines, were examined to determine whether such mobility had led or is likely to lead to a reduction of surplus fishing labor or improvements in the productivity and incomes of those fishermen who remain.

Existing alternatives to fishing offer only limited potential for absorbing labor from the fisheries sector. Examination of census data at the community level for the period 1939-80 using census-survival techniques indicated substantial net out-migration from the San Miguel Bay area. Nonetheless, in absolute terms, numbers of fishermen have increased during this period, contributing to heavy pressure on the Bay's marine resources. Equally significant in terms of fishing effort were trawlers, which began operating within the Bay during the 1970s.

The issue of competition between

small-scale fishermen and trawler operators in San Miguel Bay was discussed. In the long term, the encouragement of economic alternatives to fishing was found to be essential, but in the short term, efforts to improve conditions among small-scale fishermen might more effectively be based on better enforcement of current management regulations, which are designed to limit competition between small-scale fishermen and trawlers.

Published in: ICLARM Technical Reports
10, 57p. Institute of Fisheries
Development and Research, College
of Fisheries, University of
Philippines in the Visayas, Quezon
City, Philippines; International
Center for Living Aquatic Resources
Management, Manila, Philippines;
and the United Nations University,
Tokyo, Japan.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

SMALL SCALE FISHERIES OF SAN MIGUEL
BAY, PHILIPPINES: SOCIAL
ASPECTS OF PRODUCTION AND MARKETING

Conner Bailey

This report presents the edited findings of a team of sociologists working on an inter-disciplinary research project on the fisheries of San Miguel Bay, Philippines. The papers in this report describe the mix of field research methodologies employed (including detailed survey and participant observation); discuss characteristics of fishing families and the economic roles of women and children within the study area; assess the impact of seasonality of fishing, marketing, and processing activities; analyze variations found in sharing systems and ownership patterns; discuss problems in the fisheries of San Miguel Bay as perceived by local fishermen.

Published in: ICLARM Technical Re-
ports 9. Conner Bailey, ed. 57 p.
Institute of Fisheries Development
and Research, College of Fisheries,
University of Philippines in the
Visayas, Quezon City, Philippines;
International Center for Living
Aquatic Resources Management,
Manila, Philippines; and the United
Nations University, Tokyo, Japan.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

SYSTEMS OF SHARING AND
PATTERNS OF OWNERSHIP

Conner Bailey and E.D. Villafuerte

Sharing systems and patterns of ownership of several common small-scale gears are analyzed and compared with those of the small and medium trawlers operating in San Miguel Bay, Philippines. Significant differences between these two groups were found in concentration of ownership, presence of non-economic social relationships between owners and crewmen, and flexibility of sharing arrangements. These differences are discussed in terms of existing legal definitions of "municipal" and "commercial" fisheries.

Published in: Small Scale Fisheries of San Miguel Bay, Philippines: Social Aspects of Production and Marketing. C. Bailey, ed. ICLARM Technical Reports 9, 25-41. Institute of Fisheries Development and Research, College of Fisheries, University of Philippines in the Visayas, Quezon City, Philippines; International Center for Living Aquatic Resources Management, Manila, Philippines; and the United Nations University, Tokyo, Japan.

Supported by: Pew Memorial Trust and Marine Policy and Ocean Management Program.

MANAGING AN OPEN ACCESS RESOURCE:
THE CASE OF COASTAL FISHERIES

Conner Bailey

Coastal fisheries in many parts of the developing world are coming under increasingly intense pressure both by local artisanal fishermen and by operators of commercial trawlers fishing for penaeid shrimp found in such waters. It is argued that where such competition exists, access to the resource may be "open" but, given the greater fishing power of commercial trawlers, is inherently unequal. The case of San Miguel Bay, Philippines, is used to describe the nature of this competition.

It is argued that artisanal fishermen need to organize themselves politically to effectively assert and protect their traditional rights to this fishery. Fisheries cooperatives provide an existing institutional basis for such legitimate political action, but have had distinctly limited success in attracting active memberships. Articulation of this locally important concern could promote more active participation by fishermen in fisheries cooperatives. If effectively organized,

the over 5,000 artisanal fishermen in the San Miguel Bay area could be an important political constituency capable of influencing local municipal governments, which legally are responsible for managing approximately half of the Bay's waters.

In press: People-centered Development: Contributions Toward Theory and Planning Frameworks. David C. Korten and Rudi Klaus, eds. Kumarian Press, Spring 1984.

Supported by: Pew Memorial Trust and Marine Policy and Ocean Management Program.

DEVELOPING A U.S. RESEARCH STRATEGY
FOR MARINE POLYMETALLIC SULFIDES

James M. Broadus and Robert E. Bowen

Marine polymetallic sulfides (MPS) are recently discovered mineral occurrences of hydrothermal origin. Design of an appropriate research strategy to investigate the MPS deposits is a policy issue that the paper seeks to help resolve. The paper describes the relevant nature of MPS, reviews what has happened in MPS research so far, examines market conditions facing the contained metals, and presents some useful principles for creation of a scientifically and economically effective research strategy. This strategy is examined in several key dimensions: science versus R&D; role of government and private industry; geographic concentration; focus in time; and the locus of responsibility and scope of participation. Because of the early stage of work on MPS and the scant scientific evidence available, expected results of R&D are extremely uncertain. In this context, a sequential scientific approach is favored for government support. The strategy suggested by the paper's analysis would permit geographically varied work sites and would incorporate diversification of research approaches, decentralization of discretionary decision making, and a broad base of participation.

In press: Ocean Development and International Law.

Supported by: Pew Memorial Trust; NOAA Grant NA80AA-D-00077 (R/G-6); and Marine Policy and Ocean Management Program.

EVALUATING THE ECONOMIC SIGNIFICANCE
OF POLYMETALLIC SULFIDES DEPOSITS

James M. Broadus and Robert E. Bowen

The current economic significance of ocean ridge polymetallic sulfide deposits is evaluated. Present and future sources of demand and supply for the contained metals are considered along with the potential for and conditions likely to foster substitution away from their use and the implications of technical change for production costs over time. Estimates of future value are discounted into present terms. International legal and political parameters are examined. The deposits appear to have negligible present resource value, but knowledge about them is valuable in itself.

Published in: Proceedings of 15th
Annual Offshore Technology
Conference, Houston, Texas, May
2-5, 1983, OTC 4493.

Supported by: Pew Memorial Trust;
NOAA Grant NA80AA-D-00077 (R/G-6);
and Marine Policy and Ocean
Management Program.

CONFLICT RESOLUTION IN THE ASSIGNMENT
OF AREA ENTITLEMENTS FOR SEABED MINING

James M. Broadus and Porter Hoagland, III

The paper describes conditions creating a need for, and the procedures being implemented to achieve, resolution of conflicts over areas of the deep seabed being sought for exclusive exploration entitlements. If and when deep seabed mining takes place, certain tangible advantages will accrue to nations or organizations that have obtained entitlements to exclusivity of activity within areas of the seabed. Development of legal assurances of exclusivity and security of work within a claimed area has been proceeding along two different tracks: (1) The LOS Convention; and (2) a prospectively complementary, but currently separate and potentially competing entitlement system, based on domestic laws, multilateral arrangements and private agreements. Because the areas sought by prospective seabed miners under both tracks overlap in varying degrees, attempts to resolve conflicts are a part of each track. Successful resolution in either track could facilitate progress in the other, and there is a possibility that arrangements developed through efforts in the separate tracks could contribute eventually to a convergence between the two legal-regulatory regimes.

In press: San Diego Law Review.

Supported by: Pew Memorial Trust; NOAA
Grant NA80AA-D-0007 (R/S-7) and
Marine Policy and Ocean Management
Program.

THE MONTEGO BAY CONVENTION OF 1982
AND THE CUSTOMARY LAW OF THE SEA:
OBSERVATIONS, A FRAMEWORK
AND A WARNING

John King Gamble, Jr.

Now that a new Law of the Sea Convention has been signed by more than 130 states, it becomes especially important to inquire about the relationship between that Convention (the Montego Bay Convention) and the customary law of the sea. It is tempting, yet counterproductive and naive, to reduce this matter to the simple question, does the M.B.C. codify existing customary law? The M.B.C. is so comprehensive and heterogeneous that provisions will differ in the degree to which they codify existing customary international law. This suggests the need for at least three categories into which provisions can be placed: (a) those reinforcing traditional (old) customary law; (b) those giving written expression to new customary law developing since 1958; and (c) those directing the development of future customary international law. The paper makes suggestions about how this framework can be applied to the M.B.C. Further, national perceptions of the customary law reflectivity of the M.B.C. are examined along with national legislation concerning seabed mining. The interface between the M.B.C. and customary law will be complex and dynamic -- it will be years before the major forces have run their course and a balance is achieved.

In press: San Diego Law Review, 21,
1984.

Supported in part by: Pew Memorial
Trust; Johnson Endowment Fund;
Marine Policy and Ocean Management
Program.

THE SIGNIFICANCE OF SIGNATURE TO THE 1982
MONTEGO BAY CONVENTION ON THE LAW
OF THE SEA

John King Gamble, Jr.

On December 19, 1982 in Montego Bay, Jamaica, 117 states signed a new and comprehensive Law of the Sea Convention. The complexity of this Convention (the M.B.C.) coupled with the lethargy with

which states generally follow through with ratification suggests it may be years before the M.B.C. enters into force. This paper examines the significance of signature to the Convention. For many states an obligation not to defeat the object and purpose of a treaty between signature and force exists; the degree of this obligation is examined. Additionally, certain states continue to be bound by the 1958 Geneva Conventions; this may affect the M.B.C. prior to entry into force. A wide range of behavior is permissible without defeating the object and purpose of a treaty (an irreversible harm test has been suggested) -- there are relatively few actions that clearly will defeat the object and purpose of the M.B.C. -- the best examples are in uses of the deep sea bed. Overall, signature does create obligations, limited in scope and application, but still important.

In press: Ocean Development and International Law Journal, 13, 1984.

Supported by: Pew Memorial Trust; Johnson Endowment Fund and Marine Policy and Ocean Management Program.

MULTIATTRIBUTE ANALYSIS AND THE DEFINITION OF OPTIMUM YIELD

Michael C. Healey

This paper reviews the origin and operational definition of the optimum yield (OY) concept and demonstrates how technique of decision analysis can provide an analytical model for OY. The concept of OY was formalized as the guiding principle of fisheries management in the U.S. and Canada in 1976. The policies of both countries make it clear that a wide range of biological, economic and social factors are to be taken into account in determining OY. Confusion exists, however, about precisely which of these factors should determine OY in any fishery and what are their relative importance. Uncertainty also exists about how to take biological, economic and social factors jointly into account as the concept of OY implies one must. Established biological and economic models in fisheries are not adequate for such analysis because their focus is single -- rather than multiobjective. Operational techniques of decision analysis, such as multiattribute utility analysis, are specifically designed to deal with multiobjective problems like OY. I propose that a simple, linear, utility model be used to assess the optimality of alternative yield strategies in fisheries management. I illustrate the application of the model by assessing OY options in the Skeena River salmon fishery and the

New England herring fishery. The advantages of the model are that it is simple and intuitively appealing, that it permits a wide range of types and qualities of data to be incorporated into the evaluation of management options, that it is amenable to sensitivity analysis and that it is adaptable to a variety of decision rules.

In press: Canadian Journal of Fisheries and Aquatic Sciences 41, 6.

Supported by: Pew Memorial Trust; Johnson Endowment Fund; NOAA Grant NA80AA-D-00077 (E/L-1) and Marine Policy and Ocean Management Program.

INTER- AND INTRAPOPULATION VARIATION IN THE FECUNDITY OF CHINOOK SALMON (*Oncorhynchus tshawytscha*) AND ITS RELEVANCE TO LIFE HISTORY THEORY

Michael C. Healey

Chinook salmon varied significantly in fecundity both between years within populations and between populations throughout their range. Generally, less than 50% of the variation in fecundity between individuals within populations could be explained by variation in length. A small additional amount of variation could be attributed to racial differences (e.g., red or white fleshed types) but age, seasonal timing of sub-populations and stream or ocean type life history pattern did not contribute significantly to variation in fecundity beyond their correlation with length. A great deal of individual variation in fecundity remains to be explained in chinook salmon. The slopes of the regression of fecundity on length for all populations were low in comparison with other fishes, indicating that fecundity increases slowly with increasing size in chinook. The mean age of reproducing females varied among populations, and populations which reproduced at an older age were more fecund at a common length than populations which reproduced at a younger age. The increase in fecundity with increasing age of maturity was consistent with theoretical predictions about the trade-off between fecundity and mortality in fish of reproductive age. The mean age of reproduction within a population, however, was considerably older than the predicted optimum age of reproduction based on the trade-off between increasing fecundity with age, and natural mortality. These observations suggest that chinook have sacrificed fecundity for increased size in the allocation of surplus energy, a life history characteristic that is consistent with the survival value of size in anadromous salmon.

In press: Canadian Journal of Fisheries and Aquatic Sciences 41, 4.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund; NOAA Grant
NA80AA-D-00077 (E/L-1) and Marine
Policy and Ocean Management Program.

VARIATION IN FECUNDITY OF LAKE WHITEFISH
(Coregonus clupeaformis) FROM LESSER
SLAVE AND UTIKUMA LAKES IN
NORTHERN ALBERTA

Michael C. Healey and K. Dietz

The fecundity of lake whitefish from Lesser Slave Lake varied significantly between 1971 and 1979. Fecundity adjusted to 400 mm fork length was lowest in 1971 (12900) and highest in 1978 (21200). Utikuma Lake whitefish were only sampled in 1976, but in that year had the highest fecundity of any population yet recorded (44700 at 400 mm). These data extend the observed range of average fecundity among whitefish populations from 2 fold to 3 fold. Fecundity and condition were positively correlated within and between years in Lesser Slave Lake and between populations of whitefish. Condition adjusted for gonad weight, however, was not correlated with fecundity. These correlations indicate that more fecund whitefish have accumulated more surplus energy and that once maintenance demands are met, maturing whitefish put all excess energy into gonad growth.

In press: Copeia 1, 1984, pp. 238-242.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund; NOAA Grant
NA80AA-D-00077 (E/L-1) and Marine
Policy and Ocean Management Program.

FISH PREDATION ON AQUATIC
INSECTS

Michael C. Healey

The purpose of this chapter is to explore three aspects of the interaction between fish and insects: functional adaptations of fish as predators, methods for calculating the number of insects eaten by fish in natural populations, and the impact of fish predation on the abundance and composition of the insect community. It concentrates on the fundamental problems and shortcomings in our current understanding in the hope of stimulating some creative research in this field.

In press: Ecology of Aquatic Insects:
A Life History and Habitat Approach.
Vincent H. Resch and David M.
Rosenberg, eds. Praeger Special
Studies, Praeger Scientific, New
York, N.Y. Spring 1984. Chapter 9.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund; NOAA Grant
NA80AA-D-00077 (E/L-1) and Marine
Policy and Ocean Management Program.

THE U.S. FISHING INDUSTRY AND
REGULATORY REFORM

Timothy Hennessey

This book contains a series of articles by government officials and academics regarding the relationship between government and the fishing industry. Topics include the regulatory review process, how the fisheries management councils view regulatory reform, prospects for rational regulation and de-regulation and trade and development issues.

Published by: Wilson Publishing Co.,
Wakfield, R.I. 1983. 229pp.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund and Marine
Policy and Ocean Management Program.

HOW U.S. OCEAN POLICY IS MADE AND UNMADE

Robert W. Knecht

Policy-making can result from carefully structured and well supported studies of realistic options done significantly in advance of any crisis or it can be done as a reactive exercise when events conspire to require more-or-less immediate action on new or unanticipated problems. While examples of both types can be found in ocean policy-making in the U.S. in recent decades, ad-hoc policy making in response to specific well-developed problems is by far the most common. The origins and salient features of existing U.S. ocean policy are reviewed and the forces likely to bring about change are examined. It is argued that certain of these factors are such that a significant rethinking of U.S. policy with regard to the oceans is likely to be needed.

In press: Proceedings, Coastal Zone
'83, San Diego, June 1983.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund and Marine
Policy and Ocean Management Program.

OCEAN POLICY-MAKING IN A CHANGING WORLD

Robert W. Knecht and Biliiana Cicin-Sain

A number of elements in the U.S. ocean policy-making landscape have undergone considerable change in the last decade or so. Foremost among these, of course, are changes in the international legal framework brought about as a result of the Law of the Sea (LOS) Convention and its prolonged deliberations. But important changes in the domestic ocean policy picture have occurred as well; the large body of ocean-oriented legislation enacted by the U.S. Congress in the 1970s attests well to this fact. These changes, both on the international and domestic fronts, will significantly structure discussion of ocean policy issues in the years ahead.

In the first part of this paper, "The Changed Ocean Context," we examine, with a broad brush approach, the major ocean policy changes, both domestic and international, that have taken place in recent years. In a second section, "Recurrent Tensions in U.S. Ocean Policy-Making," we assess those factors or influences which appear to have been most important in shaping U.S. ocean policy over time. In the concluding part, "Emerging Ocean Policy Issues and Opportunities," we describe some of the new ocean policy issues and problems that will need to be addressed in the near future. We suggest, as well, certain changes in the ocean policy development process that will need to take place to properly respond to both the complexity and volatility of some of the ocean issues that lie ahead.

In press: Proceedings, Marine Technology Society Oceans '83, San Francisco, September 1983.

Supported by: Pew Memorial Trust; Johnson Endowment Fund and Marine Policy and Ocean Management Program.

ECONOMIC INCENTIVES FOR ENVIRONMENTAL PROTECTION: OCEAN DUMPING CHANGES

William L. Lahey

The question of where to dispose of society's waste has become an issue of major social import. Ocean dumping is one waste disposal alternative which is being widely considered, especially in light of the problems associated with land-based disposal. The existing regulatory regime, however, appears ill-suited to the sound management of the ocean as a waste disposal medium.

This article considers a new approach

toward regulating ocean disposal -- one based on economic incentives. The theoretical rationale behind the pollution charge approach, which have been elaborated in detail in the economic literature, are summarized. Efforts, both at the state level and abroad, to translate the theory into practice are selectively critiqued. The article presents the justifications for adopting an ocean dumping charge. Finally, a number of issues, especially associated with implementing this approach are evaluated in detail.

The article concludes that there are a number of compelling justifications for adopting a charge system for ocean dumping. There are, however, a number of administrative, political and legal questions which must be answered as a prerequisite to a viable charge system.

In press: Ecology Law Quarterly.
Spring 1984.

Supported by: Pew Memorial Trust and Marine Policy and Ocean Management Program.

THE CASE FOR OCEAN WASTE DISPOSAL

William Lahey and Michael Connor

The use of the ocean for waste disposal is growing. What are the reasons for this trend? Are the scientific arguments for this trend sound? Are our laws and regulations equipped to effectively respond to this trend? This article offers answers to each of these questions.

The increasing interest in ocean disposal of waste is an outgrowth of a number of economic, political, legal and scientific factors. Vociferous public opposition to land disposal, for instance, has resulted in increased attention to ocean disposal. The cost of land disposal is often two or three times as much as ocean dumping. These factors along with some recent judicial rulings and scientific findings have spurred more ocean waste disposal.

There are, however, a number of reasons why expansion of our use of the ocean for waste disposal should proceed cautiously. Our ability to determine the assimilative capacity of a certain ocean disposal site is very rudimentary. A comparison of the health risks associated with land-based disposal versus ocean disposal, albeit crude estimations, suggests that ocean dumping presents a higher risk than land disposal.

The article concludes with a recommendation for a new federal program for regulatory ocean disposal, one based on a sliding fee schedule based on where the dumping takes place and what is being dumped.

Published in: Technology Review, 86,
6, August/September 1983.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

AMERICA AND THE MIDDLE EAST: HOLDING THE CENTRE

Linda B. Miller

The September 1982 Reagan peace proposals are examined in terms of background, content and reaction in order to explore the validity of the idea of a political centre in U.S. foreign policy. Given the fact that partition has no roots in contemporary Arab nationalism, the Reagan plan is unlikely to break the protracted deadlock in the region. Nevertheless, well publicized peacemaking efforts are always good politics domestically. This proposal's major importance will be felt in Washington rather than in the Middle East.

Published in: The World Today,
39, 1, January 1983.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

ENERGY AT THE SUMMIT

Linda B. Miller

Since 1975, the annual economic summits of leaders from seven industrial countries and the European Community (EC) have joined the already luxuriant field of overlapping institutions through which the Western alliance partners' energy policies and practices are supposed to be reconciled.

Ideally, these gatherings, plus the established vehicles of the International Energy Agency (IEA), the OECD, and NATO, could constitute a smoothly functioning, "well-oiled" machine that would produce a comprehensive energy strategy integrating political, economic, and military factors. But organizational neatness, even if it did exist, does not guarantee coherence. An analysis of the summit process and related forums illustrates the persistent gap between theory and practice in Western energy planning.

Published in: Geopolitics of Energy,
5, 10, October 1983.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

THROUGH A GLASS DARKLY: WESTERN EUROPE AND THE MIDDLE EAST

Linda B. Miller

The impact of the Iran-Iraq war and the Israeli intervention in Lebanon on West European foreign policies is analyzed. Elements of continuity and change in national attitudes and in the role of the European Community in the Middle East are highlighted. While major shifts occurred during the "oil crisis" of 1973-74, more recent events have underscored the circumscribed nature of European influence in the region as the former colonial powers rely on arms supplies, trade and exports of manufactured goods for influence.

Published in: Middle East Review,
XV, 1 and 2, Fall/Winter, 1982-83.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

OIL, SHRIMP, MANGROVES: AN EVALUATION OF CONTINGENCY PLANNING FOR THE GULF OF GUAYAQUIL, ECUADOR

Ivon d'Almeida Pires Filho

The possibility of finding oil in the Gulf of Guayaquil has led several Ecuadorian agencies to prepare contingency plans to deal with the eventuality of an oil spill in the area. This report characterizes the importance of the oil and fisheries industries to the Ecuadorian economy, and describes the region where these activities may conflict. It also elaborates on the biological effects of oil in tropical environments, and on aspects of prevention, control/clean-up and oil spill contingency planning. Compensation for oil pollution damages and methods for damage assessment are also discussed herein.

The analysis comments on specific issues of the Ecuadorian plans, such as their oil spill response organization and operational guidelines. It notes the willingness of the government and industry to handle the problem jointly and to do so prior to actual oil production. The combination of control/clean-up methods considered demonstrates the seriousness and sophistication of these plans. However, the planning process seems to overly rely on these clean-up measures as an antidote to oil spills. This report emphasizes prevention as the cheapest and the most efficient approach to protect the marine environment and resources of the Gulf of Guayaquil from oil pollution damages.

Supported by: Pew Memorial Trust;
Johnson Endowment Fund; William H.
Donner Foundation and Marine Policy
and Ocean Management Program.

WHOI Technical Report 83-38.

PROSPECTIVE MARITIME JURISDICTIONS IN THE POLAR SEAS

Richard L. Price and Marc E. Vaucher

The maps illustrate the possible extent of the 200-nautical-mile exclusive economic zone in the polar regions. They also show the 500 m and 2,500 m isobaths as they relate to prospective economic zones. Equidistance lines are shown between countries, and where negotiated maritime boundaries exist these too are shown. The maps and accompanying text indicate where disputes may eventually arise over maritime jurisdictions as nations begin claiming sovereignty over offshore areas.

Published in: Annals of the Association of American Geographers, 73, 4, 617-618, 1983.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

WHOI Technical Report 83-1.

EFFECTIVE USE OF THE SEA OVERCOMING THE LAW OF THE SEA PROBLEMS

David A. Ross

It is evident that the Law of the Sea Treaty and the extended jurisdictional claims already made by many coastal countries have led to various problems for U.S. marine scientists who wish to work in foreign waters. Likewise, there is a potential for problems in the future, regardless of the eventual U.S. position on the treaty. The basic premise of this paper is that in spite of these problems there will also be many valuable opportunities for marine science and marine scientists, but to realize these will require new approaches and skills.

In spite of the important need for cooperative marine science programs there is no coherent, focused U.S. effort to encourage work with individual foreign countries. Although U.S. projects and contacts with individual foreign countries are common, they are spread among many federal and private organizations who are often unaware of what others are doing. As confusing as this is for U.S. scientists or organizations, it is even worse for foreign

scientists or countries. A specific recommendation of this paper is to establish a central clearing group in the United States capable of providing information to developing countries on the types of technical assistance available and by what organizations. Such a central group would provide a point of contact for those seeking assistance and for those who can provide it. The need for a coordination and information office for cooperative U.S. marine assistance activities has also been often stated by both U.S. and foreign officials.

In press: Proceedings of Marine
Technology Society Oceans '83,
San Francisco, September 1983.

Supported by: Pew Memorial Trust; NOAA
Grant NA80-AA-D-00077 (E/L-1) and
Marine Policy and Ocean Management
Program.

THE SHELFBREAK: SOME LEGAL ASPECTS

David A. Ross and K.O. Emery

Conferences on Law of the Sea have had the objective of increasing the area of ocean floor subject to control by adjacent coastal countries. These extensions of jurisdiction have paid little attention to carefully defined and relatively easily identified geological boundaries such as the shelfbreak. Indeed, a geological term often is used in a legal sense that far exceeds the geological meaning, resulting in unnecessary confusion. The recently concluded Third United Nations Conference on the Law of the Sea adds an area of the ocean subject to national control essentially equal to that of the land area of the world. Certain aspects of the remaining area of deep-ocean floor, such as mining, will also be controlled and taxed by an international authority. It is possible future oceanographers may have little opportunity for research without permission and regulation by governments of either coastal nations or the United Nations. One result could be increased research and knowledge of the ocean floor that is under the jurisdiction of industrialized countries and decreased effort in the rest of the ocean.

Published in: SEPM Special Publication,
33, 437-441, June 1983.

Supported by: Pew Memorial Trust; NOAA
Grant NA80AA-D-00077 (E/L-1) and
Marine Policy and Ocean Management
Program.

WHOI Contribution No. 5198.

U.S.-MEXICO MARINE SCIENCE BILATERAL
AGREEMENTS: WHAT SHOULD BE INVOLVED?

David A. Ross

In a meeting that considers U.S.-Mexico relations on marine resources, it certainly is appropriate to also discuss marine science and, in particular, the possibilities of bilateral scientific relationships between our two countries.

Bilateral scientific arrangements between adjacent countries is not a new subject; it has been discussed within the Law of the Sea Treaty context, within the U.S. Presidential Proclamation of 10 March 1983 and in various forums between representatives of our two countries. In general, bilateral arrangements or agreements governing marine science and scientific operations can often be of value for adjacent countries. In the specific case of the United States and Mexico, there are many areas of marine scientific research that are of mutual interest and that could play an extremely important role in furthering economic, social and scientific development in the region.

It is well known that oceanographic phenomena and processes rarely respect political boundaries. Many important aspects of the ocean extend beyond a country's territorial sea or EEZ and impact on other adjacent or even further distant countries. If the ocean and its resources are to be used effectively, consideration must often be given to the marine conditions of an entire region and how one country's use will impact on adjacent countries. To do otherwise may prove to be shortsighted and may result in unnecessary political, economic or scientific costs.

In press: Proceedings of U.S. Mexican Relations on Marine Resources: A Binational Conference, Univ. of Calif. at San Diego, Sept. 15-17, 1983.

Supported by: Pew Memorial Trust; NOAA Grant NA80-AA-D-00077 (E/L-1) and Marine Policy and Ocean Management Program.

INTERNATIONAL JURISDICTION ISSUES
IN THE ARCTIC OCEAN

Kurt M. Shusterich

Awareness in the United States of the strategic value of the Arctic has been well known for many years. Appreciation of its economic, political, and scientific value, as well as its juridical importance is rapidly increasing. For several years, both the Soviet Union and Canada have had

more ongoing commercial, scientific, and policy activities in the Arctic region than the United States. Resource development, national security, and marine boundary issues promise to raise the level of awareness and importance of the Arctic region to the United States over the next twenty years. The purpose of the article is to indicate the range and complexity of marine boundary jurisdiction issues in the Arctic.

In press: Ocean Development and International Law.

Supported by: Pew Memorial Trust and Marine Policy and Ocean Management Program.

WHOI Contribution No. 5333.

ASSESSING THE OCEAN'S ROLES IN
ANTILLEAN PREHISTORY

David R. Watters

Antillean archaeologists have been overly concerned with relationships between prehistoric human populations and the land, such that potentially significant interactions with marine and estuarine environments have been neither adequately explored nor sufficiently considered. Preoccupation with a landward orientation has created a "terrestrial bias" perceptible in some interpretations of the region's prehistory. The central theme and expressed intent of this paper is to promote the adoption of a seaward perspective to complement, but not to replace, the landward view. We advocate research incorporating both viewpoints in order to comprehend better the cultural adaptations developed by prehistoric Antillean peoples.

Published in: The Proceedings of the Ninth International Congress for the Study of the Pre-Columbian Cultures of the Lesser Antilles. Centre de Recherches Caraibes, Universite de Montreal, Montreal (1983).

Supported by: Pew Memorial Trust and Marine Policy and Ocean Management Program.

WHOI Contribution No. 4982.

HIGHLAND HOUSE, BARBUDA:
AN 18th CENTURY RETREAT

David R. Watters and Desmond V. Nicholson

Barbuda is situated between 17°31' -

17°45'N and 61°53'W in the northern Lesser Antilles. It has a total area of approximately 160 km² and two readily distinguishable topographic units: (a) the Highlands plateau in the east-central part reaching a maximum elevation of about 45 m and (b) a surrounding marginal plain that occupies about 75% of the island's landmass. A number of edaphic and climatic constraints effectively precluded large-scale production of sugar, tobacco, and other commercially important crops during the colonial era. Shallow, discontinuously distributed, poorly drained soils, inadequate and unreliable rainfall, large expanses of bare limestone and accumulated sand, and the absence of surface streams rendered the island unsuitable for plantation crops, and, instead, Barbuda became a "provisioning station" for nearby islands, supplying items such as livestock, hides, meat, corn, and wood.

After two brothers, John and Christopher, acquired the first Codrington lease in the 1680s, the Codrington family continued to lease Barbuda from the British Crown for almost two hundred years until 1870. Most historic sites on Barbuda are from the period of the Codrington leasehold. To date, ten sites have been designated; nine were examined; surface collections were made at five; test pits were excavated at two. Other historic sites are known to exist but they have yet to be designated or observed.

Published in: The Florida
Anthropologist 35, 4.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

ALTERNATIVE REGIMES FOR FUTURE MINERAL RESOURCE DEVELOPMENT IN ANTARCTICA

William E. Westermeyer

The operation of the Antarctic Treaty is subject to review in 1991. It has functioned as a model of international cooperation for twenty-two years, but stresses on the Treaty system are increasing. In particular, interest is growing concerning the mineral resource potential of Antarctica, but the Antarctic Treaty does not address resource allocation issues. Efforts are currently underway to devise a minerals regime for Antarctica, but negotiations could take several years. If negotiations do not produce a new regime by the review date, some countries may decide to withdraw from the Treaty, signaling the end of the cooperative era in Antarctica. This paper reviews and evaluates some of the major options for a minerals regime. It is suggested that the most likely alternatives for Treaty states are those joint jurisdic-

tion options which sidestep the contentious issue of sovereignty. External accommodation with the international community will remain a difficult problem. While the joint jurisdiction options contain several provisions for sharing with outsiders, there is room for improvement.

Published in: Ocean Management, 8,
197-232, 1982/83.

Supported by: Pew Memorial Trust and
Marine Policy and Ocean Management
Program.

WHOI Contribution No. 5358.

GRADUATE STUDENTS

Abstracts of papers or theses submitted in 1983 by graduate students of the Woods Hole Oceanographic Institution Doctoral Degree Program and the Woods Hole Oceanographic Institution/Massachusetts Institute of Technology Joint Program in Oceanography/Oceanographic Engineering. Other papers authored or coauthored by graduate students are included in the departmental sections. Students are indicated by an asterisk in the Author Index.

AN IMPROVED LORAN-C DRIFTING BUOY
AND DROGUE FOR COASTAL APPLICATIONS

William J. Burke

The engineering development of a second generation Loran-C drifting buoy system is presented. This system is designed for use in the coastal ocean environment and has the capability to take Loran-C position fixes hourly and to transmit these fixes back to a shore station on a daily basis.

A custom surface buoy was designed for use with the system and was found to have favorable hydrodynamic drag and stability characteristics while providing a secure, watertight, and protected platform for the system electronics and power supply.

An improved drogue was also developed and tested on both a scale model and full scale basis. It was found to have a drag coefficient on the order of $2.15 + .14$, to be free from error-inducing behavior such as kiting or towing at an angle to the current, and to have minimal ballast requirements.

The system was field tested and a sample experiment performed, from which it was determined that the system is relatively simple to deploy and recover and has a somewhat uncertain positioning error of approximately 100 feet.

A hydrodynamic analysis of a drifting buoy in wind, waves and a current was performed, indicating that surface waves do have an important effect on drifting buoy performance.

Supported by: NOAA, Office of Sea Grant,
Grant NA 80-AA-D-00077 (RIE-9) and
NASA Grant NGT 22-014-800.

INVERSE METHODS AND RESULTS FROM THE 1981
OCEAN ACOUSTIC TOMOGRAPHY EXPERIMENT

Bruce D. Cornuelle

Ocean acoustic tomography was proposed in 1978 by Munk and Wunsch as a possible technique for monitoring the evolution of temperature, density, and current fields over large regions. In 1981, the Ocean Tomography Group deployed four 224 Hz acoustic sources and five receivers in an array which fit within a box 300 km. on a side centered on 26°N, 70°W (southwest of Bermuda). The experiment was intended both to demonstrate the practicality of tomography as an observation tool and to extend the understanding of mesoscale evolution in the low-energy region far from the strong Gulf Stream recirculation.

The propagation of 224 Hz sound energy in the ocean can be described as a set of rays travelling from source to receiver, with each ray taking a different path

through the ocean in a vertical plane connecting the source and receiver. The sources transmitted a phase-coded signal which was processed at the receiver to produce a pulse at the time of arrival of the signal. Rays can be distinguished by their different pulse travel times, and these travel times change in response to variations in sound speed and current in the ocean through which the rays passed.

In order to reconstruct the ocean variations from the observed travel time changes, it is necessary to specify models for both the variations and their effect on the travel times. The dependence of travel time on the oceanic sound speed and current fields can be calculated using ray paths traced by computer. The vertical structure of the sound speed and current fields in the ocean were modeled as a combination of Empirical Orthogonal Functions (EOFs) from MODE. The horizontal structure was continuous, but was constrained to have a gaussian covariance with a 100 km e-folding scale. The resulting estimator closely resembles objective mapping as used in meteorology and physical oceanography. The tomographic system has at present only been used to estimate sound speed structure for comparison with the traditional measurements, especially the first two NOAA CTD surveys, but the method provides means for estimating density, temperature or velocity fields, and these will be produced in the future.

The sound speed estimates made using the tomographic system match the traditional measurements to within the associated error bars, and there are several possibilities for improving the signal to noise ratio of the data given structures at small scales, such as in the Gulf Stream, or at large scales, over entire ocean basins. Work is in progress to evaluate the usefulness of tomography as an observation tool in these applications.

Supported by: NSF Grant OCE-80-17791.

FORAMINIFERAL TRACE ELEMENTS:
UPTAKE, DIAGENESIS, AND 100 M.Y.
PALEOCHEMICAL HISTORY

Margaret L. Delaney

Crustal generation rates from 80-110 m.y.b.p. have been suggested to be a factor of two times those at present from geological and geophysical evidence about eustatic sea levels, sea floor magnetic lineations, and plate configurations. Other geophysical studies have suggested that crustal generation rates in the past have not been more than ~20 percent larger than those at present. Faster crustal generation should result in increased chemical fluxes

to the ocean from hydrothermal circulation. The oceanic balance of Li is dominated by hydrothermal input; oceanic Li concentrations in the past could have been as much as a factor of two higher than those at present. This thesis investigates the history of oceanic Li concentrations over the last 100 m.y. in order to add independent geochemical evidence to the debate about sea floor spreading rate variations.

Li/Ca ratios in the calcite shells of planktonic foraminifera are used in this study to determine oceanic Li/Ca ratios. The elemental uptake from seawater and the diagenetic changes in the sediment of foraminiferal calcite were investigated to assess the utility of these shells as such indicators.

Laboratory culture experiments on planktonic foraminifera showed that Li, Sr, and Mg elemental ratios to Ca in foraminiferal shells are directly proportional to seawater ratios. The mean distribution coefficients determined in this study for foraminiferal calcite are: Li, $(5.2 \pm 0.6) \times 10^{-3}$; Sr, 0.16 ± 0.02 ; and Mg, $(0.89 \pm 0.18) \times 10^{-3}$. For Na, the ratios in foraminiferal shells did not vary in response to changes in seawater solution Na/Ca ratios. In other laboratory culture experiments, Cd and Zn distribution coefficients in foraminiferal calcite were measured by the use of radiotracers. The best estimate from these measurements of the Cd distribution coefficient in foraminiferal calcite is 1.8 ± 0.1 , which compares well with previous estimates from measurements on planktonic and benthic foraminifera from sediment core samples. Planktonic foraminifera were cultured in the laboratory at different temperatures to investigate the effect of temperature on the trace elemental composition of foraminiferal shells. From these experiments, the interspecific differences in Sr/Ca, Mg/Ca, and Na/Ca ratios of foraminiferal calcite cannot be due solely to differences in calcification temperatures.

Li/Ca, Sr/Ca, Mg/Ca, and Na/Ca ratios were measured in planktonic foraminiferal shells from five DSDP sites; sample ages extended to 100 m.y. The effect of diagenesis on foraminiferal calcite composition was considered from a simple model comparing the composition of the foraminiferal calcite with that of inorganic calcite in equilibrium with the present-day interstitial water. From these comparisons, some foraminiferal calcite appears to be in inorganic equilibrium with the interstitial water. For other samples, the comparison of the various indicators shows that these foraminiferal shells may be substantially unrecrystallized. Quantification of diagenetic changes in foraminiferal calcite chemistry is hampered by the lack of knowledge about inorganic calcite distribution

coefficients applicable to sedimentary recrystallization reactions. The open system nature of the chemical balances between sediments and interstitial water must be explicitly included in diagenetic models of changes in the chemical composition of foraminiferal calcite. As an example, the effect of recrystallization on oxygen isotopic records of foraminiferal calcite is determined in a model with typical interstitial water oxygen gradients.

From a Li mass balance model and the paleochemical records, the possible variations in crustal generation rates over the last 100 m.y. are evaluated. Alternate global extrapolations of the importance of hydrothermal circulation to oceanic mass balances, temporal changes in river fluxes of Li, the nature of the processes removing Li from the ocean, and the potential of coupled variations in the cycles of Ca and Li are considered with this model. This study concludes that at no time in the past 100 m.y. have oceanic Li concentrations been significantly greater than those at present. This geochemical evidence suggests that hydrothermal fluxes and sea floor spreading rates from 80-100 m.y. were not significantly greater than those at present.

Supported by: NSF Grants OCE 8209362
and OCE 8018665; and Shell Companies Foundation, Inc.

ATMOSPHERIC INTERACTIONS WITH GULF STREAM RINGS

William K. Dewar

Four different problems concerning Gulf Stream Rings are considered. The first deals with the particle trajectories of, and advection-diffusion by, a dynamic model of a Ring. It is found that the streaklines computed from the assumptions that the Ring is a steadily propagating and permanent form structure accurately describe its Lagrangian trajectories. The dispersion field of the Ring produces east-west asymmetries in the streaklines, not contained in earlier kinematic studies, which are consistent with observed surface patterns. In the second problem, we compute the core mixed layer evolution of both warm and cold Rings, and compare them to the background SST, in an effort to explain observed SST cycles of Rings. We demonstrate that warm Rings retain their anomalous surface identity, while cold Rings do not, because of differences in both the local atmospheric states of the Sargasso and the Slope and the typical mixed layer structures appropriate to each. The third and fourth problems concern the forced evolution of Gulf Stream Rings as effected by

atmospheric interaction. First, we compute the forced spin down of a Gulf Stream Ring. The variations in surface stress across the Ring necessary to spin it down are caused by the variations in relative air-sea velocity, of which the stress is a quadratic function. From numerical simulations, we find the forced decay rates are comparable to those inferred from Ring observations. In the final problem, it is suggested that a substantial fraction of meridional Ring migration is a forced response, caused by Ring SST and the temperature dependence of stress. The warm central waters of anticyclonic Rings are regions of enhanced stress, producing upwelling to the north, and downwelling to the south, which shifts the Ring to the south. A similar, southward shift is computed for cyclonic Rings with cold centers, which tends to reconcile their numerically computed propagation with observations.

Supported by: NOAA Contract NA80AA-D-0057; NSF Grant OCE-82-40455; and ONR Contract N00014-79-C-0838.

PROCESSING AND INVERSION OF ARCTIC OCEAN REFRACTION DATA

Gregory L. Duckworth

Several seismic refraction and long-range propagation experiments were carried out during the Fram 2 experiment in the Pole Abyssal Plain of the Eastern Arctic Ocean (May 1980). This thesis concerns the digital signal processing and inversion methods used to analyze and interpret these data for crustal and oceanic structure, and the results of these analyses. Examined are: i) methods for obtaining the short aperture plane wave decomposition of the data subject to the experimental limitations imposed by the arctic environment, and ii) methods for inverting these decompositions for oceanic and crustal structure.

For the plane wave decomposition, conventional time and frequency domain array processing techniques for velocity analysis and slant stacking are contrasted to an optimal adaptive technique. The criteria of phase velocity, group velocity (time of arrival), and amplitude estimator bias and stability are discussed.

To recover the structure, several inversion methods for estimation of the ocean/crust velocity structure from the refraction data are examined. In this work, an emphasis is placed on techniques which fully utilize the data obtained in this experiment. The incorporation of the directly estimated delay time, $(\tau)(p)$, and offset, $x(p)$, data as a function of horizontal slowness, P , is carried out. In addition, the high processing gain of the data adap-

tive short aperture slant-stacking technique allows the inclusion of late breaking multiples in the inversions. A primary component of this work deals with the adaptation of a velocity/depth migration technique to the sparsely shot lines available from the arctic.

In addition to the refraction experiments, long-range (341 km) acoustic propagation data are used for the determination of water column and near-bottom sediment velocity structure. The use of both normal mode and WKBJ theory allows development of inversion algorithms for the dispersive water column arrivals and bottom interacting multiples which tightly constrain these velocity structures. The methods allow structural determination from single-shot data as long as the lateral homogeneity of the structure to be examined is sufficient.

Using the techniques developed, refraction and long-range propagation data taken on the Fram 2 experiment are analyzed. These results are some of the most comprehensive determinations of crustal structure in the Fram basin of the Arctic Ocean near the Morris Jessup Rise.

Supported by: ONR Contracts N00014-77-C-0266, N00014-80-C-0161, N00014-81-C-0161 and N00014-82-C-0161; and ARPA Contract N 00014-81-K-0742.

A CONTINENTAL SHELF BOTTOM BOUNDARY LAYER MODEL: THE EFFECTS OF WAVES, CURRENTS, AND A MOVEABLE BED

Scott M. Glenn

A simple model for the bottom boundary layer on the continental shelf is presented. The governing equations are developed for a stratified, turbulent Ekman layer in a combined wave and current flow over a moveable sediment bed. An eddy diffusivity closure scheme that includes the effect of suspended sediment, temperature, and salinity induced stratification on the vertical turbulent diffusion of mass and momentum couples the resulting unsteady conservation equations for fluid momentum, fluid mass, and suspended sediment mass. The wave velocity, current velocity, and suspended sediment concentration profiles predicted by the simultaneous solution of the conservation equations require the physical bottom roughness and a sediment reference concentration to be specified as boundary conditions. The physical bottom roughness associated with biologically generated bedforms, wave generated ripples, and near bed sediment transport are calculated as functions of the flow and sediment conditions. Using expressions for the height of sediment transporting layer and sediment velocity, an expression for the sediment refer-

ence concentration is developed by matching laboratory measurements of sediment transport rates in oscillatory flow. The model predicts that the bottom flow field is highly dependent on (1) the nonlinear wave and current interaction, which increases the boundary shear stress and enhances vertical turbulent diffusion, (2) the effect of the boundary shear stress on a moveable sediment bed, which determines the physical bottom roughness and the amount of sediment in suspension, and (3) the effect of stable stratification, which inhibits vertical turbulent transport and couples the flow to the suspended sediment and fluid density profiles. The validity of the theoretical approach is supported by model predictions that are in excellent agreement with high quality data collected during two continental shelf bottom boundary layer experiments for a wide range of flow and bottom conditions.

Supported by: The American Gas Association Project PR-153-126; NSF Grant OCE-8014930; and NOAA Office of Sea Grant Contracts NA-79AA-D-00101 and NA 79AA-D-00102.

PURIFICATION AND CHARACTERIZATION
OF THE HEPATIC MICROSOMAL MONOOXYGENASE
SYSTEM FROM THE COASTAL MARINE FISH
Stenotomus Chrysops

Alan V. Klotz

Three cytochrome P-450 forms were highly purified (8-12 nmol/mg) from the hepatic microsomes of the untreated coastal marine fish Stenotomus chrysops (scup) by detergent solubilization and column chromatography. Scup cytochromes P-450A, P-450B, and P-450E (labeled in order of elution from the first ion exchange column) had distinct spectroscopic properties, substrate profiles and minimum molecular weights on SDS-polyacrylamide gels (5.7, 45.9 and 54.3 K, respectively). The three purified cytochrome P-450 isozymes yielded different peptide maps when digested with α -chymotrypsin of S. aureus V8 protease. An additional hemoprotein fraction called cytochrome P-450 fraction D was also resolved and partially purified. This cytochrome P-450 preparation was characterized by a red shift in the CO-ligated, reduced difference spectrum with a chromophore near 451 nm. The scup NADPH-cytochrome P-450 reductase was purified to electrophoretic homogeneity ($M_r = 82.6$ K), had a specific activity of 45-60 U/mg with cytochrome c, and contained both FAD and FMN.

Scup cytochrome P-450E ($M_r = 54.3$ K) is the major aryl hydrocarbon hydroxylating form in untreated hepatic microsomes as judged by both its abundance (30-50 percent

of the total resolved cytochromes P-450) and catalytic activity with benzo[a]pyrene (turnover number = 0.9 nmol product/nmol P-450/min). Further, reconstituted cytochrome P-450E was inhibited 70-80 percent by 100 μ M 7,8-benzoflavone in catalytic assays, similar to the 80-90 percent inhibition of benzo[a]pyrene hydroxylase in microsomal incubations. Analysis of benzo[a]pyrene products derived from reconstitutions of purified cytochrome P-450E revealed that greater than 50 percent of the oxidation occurred at benzo-ring positions, like the product profile observed in incubations with microsomes. The molecular weight of the purified cytochrome P-450E is identical to the major microsomal hemoprotein induced by 3-methylcholanthrene pretreatment, suggesting cytochrome P-450E is the major aromatic hydrocarbon-inducible cytochrome P-450 form in scup. Rabbit antisera raised against purified scup cytochrome P-450E reacts in Ouchterlony diffusion analysis with cytochrome P-450E antigenic determinants in microsomes but not purified cytochrome P-450A. Further, the antisera cross-reacts with an apparent 3-methylcholanthrene-inducible cytochrome P-450 isozyme purified from trout liver (TLM-4a; Williams and Buhler, Comp. Biochem. Physiol. 75C: 25-32, 1983), giving a pattern of fusion without visible spurring in Ouchterlony analysis. These observations imply common antigenic determinants for the apparent major 3-methylcholanthrene-inducible cytochrome P-450 forms in trout and scup.

Monooxygenase reconstitution experiments indicated that purified scup cytochrome P-450A actively hydroxylated testosterone at the 6 β position (turnover number = 0.8 nmol/min/nmol cytochrome P-450). Reconstituted cytochrome P-450B oxidized testosterone at two different sites tentatively identified as the 2-a and 15-a positions (total turnover number = 0.07 min⁻¹). Cytochrome P-450 fraction D produced several metabolites upon reconstitution (sum turnover number = 0.2 min⁻¹) including two chromatographically similar to 16a - and 16 β -hydroxytestosterone. Reconstituted cytochrome P-450E was a poor catalyst of testosterone hydroxylase but the only detectable product was 6 β -hydroxytestosterone (turnover number = 0.04 min⁻¹). However, besides benzo[a]pyrene, reconstituted cytochrome P-450E was active with 7-ethoxycoumarin, acetanilide and 7,8-benzoflavone as a substrates.

Addition of purified scup cytochrome b₅ to mono-oxygenase reconstitutions had a stimulatory effect on some catalytic rates. The magnitude of the cytochrome b₅ stimulation depended on the P-450 isozyme and the substrate used in the reconstitution; cytochrome P-450A was generally influenced by the presence of cytochrome

b₅. This rate stimulation was greater than the effect obtained with purified rabbit liver cytochrome b₅. In an extreme example, cytochrome P-450E was completely inactive in reconstitutions of 7-ethoxyresorufin O-deethylase (an activity associated in microsomes with aromatic hydrocarbon induction) in the presence or absence of rabbit cytochrome b₅, but the addition of scup cytochrome b₅ to the reconstitution led to an observed O-deethylation rate of 7.0 min⁻¹. It is uncertain whether these cytochrome b₅ effects are exhibited in microsomes or in vivo but the stimulation in reconstitutions appears to be important in the evaluation of catalytic activity with purified isozymes.

Supported by: Woods Hole Oceanographic Institution; WHOI Coastal Research Center Grant 67.08; NSF Grant OCE 80-18569; and NIH Grant GM 21643.

STABLE ISOTOPE GEOCHEMISTRY OF NITROGEN IN MARINE PARTICULATES

Susan M. Libes

Isotope studies of nitrogen and carbon were undertaken to investigate the fate of particulate organic matter (POM) during its residence in the water column and after deposition on the seafloor. The processes focused on were water-column transformation and sedimentary diagenesis. Sampling sites were chosen to provide POM subject to different specific mineralization processes (nitrification, denitrification, and sulfate reduction), different lengths of water column (duration of the mineralization process), and differences in the size of the organic-matter flux. The $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$ of plankton, POM, and sediments from several oceanic sites were related to biological and hydrographic processes identified from nutrient, temperature, and salinity profiles. This was done to determine what effect these processes have on the $\delta^{15}\text{N}$ of POM. Four stations were studied in the upwelling area off the coast of Peru and one station was studied in the Gulf of Maine.

Important factors controlling the $\delta^{15}\text{N}$ of plankton appear to be the concentration and $\delta^{15}\text{N}$ of nitrate in the surface waters and the relative zooplankton and phytoplankton abundances. Plankton from the Peru Upwelling area are enriched in ^{15}N as compared to plankton from other parts of the world's oceans where denitrification is absent. This enrichment may be due to the assimilation of ^{15}N -enriched nitrate, produced by the selective reduction of ^{14}N during denitrification. Zooplankton are 3 to 4 o/oo enriched in ^{15}N as compared with phytoplankton. Production

of ^{14}N -enriched fecal pellets is suggested as a mechanism for this trophic enrichment.

In the surface waters, the $\delta^{15}\text{N}$ of POM is similar to that of plankton. In the Peru Upwelling Area, the $\delta^{15}\text{N}$ of POM from the oxygen-deficient waters decreases with increasing depth. In the Gulf of Maine, below the euphotic zone in the oxic deep waters, the $\delta^{15}\text{N}$ of POM increases with increasing depth. The difference in isotopic alteration may be due to the effect of different redox conditions on the mechanism and sequence by which specific organic nitrogen compounds, variably enriched in ^{15}N , undergo degradation. Furthermore, bacterial growth on nitrogen-poor particles in the deep waters of the Peru Upwelling Area may contribute to the low $\delta^{15}\text{N}$ of POM.

In contrast to the large range in $\delta^{15}\text{N}$ (-2 to +17 o/oo) of the POM, the range of $\delta^{15}\text{N}$ in the sediments is small (+5 to +9 o/oo). Within a core, the average variation in $\delta^{15}\text{N}$ was only 1.8 o/oo. Temporal variability in the $\delta^{15}\text{N}$ of sedimenting POM and benthic activity appear to be important in determining the $\delta^{15}\text{N}$ of the sediments. The large changes in POM concentration and isotope content at the sediment/water interface as compared with the more constant values found down-core, suggest that processes occurring at the sediment/water interface are critical, although bioturbation may also be important in determining the $\delta^{15}\text{N}$ of oxic sediments. If diagenesis causes a significant loss of organic matter, profiles of organic carbon and nitrogen contents should show decreases with increasing depth and C/N ratios should increase with increasing depth (Reimers, 1981). Since none of the sedimentary profiles exhibited such trends, it is concluded that diagenesis was insufficient to erase the percent carbon, nitrogen and C/N ratio signatures generated by the POM flux and alterations at the sediment/water interface. Temporal variability in the $\delta^{15}\text{N}$ of bottom-water POM may be caused by changes in deep-water currents which transport POM horizontally and to changes in bacterial and possibly other biological activity in the water column.

This thesis work suggests that $\delta^{15}\text{N}$ may be a useful tool in studying the geochemistry of POM in the marine environment. In addition, this research has shown that interpretation of the sedimentary ^{15}N record must include consideration of isotopic alteration associated with bacterial remineralization of POM and benthic activity.

Supported by: NSF Grant OCE-8024442; The Andrew W. Mellon Foundation; and the WHOI Coastal Research Center.

EFFECTS OF SIZE, AGE AND PHOTOPERIOD
ON HYPOOSMOREGULATION IN BROOK TROUT,
Salvelinus fontinalis

Stephen D. McCormick

Brook Trout (Salvelinus fontinalis) raised from eggs under two photoperiod and two feeding regimes were tested for physiological changes preparatory for transition from freshwater to seawater. Size, age, growth rate, photoperiod, and diel rhythms were examined for possible influences on plasma osmolality, $[Na^+]$, $[Cl^-]$, $[K^+]$, $[Mg^{2+}]$, thyroxine concentration, hematocrit, and gill Na^+, K^+ -ATPase activity of brook trout in freshwater. Significant diel cycles were found in plasma osmolality, $[Na^+]$ and thyroxine concentration. Significant size and/or age related changes occurred for plasma osmolality, $[Na^+]$, $[K^+]$ and hematocrit, but could explain little of their total variation ($0.02 < r^2 < 0.18$). A sexually dimorphic response to photoperiod was observed in hematocrit for both mature and immature fish, with hematocrit of mature females declining in autumn and hematocrit of immature males increasing in autumn. Gill Na^+, K^+ -ATPase activity did not respond to photoperiod or feeding treatment and showed no change with size or age. Plasma thyroxine levels responded to feeding and photoperiod treatment. There was a significant correlation between the percent mean difference in plasma thyroxine and the mean difference in growth rate between high and low feed fish ($r^2 = 0.51$), suggesting a relationship between thyroxine and growth.

In 11 experiments over 1.5 years, brook trout were gradually exposed to 32 ppt seawater for 20 d to investigate the ontogeny of salinity tolerance. A single experiment examined daily changes in plasma osmolality, $[Na^+]$, $[Cl^-]$, $[K^+]$, $[Mg^{2+}]$, thyroxine, hematocrit and gill Na^+, K^+ -ATPase during adaptation to 10 ppt, 20 ppt and 32 ppt. Size was the primary determinant of seawater survival ($r^2 = 0.77$); the effect of size on seawater survival slowed after fish reached a fork length of 14 cm. The effect of age on seawater survival ($r^2 = 0.65$) was through its covariance with size. Photoperiod affected seawater survival only through its influence on the timing of male maturation, which decreased salinity tolerance. Hypoosmoregulation of plasma osmolality, $[Na^+]$, $[Cl^-]$, $[K^+]$, $[Mg^{2+}]$, and hematocrit increased linearly with size over the entire range of sizes (6-32 cm). Gill Na^+, K^+ -ATPase activity after 20 d in seawater decreased with increasing size of brook trout, possibly reflecting decreased demand for active ion transport in larger fish. Plasma thyroxine concentrations generally declined in seawater, but no de-

finite role of this hormone in seawater adaptation was found. Size dependent survival and osmoregulatory ability of brook trout is compared to other salmonids and a conceptual model is developed.

Decreased salinity tolerance and hypoosmoregulatory ability was found in mature male brook trout and was not found in females or immature males. Lowered salinity tolerance of adult males becomes acute during autumn photoperiod when normal spawning occurs. Plasma $[Cl^-]$, $[Mg^{2+}]$, osmolality and hematocrit are significantly higher in mature males after transfer to seawater, relative to mature females. It is postulated that reduced adult male hypoosmoregulatory ability explains skewed sex ratios in anadromous populations, limits the extent of anadromy, and was a significant phase in the evolution of extended salmonid migration.

Anadromous brook trout of Riviere a la Truite, Quebec, were examined for physiological changes associated with salmonid smoltification, and compared to non-anadromous brook trout of the Matamek River. There were no significant differences in plasma thyroxine concentration, gill Na^+, K^+ -ATPase activity, hematocrit or osmoregulatory ability of anadromous and non-anadromous brook trout. Moisture content was significantly different between fish from the two river systems, but had the same pattern of declining moisture content as summer progressed. Silver coloration of brook trout in Riviere a la Truite was significantly associated with larger fish and higher gill Na^+, K^+ -ATPase activity, but not with changes in plasma thyroxine, moisture content, hematocrit or condition factor. Silver coloration was absent in Matamek River brook trout. Brook trout at high salinity estuarine sites had significantly greater gill Na^+, K^+ -ATPase activity and hypoosmoregulatory ability than brook trout at low salinity sites. Atlantic salmon (Salmo salar) in high salinity estuarine site had significantly higher plasma thyroxine and gill Na^+, K^+ -ATPase activity than brook trout. The results indicate that smoltification is relatively undeveloped in brook trout and that estuarine residence is important in salinity adaptation and eventual seaward migration.

Supported by: NOAA Office of Sea Grant.

LATE PALEOGENE (EOCENE TO OLIGOCENE)
PALEOCEANOGRAPHY OF THE NORTHERN NORTH
ATLANTIC

Kenneth G. Miller

Seismic stratigraphic evidence from the western and northern North Atlantic indi-

cates that a major change in abyssal circulation occurred in the latest Eocene to earliest Oligocene. In the northern North Atlantic, the widely-distributed reflector R4 correlates with an unconformity that can be traced to its correlative conformity near the top of the Eocene. This horizon reflects a change from weakly circulating (Eocene) to vigorously circulating bottom water (early Oligocene). Sediment distribution patterns provide evidence for strong contour-following bottom water flow beginning at reflector R4 time; this suggests a northern source for this bottom water, probably from the Arctic via the Norwegian-Greenland Sea and Faeroe-Shetland Channel. Erosion and current-controlled sedimentation continued through the Oligocene; however, above reflector R3 (middle to upper Oligocene), the intensity of abyssal currents decreased. Above reflector R2 (upper lower Miocene) current-controlled sedimentation became more coherent and a major phase of sedimentary drift development began. This resulted from further reduction in speeds and stabilization of abyssal currents.

Paleontological and stable isotopic data support these interpretations. In the Bay of Biscay/Goban Spur regions, a major $\delta^{18}O$ increase began at ~38 Ma (late Eocene), culminating in a rapid (< 0.5 my) increase in $\delta^{18}O$ just above the Eocene/Oligocene boundary (~36.5 Ma). A rapid $\delta^{13}C$ increase also occurs at ~36.5 Ma in these sites. Major changes in benthic foraminiferal assemblages also occurred between the middle Eocene and the earliest Oligocene: 1) In the Labrador Sea, a predominantly agglutinated assemblage was replaced by a calcareous assemblage between the middle Eocene and early Oligocene; 2) In the abyssal (> 3 km) Bay of Biscay, an indigenous Eocene calcareous fauna including Nuttallides truempyi, Clinapertina spp., Abyssammina spp., Aragonia spp., and Alabamina dissonata became extinct between the middle Eocene and earliest Oligocene; 3) In shallower sites (< 3 km paleodepth) throughout the Atlantic, a Nuttallides truempyi-dominated assemblage was replaced by a Globocassidulina subglobosa-Gyroidinoides-Cibicidoides ungerianus-Oridorsalis assemblage in the early late Eocene (~40-38.5 Ma). These faunal and isotopic changes represent the transition from warm, old, corrosive Eocene bottom waters to colder, younger (lower CO_2 and higher pH, hence less corrosive) early Oligocene bottom waters.

Supported by: ONR Contract N00014-79-C-0071; Phillips Petroleum Company; and WHOI.

THE NUMERICAL SYNTHESIS AND INVERSION OF ACOUSTIC FIELDS USING THE HANKEL TRANSFORM WITH APPLICATION TO THE ESTIMATION OF THE PLANE WAVE REFLECTION COEFFICIENT OF THE OCEAN BOTTOM

Douglas R. Mook

The plane wave reflection coefficient is an important geometry independent means of specifying the acoustic response of a horizontally stratified ocean bottom. It is an integral step in the inversion of acoustic field measurements to obtain parameters of the bottom and it is used to characterize an environment for purposes of acoustic imaging. This thesis studies both the generation of synthetic pressure fields through the plane wave reflection coefficient and the inversion of measured pressure fields to estimate the plane wave reflection coefficient. These are related through the Sommerfeld integral which is in the form of a Hankel transform. The Hankel transform is extensively studied in this thesis and both theoretical properties and numerical implementations are considered. These results have broad applications. When we apply them to the generation of synthetic data, we obtain hybrid numerical-analytical algorithms which provide extremely accurate synthetic fields without sacrificing computational speed. These algorithms can accurately incorporate the effects of trapped modes guided by slow speed layers in the bottom. We also apply these tools to study the inversion of measured pressure field data for the plane wave reflection coefficient. We address practical issues associated with the inversion procedure including removal of the source field, sampling, field measurements over a finite range, and uncontrolled variations in source-height. A phase unwrapping and associated interpolation scheme is developed to handle improperly spaced data.

A preliminary inversion of real pressure field data is performed. In parallel, an inversion of a synthetically generated field for similar bottom parameters is also performed and the results of processing the real and synthetic data are compared. The estimate for the depth dependent Green's function obtained from the real data shares many features with the depth dependent Green's function estimated from the synthetic data, suggesting that the total inversion to obtain the plane wave reflection coefficient will soon be possible. Errors in the present estimate of the plane wave reflection coefficient are associated with uncontrolled source-height variations during the acquisition of data.

Supported by: ONR Contracts N00014-C-0152, N00014-81-K-0742 and NR 049-506; and NSF Grant ECS 80-07102.

FLOW AND SKIN FRICTION
OVER NATURAL ROUGH BEDS

Christopher Paola

When a boundary layer develops over a bed that is hydrodynamically rough at a length scale or scales larger than the grain size (a macrorough bed), as is usually the case where bed forms are present, it is necessary to distinguish among total boundary shear stress and its components, form drag and spatially averaged skin friction. It is known that the mean-velocity field reflects the composite boundary shear stress. Above about one roughness height above the tops of the roughness elements, the velocity does not vary horizontally. Its vertical profile is semilogarithmic and scales with the total friction velocity u_{τ} and total roughness length z_{0t} . This region is here called the integrated logarithmic layer (ILL). Below the ILL the velocity varies horizontally in response to the irregular boundary; this region is called the surface layer.

In the first of two sets of experiments reported here, skin-friction measurements were made with an array of flush-mounted hot films at four points on the stoss slope of one of a field of two-dimensional immobile current ripples. Total boundary shear stress was also measured, as were mean-velocity profiles in the ILL and the surface layer. The ILL behaves as described above. Although surface-layer velocity profiles are semilogarithmic, their semilogarithmic slope is not proportional to the local skin-friction velocity, so they do not locally obey the law of the wall. Rather, the velocity field can be decomposed into a spatially averaged rotational component and a local inviscid perturbation. The measured skin-friction field is consistent with a simple model for sediment transport over the bed forms except near reattachment, where the fluctuating skin friction is important. The data are also consistent with the drag-partition theories of Einstein and Barbarossa (1952) and Englund (1966). Normalized skin-friction spectra do not vary with streamwise position but do vary with Reynolds number; skin-friction probability density functions show significant increases in skewness and kurtosis near reattachment but do not vary strongly with Reynolds number.

In the second set of experiments the skin-friction vector field was measured around isolated hemispheres, with model sedimentary tails one and four obstacle heights long and without tails. The measured skin-friction fields are not consistent with deposition along the obstacle-flow centerline downstream of reattachment, which occurs about two obstacle heights downstream of the trailing edge of the hem-

isphere. This applies for local bed-load erosion and deposition and for general deflation of the bed and is not substantially altered by the presence of either tail. Measurements were also made of skin friction, total boundary shear stress and ILL velocity profiles over h,B-rough arrays of hemispheres with and without tails four roughness heights long, at two areal densities. The skin-friction field in the denser array is significantly distorted from that around an isolated element. The measured skin friction in both arrays is significantly greater than that given by the drag-partition formula proposed by Wooding et al. (1973). The roughness length z_{0t} for both densities is not changed by addition of the tails.

Supported by: NSF Grant OCE 77-20437
and ONR Contract N00014-80-C-0273.

A STUDY OF THE SEISMIC STRUCTURE
OF UPPER OCEANIC CRUST
USING WIDE-ANGLE REFLECTIONS

Kristin M. M. Rohr

The lateral homogeneity of oceanic crust on the scale of a seismic experiment is a condition that most methods of seismic interpretation depend on. Whether this condition is in fact true is largely unknown and only recently have efforts been made to test this hypothesis. This thesis is part of that effort and is focused on determining with as much resolution as possible the seismic structure of upper oceanic crust, i.e., Layer 1 and the uppermost part of Layer 2. This portion of the crust is of interest because of the effect of the sediment-basement interface on the transmission and conversion of seismic energy and because of the possibility of detecting lateral heterogeneities in upper Layer 2 caused by faulting, hydrothermal circulation, etc. The data employed are a set of wide-angle reflections from oceanic crust 130 m.y. old in the western North Atlantic Ocean southwest of Bermuda. First, the sedimentary structure is determined by stacking the data along hyperbolae and interpreting the stacking velocities and two-way normal incidence travel-times for interval velocities. This method has not been applied to deep sea marine data before; it gives a more detailed velocity structure of the sediments than does a traditional study of the basement reflections' travel times. Second, the same data are mapped into tau-p space in order to measure the velocity gradient in oceanic basement; unfortunately the scatter in the tau-p picks caused by the topography of the basement reflector combine with the properties of the tau-sum inversion to make such a

measurement impossible. Third, the amplitudes of the basement reflections observed on three seismic lines are modeled by synthetic seismograms; each can be matched by velocity-depth models which contain a transition zone between the sediments and the basement. The different thicknesses of this transition zone near the three receivers is an indication that the top few hundred meters of Layer 2 are laterally heterogeneous on a scale of 3 to 8 km.

Supported by: NSF Grant OCE 79-09464.

WAVE-INDUCED TURBULENT FLOW
NEAR A ROUGH BED: IMPLICATIONS OF A TIME-
VARYING EDDY VISCOSITY

John H. Trowbridge

Approximate equations governing the turbulent flow produced near a fixed, rough bed by plane, progressive, monochromatic, small amplitude surface waves are derived in detail. The equations involve two approximations: the boundary layer approximation and the Stokes expansion in powers of the wave steepness. An order-of-magnitude analysis of the flow very near the bed provides the basis for two time-varying eddy viscosity models with different vertical structures. In both models, the temporal variation of the viscosity is related to the first few Fourier components of a shear velocity based on the instantaneous bed shear stress. Approximate analytical expressions are derived for the near-bed flow produced by first and second order Stokes waves.

Comparison of the first-order solution with available measurements indicates that the theoretical models are physically sound. At first order, the solution is insensitive to the assumed vertical structure of the time-varying part of the viscosity, and temporal variation of the viscosity has a noticeable but relatively small effect on quantities of practical interest. The most dramatic result of the analysis is a predicted reversal of the second-order steady streaming produced by long waves. The prediction of the steady streaming is quite sensitive to the assumed vertical structure of the time-varying part of the eddy viscosity. Qualitative support for the reversal of the steady streaming is found in the available experimental data. In contrast to the steady streaming, the second-order bed shear stress is found to be insensitive to assumptions regarding the temporal variation of the eddy viscosity.

Supported by: NSF Grant OCE 80-14938;
NOAA Office of Sea Grant
NA81AA-D-00069, Project R/C-22; and
WHOI.

THE OCEANOGRAPHIC AND GEOIDAL COMPONENTS
OF SEA SURFACE TOPOGRAPHY

Victor Zlotnicki

Altimetric, gravimetric and oceanographic data over the North Atlantic are combined, using techniques of optimum estimation, to infer the surface expression of the time averaged circulation (ζ) and to estimate the marine geoid (γ), both in the wavelength band 100 km-2000 km.

Optimum inverse methods in geophysics are reviewed. They are then used to analyze the estimation of the geoid from gravity data, emphasizing the wavenumber spectrum of resolution functions. It is found that accurate bandpassed versions of the geoid can be recovered from restricted data sets.

The accuracy and distribution of publicly available gravity data are shown to define an estimate γ whose expected errors, $\sigma\gamma$, range between 30 and 260 cm, assuming the Wagner and Columbo (1978) spectrum describes the average geoid behavior. The $\sigma\gamma$ underestimate the actual differences between γ and an altimetric surface (s) derived from Seasat, but the spatial variation of $\sigma\gamma$ follows closely the differences $s-\gamma$. The discrepancy is attributable to a partial failure of the spectral model at short wavelengths.

The differences $s-\gamma$ are dominated by geoid error that masks much of the signal (ζ). The main North Atlantic gyre emerges clearly only after the $\sigma\gamma$ and the simplest model for (ζ), as a spatially uncorrelated process with $(30 \text{ cm})^2$ variance, are taken into account. To obtain a corrected geoid, a hydrographic estimate of (ζ) is combined with s and γ , and their expected errors.

Supported by: NASA Grant NAG 6-9.

AUTHOR INDEX

	Page No.		Page No.
Aber, John D.	B-10	Cacchione, David A.	OE-4,5,6
Adamson, A.C.	C-7	Calabrese, Anthony C.	B-4
Agrawal, Yogesh C.	OE-8,9,11	Calloway, C. Bradford	B-19
Albarede, F.	OE-8	Cannon, Glenn A.	PO-3
Almogi-Labin, A.	C-3	Canuel, E.A.	C-7
Alt, J.C.	C-7	Capuzzo, Judith M.	B-14,15,16,17,26
Anderson, Donald M.	B-20,22,23,29	Carey, A.E.	C-16
Anderson, R.N.	C-7	Carey, Francis G.	B-3
*Asper, Vernon L.	OE-11	*Caron, David A.	B-26,27
Aubrey, David G.	GG-22,23,26,27	Casey, J.F.	GG-6,20
Backus, Richard H.	PO-14	Casso, Susan A.	C-16,18
Bacon, Michael P.	C-14,15	Caswell, Hal	B-19,20,23,24
Bailey, Conner	MP-3,4	Chapman, David C.	PO-5,17
Baker, Karen S.	PO-14	Charlou, J.L.	OE-8
Ballard, Robert D.	OE-7,8	Cheminee, J.L.	OE-8
Bandel, K.	C-3	Choukroune, P.	GG-4;OE-8
Barbour, R. Lorraine	PO-25	Cicin-Sain, Biliana	MP-8
+Bé, Allan W.H.	B-27	Clague, D.	C-15
Beardsley, Robert C.	PO-3,4,5,7,21,23	Clay, Peter R.	OE-10,11
Becker, K.	C-7	Coats, D. Wayne	B-23
Beg, M.A.A.	GG-25	Cochran, J.K.	C-6,14
Behringer, David W.	PO-23	Cole, Jonathan J.	GG-23
Berggren, William A.	GG-3,17,18	*Collins, John A.	GG-6,11;OE-11
Berteaux, H.O.	OE-9,10,12	Comita, Paul B.	C-3
*Binder, Brian J.	B-22	Connor, Michael	MP-8
Binder, Robert L.	B-3	Corliss, Bruce H.	GG-7
Birdsall, T.G.	OE-14	*Cornuelle, Bruce D.	GS-3
Blackwelder, Patricia L.	PO-14	Costello, C.E.	C-3
Boardman, David C.	PO-22	Cowles, Timothy J.	B-16,27;PO-14
Bonatti, E.	GG-6	Crane, K.	GG-4,6
Bonde, L.W.	OE-10	Cronin, Carolyn	C-5
Borges, Mark D.	PO-12	Csanady, Gabriel T.	PO-5,6,17
Boudreau, R.D.	C-15	Curry, William B.	GG-16,17
Boulegue, J.	OE-8	Dacey, John W.H.	B-11,12
Bowen, Robert E.	MP-4,5	Davis, Linda H.	B-29
Bowen, V.T.	C-16	Davis, Paul G.	B-27
Bowin, Carl	C-7;GG-9,10,11	Dean, Jerome P.	PO-21,22
Bradley, Albert M.	OE-10,11;PO-7	*DeBaar, Hein J.W.	C-14
Bratt, Steven R.	GG-11	DeBoer, J.A.	B-21
Bray, Nancy A.	PO-13	Debusk, Thomas A.	B-9
Brewer, Peter G.	C-8	Degens, Egon T.	C-4,5;GG-23
Brink, Kenneth H.	PO-3	*Delaney, Margaret L.	GS-3
Briscoe, Melbourne G.	PO-21,23,25,26	DeMaster, D.J.	GG-24
Broadus, James M.	MP-4,5	Denham, Charles R.	GG-3,8,12,13
Broecker, W.S.	C-8	Dennett, Mark R.	B-21
Brown, Otis B.	PO-14	Derby, Jennifer G. Smith	B-16
Brown, Wendell S.	PO-23	Desaubies, Yves	OE-13,15
Bruce, John G.	PO-13	Deser, Clara	PO-23
Bryan, W.B.	GG-19,21	Deuser, Werner G.	C-3,4,5,10,14
Bryant, Donald	B-12	*Dewar, William K.	GS-4;PO-17
Bryden, Harry L.	PO-9,14	Dick, Henry J.B.	GG-7,12,20,21
Buchsbaum, Robert	B-12	Dietz, K.	MP-7
Buckley, D.E.	GG-26	Dillon, D.B.	OE-10
Burke, J.C.	C-18	Doherty, Kenneth W.	OE-11
*Burke, William J.	GS-3	Drake, David E.	OE-4,5,6
Burt, Wayne V.	B-15	Druffel, Ellen R.M.	C-9
Butenko, Jorge	GG-3	*Duckworth, Gregory L.	GS-5
Butman, Bradford	PO-4,5,23	Duedall, Iver W.	B-15

Ebbesmeyer, Curtis C. PO-9
 Edmond, J.M. C-11
 Elthon, D. GG-6,20
 Embley, Robert W. GG-9
 Emeis, K. C-4
 Emery, K.O. MP-10
 Emmermann, R. C-7
 Erlank, A.J. GG-21
 Evans, Robert H. PO-14
 Eyster, Linda S. B-8
 Farmer, Harlow G. GG-8
 Farr, Barbara S. B-26
 Farrington, J.W. C-6,7
 Feuer, A. OE-3
 Fisher, R.L. GG-20
 FitzGerald, Gerry J. B-10
 Flagg, Charles N. PO-6
 Fleer, Alan P. C-14,15;GG-23
 Flynn, John J. GG-17
 Fofonoff, Nick P. PO-9
 Foreman, Kenneth B-12
 Fornari, D. GG-6
 Fox, P.J. GG-4,6
 Francheteau, Jean OE-7,8
 Francis, Erika A. PO-22
 Francis, Margaret M. B-9
 Francis, T.J.G. C-7
 Freadman, Marvin A. B-3,4
 Frey, F.A. GG-21
 Friehe, Carl A. PO-21
 Frisk, George V. OE-13,14
 Fryxell, Greta A. PO-14
 Gagosian, Robert B. C-3,4,5,6,7
 Gallagher, Scott M. B-5
 Gallo, D.G. GG-4,6
 Gamble, John King Jr. MP-5
 Gaudreault, Alaine B-10
 Gibson, Jane B-18
 *Glenn, Scott M. OE-5,6;GS-5
 Glibert, Patricia M. B-21
 Goebel, Camille A. B-14
 Goldman, Joel C. B-21
 *Goud, Margaret R. GG-26
 Gove, Lee OE-7
 Govoni, John J. B-7
 *Graham, D. C-11
 Grant, William D. OE-4,5,6
 Grassle, J. Frederick B-4
 Grosskopf, W.G. GG-27
 Grove, T.L. GG-21
 Hacker, Peter GG-25
 Haidvogel, Dale B. PO-18
 Hamblin, R.E. C-18
 Hamlyn, P. GG-6
 *Hannan, Cheryl Ann B-24
 Haq, B.U. GG-8
 Harding, A.J. GG-15
 Hart, S.R. C-9,15;GG-21
 Hayes, Stanley P. PO-21
 Healey, Michael C. MP-6,7
 Heath, Ron A. PO-14
 Hekinian, R. OE-8
 Hemleben, C. C-3

Henderson-Sellers, Brian PO-18
 Hendry, Ross M. PO-9
 Hennessey, Timothy MP-7
 Hoagland, Porter III MP-5
 Hogg, Nelson G. PO-10
 Holland, William R. PO-10
 Hollister, Charles D. GG-4,9,24
 Honjo, Susumu GG-23,26;OE-11
 Honnorez, J. C-7
 Howarth, Robert W. B-17
 Howes, Brian L. B-11,12
 Hsueh, Ya GG-25
 Hu, Dun-Xin GG-25;PO-3
 Hua, Bach-Lien PO-13
 Huebert, B. C-11
 Huh, Chih-An C-14
 Humphris, Susan E. C-9
 Hunt, J.M. C-11
 Hunt, Mary M. OE-7
 Hurd, David C. GG-26
 Irish, James D. PO-23
 Ittekkot, Venugopalan. C-4,5;GG-23
 Janecky, D.R. C-12
 Jannasch, Holger W. B-18,19
 Jenkins, William J. C-8,10,11,15
 Jickells, T.D. C-10
 Johnson, David A. GG-4,5,17,24
 Johnson, Paul W. B-27
 Jones, Glenn A. GG-5,17
 Joyce, Terrence M. PO-14,15,25
 Kanwisher, John W. B-3
 Karson, Jeffrey A. C-11;GG-4,6,7,20,21
 Kastens, K. GG-4
 Keffer, Thomas PO-10,11,18,25
 Keigwin, L.D. GG-7,8,26
 Kempton, P.D. C-7
 Kent, Dennis V. GG-3,17
 Kester, Dana R. B-15
 Kidd, W.S.F. GG-4,6
 King, Gary M. B-11
 Kinoshita, H. C-7
 Kirk, R.E. OE-12
 Kletter, Doron OE-3
 Kley, D. C-11
 Klitgord, Kim GG-16
 *Klotz, Alan V. B-4,5;GS-6
 Knap, A.H. C-10
 Knapp, George P. PO-23
 Knecht, Robert W. MP-7,8
 Knox, R.R. OE-14
 Koelsch, Donald E. GG-11;OE-11,12
 Koide, M. C-9
 Krishnaswami, S. C-14
 Kulis, David M. B-22
 Kurz, M.D. C-15
 Lahey, William L. MP-8
 Laine, E.P. GG-7
 Lambertsen, Richard B-13
 Lancaster, Bruce A. B-15
 Laverne, C. C-7
 Lawver, Lawrence A. GG-12
 Le, Ken Tang PO-3
 Ledbetter, Michael T. GG-17

Lee, Cindy C-5,6
 leRoex, A.P. GG-21
 Levanon, Nadav OE-3
 Levy, Ellen PO-24
 Liberatore, S. OE-12
 *Libes, Susan M. GS-7
 Limeburner, Richard PO-3,7
 Lindquist, Niels L. B-23
 Linkins, Alan E. B-10
 Liu, S.C. C-11
 Lively, John S. B-20
 Livingston, Hugh D. C-9,16,17
 Loder, John W. PO-4
 Lohmann, G.P. GG-16,18
 Lott, D.E. C-15
 Luyten, James R. PO-11,12
 Lynch, James F. OE-13
 Macdonald, K. GG-4
 Magnell, Bruce A. PO-3
 Malmgren, Bjorn A. GG-18
 Mallette, M. C-11
 Mann, Don R. C-16,17,18
 Mann, Roger B-5,6,7,8,24,25
 Marcus, Nancy H. B-25
 Margolis, S.V. GG-8
 Marra, John PO-22
 Marty, J.C. OE-8
 Mattie, M.G. GG-27
 McCarthy, James J. B-21
 McCartney, Michael S. PO-8
 McCave, Ian N. GG-4,24
 McCormick, M.H. PO-18
 *McCormick, Stephen D. B-6;GS-8
 McDevitt, R.E. OE-10
 McDougall, Trevor Jr. PO-20
 McDowell, Donald M. B-26
 *McElroy, Anne E. B-16
 McFarland, M. C-11
 McKee, Theresa K. PO-12
 Meade, R.H. GG-25
 Melillo, Jerry M. B-9,10
 Mellinger, Ed OE-11,12
 Metzger, K. OE-14
 *Miller, Kenneth G. GS-8
 Miller, Linda B. MP-9
 Miller, Tandy B-10
 Milliman, John D. GG-3,25
 Mills, Carol A. PO-23
 Minster, J.F. OE-8
 Montgomery, W. Linn B-10
 Moody, R. GG-4
 *Mook, Douglas R. OE-14;GS-9
 Moore, Karen E. B-13
 Morin, Rod B-24
 Morris, James T. B-12
 Mottl, M.J. C-7,9,11,12,13
 Mountain, David G. PO-14
 Naiman, Robert J. B-6,9,10,24,26
 Needell, Gerald J. PO-12
 Needham, D. GG-6
 Nelson, Douglas C. B-18
 Nelson, D.M. C-16
 Newmark, R.L. C-7

Nicholson, Desmond V. MP-11
 Nittrouer, C.A. GG-24
 Noble, Marlene PO-3,5
 Ochrymowych, Christina B-13
 Olson, Brenda L. C-5
 Olson, Donald B. PO-14
 Onbe, Takashi B-28
 Oppenheim, Alan V. OE-14
 Owens, W. Brechner PO-9,11,13
 Pang, H. C-3
 *Paola, Christopher GS-10
 Park, P. Kilho B-15
 Pashinski, David J. GG-25;PO-3
 Peal, K.R. OE-12
 Pechenik, Jan A. B-8
 Pederson, Judith B. B-26
 Pedlosky, Joseph PO-18,19,20
 Peters, K.E. C-11
 Pettigrew, Neal R. PO-23
 Pfennig, Norbert B-18
 Pilskaln, Cynthia H. GG-23
 Pinkel, Robert PO-22
 Pires Filho, Ivon d'Almeida MP-9
 Popp, T. OE-10
 Prada, K. OE-12
 Pratt, M.W. C-15
 Price, James F. PO-15,22
 Price, Richard L. MP-10
 Prothero, Donald R. GG-8
 Purcell, Jennifer E. B-9,28
 Purdy, G.M. GG-11,15,16;OE-11
 Qinchon, Xie GG-25
 Quraishie, G.S. GG-25
 Ramp, Steven R. PO-23
 Rasmussen, Kenneth A. GG-24
 Ray, G. Carleton B-14
 Remillard, James F. B-16
 Renard, V. OE-8
 Repeta, Daniel J. C-5
 Rhines, Peter B. C-8;PO-10,17
 Richardson, Philip L. PO-12
 Rietsma, Carol B-12
 *Riley, J.B. OE-9
 Riser, Stephen C. PO-9
 *Rohr, Kristin M.M. GG-12,13;GS-10
 Rona, Peter A. C-11,13
 Rooth, C.G. C-8
 Rosencrantz, Elric GG-20
 *Rosenfeld, Leslie K. PO-24
 Ross, David A. GG-8;MP-10
 Rossby, H. Thomas PO-9,15
 Ruder, Eric GG-23
 Sartori, R. GG-6
 *Sasaki, Glenn C. B-15
 Sayles, F.L. C-16
 Scavia, Donald PO-18
 Scheltema, Rudolf S. B-8
 Schevill, William E. B-13
 Schlitz, Ronald J. PO-14,23
 Schmitt, Raymond W. PO-14,20
 Schmitz, William J., Jr. PO-16
 Schneider, David L. C-17
 Schneider, J.K. C-6

Schouten, Hans GG-13,16
 Schuler, F. OE-12
 Seguret, M. GG-4
 Seyfried, W.E. C-12
 Sgouros, Thomas A. PO-25
 Sharp, Arnold G. OE-15
 Shaw, Ping Tung PO-6
 *Shen, Glen T. C-17
 Shen, Huang-ting GG-25
 Sholkovitz, Edward R. C-17,18
 Shure, Loren GG-12
 Shusterich, Kurt M. MP-11
 Sieburth, John McN. B-27
 Silva, A.J. GG-24
 Sloane, H. GG-6
 Smith, Peter C. PO-14,23
 Smith, Raymond C. PO-14
 Softley, E.J. OE-10
 *Speer, Paul E. GG-23,27
 Spiesberger, J.L. OE-14
 Spindel, R.C. OE-14,15
 Spofford, C.W. OE-14
 Stalcup, Marvel C. PO-15,25
 Stanley, Robert J. PO-23
 Stegeman, John J. B-3,4,5,7
 Stephen, Ralph A. GG-14,15;OE-7
 Stevens, E. Don B-3
 Stoecker, Diane K. B-7,28,29
 Stommel, Henry M. PO-9,11,23
 Stroup, J. GG-6
 Stuart, David W. PO-3
 Sugimoto, Takashige PO-7
 Surprenant, Lolita D. C-16
 Swift, J.H. C-8
 Takahashi, Kozo GG-18,26
 Takahashi, T. C-8
 Talley, Lynne D. PO-8
 Tappa, Eric GG-17
 Tarafa, M.E. C-11
 Tarbell, Susan A. PO-24
 Tate, George B. OE-4,5
 Taylor, Craig D. B-18
 Teal, John M. B-11,12,17
 Terry, William E. Jr. OE-10
 Thompson, Geoffrey C-7,9,11,12,13
 Thomson, John H. PO-12
 Thunell, Robert GG-17
 Thurberg, Frederick P. B-4
 Titus, Marchall GG-20
 Toole, John M. PO-12,21
 Trask, Richard P. PO-25,26
 Trehu, A.M. GG-15
 *Trowbridge, John H. GS-11
 *Trull, T.W. C-6
 Tucholke, Brian E. GG-9
 Tucker, C.S. B-9
 Turekian, K.K. C-14
 Turner, Ruth D. B-19
 Tyler, Mary A. B-23
 Vaccaro, Ralph F. B-20
 Valiela, Ivan B-12
 Van Couvering, John A. GG-3,17
 Vandermeulen, John B-17

Van Leer, John C. PO-3
 Vaucher, Marc E. MP-10
 Vermersch, John A. PO-23
 Vilks, Gustavs GG-26
 Villafuerte, E.D. MP-4
 Volkman, J.K. C-6
 *Von Damm, K. C-11
 Von der Heydt, K. OE-12
 Von Herzen, R.P. C-7
 Voorhis, Arthur D. PO-7,12
 Wakeham, Stuart G. B-12;C-6,7
 Walden, R.G. OE-9,10,12
 Walsh, Chris B-4,5
 Warren, Bruce A. PO-13
 Waterbury, John B. B-18,19
 Watkins, William A. B-13,14
 Watters, David R. MP-11
 Webb, Douglas C. PO-15
 Weinstein, Ehud OE-3,4
 Weiss, Anthony J. OE-3,4
 Weller, Robert A. PO-21,22,23
 Westermeyer, William E. MP-12
 Whelan, J.K. C-11
 White, Robert S. GG-16
 Whitehead, John A., Jr. PO-7,8,20,26
 Whoriskey, F.G. B-21
 Wiebe, Peter H. PO-14
 Williams, Albert J. III OE-5,6
 Williams, Peter M. C-3,9
 Williams, R.T. C-8
 Wilson, John B-12
 Winant, Clinton D. PO-21
 Winget, Clifford L. OE-12
 Wirson, Carl O. B-19
 Wolf, Charlotte C. B-6
 Woodin, Bruce R. B-7
 Worcester, Peter F. OE-14
 Yang, Zuo-sheng GG-25
 Yincin, Y. GG-3
 Yingst, J.Y. GG-24
 Young, William R. PO-17,20
 Zafirou, O. C-11,13
 *Zlotnicki, Victor GS-11
 Zuosheng, Yang GG-25

* student

+ deceased