WHOI-80-56

COPY 1

Woods Hole Oceanographic Institution

WHOI
DOCUMENT
COLLECTION

OBSERVED AND NUMERICALLY SIMULATED
KINETIC ENERGIES FOR MODE EDDIES

by

William J. Schmitz, Jr.
and
W. Brechner Owens

November 1980

TECHNICAL REPORT

*Prepared for the Office of Naval Research under
Contract N00014-76-C-0197; NR 083-400 and for
the International Decade of Ocean Exploration
Office of the National Science Foundation under
Grant OCE 75-03962.*

*Approved for public release; distribution un-
limited.*

WHOI-80-56

OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES
FOR MODE EDDIES

by

William J. Schmitz, Jr.
and
W. Brechner Owens

WOODS HOLE OCEANOGRAPHIC INSTITUTION
Woods Hole, Massachusetts 02543

November 1980

TECHNICAL REPORT

Prepared for the Office of Naval Research under Contract N00014-76-C-0197; NR 083-400 and for the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE 75-03962.

Reproduction in whole or in part is permitted for any purpose of the United States Government. In citing this report in a bibliography, the reference given should be to "Journal of Physical Oceanography, Vol. 9, No. 6, November 1979, pp. 1294-1297".

Approved for public release; distribution unlimited.

Approved for Distribution: Valentine Worthington, Chairman
Department of Physical Oceanography

Observed and Numerically Simulated Kinetic Energies for MODE Eddies

WILLIAM J. SCHMITZ, JR., AND W. BRECHNER OWENS

Woods Hole Oceanographic Institution, Woods Hole, MA 02543

30 March 1979 and 6 August 1979

ABSTRACT

It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the comparison is indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.

1. Introduction

A variety of numerical experiments with periodic horizontal boundary conditions on a limited oceanic area have been run as initial value (spindown) calculations, with results analyzed over a (somewhat arbitrary) time interval after a preliminary period of adjustment [for a review see Rhines (1977)]. The six-layer quasi-geostrophic model developed by Bretherton and Karweit (1975; see also Bretherton, 1975) was applied to the MODE area (near 28°N, 70°W), and statistical results compared with the data there by Owens (1975) and by Owens and Bretherton (1978). Owens and Bretherton (1978) stated that their model (i) successfully predicted the time scales, (ii) but underpredicted the baroclinic (kinetic) energy for eddies in the MODE area. In other words (Owens and Bretherton, 1978; p. 8), point (ii) is that the model results appeared to be significantly more barotropic than the data. Our objectives here are to reexamine this set of results using a more complete data set, and from a new point of view, paying particular attention to the dependence of the vertical distribution of eddy kinetic energy on frequency range.

2. Procedure

Time series of 200 days duration, sampled every quarter-day, each with least-square fits of the mean and linear trend removed, were formed at each of the model's horizontal grid points. Discrete Fourier transforms (DFT's) were then computed for each grid point and averaged over all of the 1024 model grid points for spectral estimates (Owens, 1975; pp. 53 and 57). For a 200-day time series sampled at quarter-day intervals, the DFT assigns energy to frequencies of $n/200$ cpd (cycles per day), $n = 0, 1, \dots, 400$; each at the center of a frequency band of width $(n/200 - 1/400)$ to $(n/200 + 1/400)$ cpd, except for $n = 0$ or 400, where the bandwidth is half as large. In order to compare model and

observation, attention needs to be focused on the lowest frequencies because the MODE thermocline observations contain considerable energy there, in contrast to the abyssal data [an observation not fully available when the model and model data analyses were formulated (Owens, 1975)]. The model series mean ($n = 0$), containing the energy between 0 and $1/400$ cpd, vanishes. Removal of a linear trend reduces the energy in band $n(n \geq 1)$ by $6\Delta/\pi^2 n^2$, where Δ is a measure of the energy in the trend. For example, the lowest time-dependent estimate ($n = 1$) for the model spectra, containing the energy in the period range 133 to 400 days, is reduced in energy from the original series by approximately 60% of the energy associated with the trend. The trend was removed in order to filter out the dissipative (or spindown) time scale, but the spindown spectrum is not known with precision, so that some indeterminate amount of eddy energy was also removed. The spectral data used by Owens (1975) and Owens and Bretherton (1978) have been summed for present purposes over frequency bands that match to some extent the resolution thought to be natural (see below) for the moored instrument data base in the MODE area. A detailed description of this model and of the previous intercomparison of model results with data from the MODE area may be found in Bretherton (1975), Bretherton and Karweit (1975), Owens (1975) and Owens and Bretherton (1978).

Time series of sufficient length and quality to permit the formation of spatially averaged spectra with demonstrable stability out to the energetic (in the thermocline) lowest frequencies are not available. There is a particular location in the MODE area from which the longest term and highest quality data have been obtained. The moored current meter data available at a site labeled MODE CENTER (near 28°N, 70°W) consist of gappy time series containing approximately 750 data-days each, at nominal depths of 500, 1500 and 4000 m [technical

reports by Chausse and Tarbell (1976) and Tarbell and Spencer (1978) contain a more detailed description of the data used in the following]. DFT estimates were computed by a method first suggested by Thompson (1971) except that here the most conservative choice of frequencies is made. The DFT values were then summed in frequency to obtain spectral estimates. Both model and data are split into only three or four frequency bands, primarily in analogy to the spectral plots in Schmitz (1978) but also to match the frequencies accessible with the model time series. These results are listed in Table 1 in the form of the kinetic energy contained in four frequency bands or period ranges, at two depths for both model and data. The comparison is made at two of the three depths possible because utilizing the third depth at which data are available (near 1500 m) does not add significant new information for present purposes. Two model levels [numbers 2 and 6, see Owens and Bretherton (1978, Table I, p. 2)] were selected as indicative of thermocline (near 500 m) and abyssal depths (near 4000 m).

In Table 1, and in the following, the symbol K_E will denote kinetic energy per unit mass, with the per unit mass hereafter understood. $K_E(T)$, $K_E(U)$, $K_E(V)$ will signify the total, zonal and meridional K_E respectively.

3. Discussion

The first essential point to be made is that the method of analysis used by Owens (1975) and Owens and Bretherton (1978) in the computation of model K_E spectra (and K_E summed over all frequencies) simply rules out the possibility of any comparison between model and observed vertical structure at the lowest range of periods now accessible (and, therefore, for K_E summed over all of the data available). The model time series were truncated at a length of 200 days, and then the mean and trend were removed, thereby precluding the presence of

time-dependent kinetic energy in model spectra from periods >400 days (this energy is contained in the model mean, which was removed from the model time series). But in the data for MODE CENTER, the $K_E(T)$ associated with periods >400 days [and ≤ 1500 days for the particular data sets used (Band 1, Table 1)] is $20.4 \text{ cm}^2 \text{ s}^{-2}$ in the thermocline, and $0.3 \text{ cm}^2 \text{ s}^{-2}$ at abyssal depths. This contribution, although not statistically significant, is temporally stable, highly baroclinic and $\sim 40\%$ of the observed $K_E(T)$ at thermocline depths. That is, the model time series with mean removed do not contain any energy in a range of periods that is observed to be quite energetic in the thermocline only (dramatically baroclinic in K_E). Model and data should be compared over a common frequency range, and in particular the model kinetic energies should not be compared with the kinetic energy associated with the total data set available. The model results may or may not (see below) be more barotropic for $K_E(T)$ estimates over a common frequency range. The claim that the model successfully predicts time scales in the MODE area (Owens and Bretherton, 1978) should also be qualified: removal of the mean from the model data series implies an indeterminate conclusion for the lowest frequencies and therefore for a crucial contribution to spectral shape.

For the range of periods [taken to be 45–135 days (Band 3, Table 1)] that one might expect the model to be able to simulate the best, the predicted $K_E(T)$ in the thermocline ($9.3 \text{ cm}^2 \text{ s}^{-2}$) is very close to that observed ($10.4 \text{ cm}^2 \text{ s}^{-2}$). This constitutes percentage agreement to within 11%, probably within the stability of the data-based estimate. At abyssal depths the percentage agreement is not as good (within 26%) but still favorable. Considering the fact that the model scale factor for K_E is somewhat arbitrarily dependent on the selection of the particular time interval (which determines the total

TABLE 1. Spectral decompositions for data and model at selected depths.

Period range (days)	K_E ($\text{cm}^2 \text{ s}^{-2}$)	Mode center (500 m)	Mode center (4000 m)	Model (Layer 2)	Model (Layer 6)
Band 1 (400 $\rightarrow$ 1500)	T	20.4	0.3	—	—
	U	18.7	0.2	—	—
	V	1.7	0.1	—	—
Band 2 (133 $\rightarrow$ 400)	T	13.5	0.9	9.8	1.7
	U	8.5	0.7	5.2	0.8
	V	5.0	0.2	4.6	0.9
Band 3 (44 $\rightarrow$ 133)	T	10.4	5.7	9.3	4.2
	U	4.7	2.2	5.3	2.0
	V	5.7	3.5	4.0	2.2
Band 4 (5 $\rightarrow$ 44)	T	7.2	1.9	1.7	1.5
	U	2.4	0.9	1.0	0.8
	V	4.8	1.0	0.7	0.7

energy) over which one chooses to analyze the spindown model series, attention should probably be focused on the relative vertical distribution of K_E (i.e., the ratio of thermocline to abyssal K_E). For Band 3, these ratios for $K_E(T)$ are 2.2 and 1.8 for model and data, respectively, a comparatively good agreement in our opinion. The influence of removing the mean from the model series on the Band 1 comparison was very straightforward, but the influence of the removal of the trend on the other bands is much less so. For Band 3, the trend energy removed was $\sim 50\%$ of that retained. If one-half of this were eddy energy as opposed to energy related to spindown, then the associated uncertainty would be rather close to the percentage agreement between model and data. However, since the trend removal always acts at both depths to decrease K_E , the ratio of thermocline to abyssal energies should be relatively less affected. This ratio, as noted above, is probably the key inter-comparison parameter.

There is a small negative result for the Band 3 comparison that is of potential significance (Band 3 averages are over 11 natural bandwidths). A discrepancy exists between model and data in the zonal/meridional partitioning of K_E (Table 1). In the thermocline, the data for Band 3 are slightly more meridional (in sharp contrast to the case for Bands 1 and 2) and the model results slightly more zonal. That is, the often quoted (e.g., Rhines, 1977, p. 303) predominance of zonal over meridional K_E in the MODE area thermocline is a dominant feature of the data associated with the longest time scales (Band 1, and to some extent Band 2), but not Band 3 (or the temporal mesoscale as defined here). At abyssal depths the data are more meridional, while model results are in approximate equipartition.

Although the detrending technique used on the model series degrades the comparison percentage-wise the most ($\sim 60\%$) for the frequencies in Band 2 (the energy removed with the trend is approximately twice that retained, but again the ratio of energy lost to spindown to that due to eddies is unknown), it is nevertheless not ridiculous (Table 1). The model $K_E(T)$ for the thermocline is down from the data by $\sim 27\%$ and model $K_E(T)$ at abyssal depths is high by a factor of about 2. The vertical structure ratios are thus 5.8 and 15 for model and data, respectively, for Band 2. The data are more relatively zonal in Band 2 than the model results.

Even though the model estimates were calculated using 1024 model time series (with mean and linear trend removed), the number of truly independent estimates might be as large as 10, and the data-based Band 2 estimates are over only four natural bandwidths, so that the comparison is marginally significant for this reason as well as for reasons associated with trend removal. This observation suggests that Bands 2 and 3 might be combined

before comparison. Doing so, the vertical $K_E(T)$ ratios are 3.2 and 3.6 for model and data, respectively, again, good agreement. Model amplitudes for K_E are low for both depth ranges, by only 10–20% however. The zonal/meridional K_E partitioning is also favorable.

In Band 4 [periods 5–45 days nominal (Table 1)], the abyssal $K_E(T)$ agreement is reasonably good, but the model thermocline $K_E(T)$ is less than that observed by a factor of about 4 (the model is then much more barotropic than the data in this frequency band). Averages for the data-based Band 4 estimates are over 136 natural bandwidths so this divergence between model and data is statistically significant. In Band 4, the energy removed with the model trend is approximately equal to that retained. The disagreement between model and data in this range of periods could be rationalized if the relevant energy were associated with horizontal scales near the model grid spacing, and therefore not resolved [Owens and Bretherton (1978) noted that the model appeared to underpredict the energy at small horizontal scales]. Another possible reason for the disagreement in the 5–45 day period range could be associated with the lack of atmospheric forcing in the model.

4. Summary and conclusions

A numerical model designed to reproduce the statistical properties of the mesoscale eddy field in the MODE area has been compared with long-term moored instrument data. The amplitudes and ratio of eddy kinetic energy between thermocline and abyssal depths predicted by the model are frequency dependent: (i) in good agreement with the data in the period range of 50 to either 150 or to 400 days; (ii) in poor agreement with the data for the range of periods 5–50 days; (iii) in indeterminate agreement with the data at periods longer than 150 or 400 days because energy in this frequency range was pre-filtered from the model data series. Amplitudes for (i) above agree to within 10–30%, but this is not a crucial point because there is a somewhat arbitrary scale factor specified for the model. Of more significance, the ratios of thermocline to abyssal K_E in frequency band (i) agree to within 20%, and these ratios were least affected by removal of a trend from the model data series. Previous conclusions (Owens and Bretherton, 1978) that the model is too barotropic in K_E should be qualified by stipulating frequency range. It should not be claimed that the model successfully predicts time scales (Owens and Bretherton, 1978) for the same reason that led to conclusion (iii): the way that model results were filtered make this a moot point at present.

New model runs in regions like MODE should attempt to cope with the longer time scales [in

principle, new model runs should attempt to address time scales greater than mesoscale (periods longer than ~150 days) in any oceanic region because these scales are energetic and interesting everywhere that data are available, becoming dominant in the thermocline as one progresses into the ocean interior (Schmitz, 1978)]. Definitive comparisons for these longer time scales will also require a much more extensive and higher quality data base than presently available. One of us (WBO) is more strongly negative than the claim of indeterminate agreement between model and data at the longest time scales noted above, because 1) the original time series might not have contained the energetic, longer, baroclinic scales even before the mean and trend were removed (these raw model data are not now easily available) and 2) the very nature of the numerical model under discussion may preclude a comparison at the longest time scales because nonlocal influences (from outside the modeled region) could be important. In addition, the dissipative mechanism used might not be realistic, especially for the longer time scales. In any event, it may not be feasible to devise techniques for the model data analysis that separate with sufficient accuracy the energy associated with spin-down from longer period eddy energy.

We wish to emphasize that there is a sharp distinction in our minds between the three types of results of this investigation. First, we conclude that the intercomparison that we have made between the model results available and the MODE CENTER data is frequency dependent, with the outcome stated in some detail above. Second, since the intercomparison is frequency dependent, previous investigations that did not compare model and data over common frequencies should be qualified, and as an inference, future intercomparisons of other models with data should be cognizant of frequency domain. These two sets of results, our main points, are relatively firm and straightforward. There is a third type of result, related to the appropriateness or likely relevance of the entire class of models being considered to regions (like MODE) with energetic longer (than mesoscale) time scales. The essential point is that this question is at present indeterminate relative to our intercomparison, or with respect to any other with which we are familiar. As a minimum, improvement on an indeterminate result will require the ability to distinguish between the scales associated with spin-down and the longer eddy time scales. We do have the prejudice that the question of relevance for temporal scales longer than mesoscale will more likely be answered with a different class of models.

Finally, we wish to point out some of the ways that our methodology influences or restricts the results noted above. First, the removal of the trend from the model data series, designed to remove the

energy associated with the spindown of the model, also removes some unknown fraction of the eddy energy. To reiterate, we can calculate with good precision the amount of energy removed with the trend but do not know how much of that (thought to be small, however) is eddy energy. Nevertheless, the key intercomparison parameter, the ratio of thermocline to abyssal energy, is affected only to the extent that the energy percentage removed was depth dependent. Second, the model spectra are areal averages whereas the data used were from a single location (MODE CENTER). Spatial inhomogeneity is a prominent observed feature of the western North Atlantic (Schmitz, 1978). Kinetic energies from a site even as close as 100 km to the east of MODE CENTER are more baroclinic, but the difference may not be statistically significant. In any event, we don't have the data needed to compute spatially averaged spectra that would be significantly more appropriate than the site spectra used, and even given the areal data, we doubt that the outcome of our investigation would be qualitatively (or possibly even semi-quantitatively) different.

Acknowledgments. This investigation was supported by the Office of Naval Research under Contract N00014-76-C-0197, NR 083-400 and by the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE75-03962. This is Contribution No. 4341 from the Woods Hole Oceanographic Institution and No. 120 from the open ocean dynamics experiments, MODE and POLYMODE.

REFERENCES

- Bretherton, F. P., 1975: Recent developments in dynamical oceanography. *Quart. J. Roy. Meteor. Soc.*, **101**, 705-721.
- , and M. Karweit, 1975: Mid-ocean mesoscale modelling. *Numerical Models of Ocean Circulation*, Ocean Affairs Board, Nat. Res. Coun., Washington, DC, 237-249.
- Chausse, D., and S. Tarbell, 1976: A compilation of moored current data and associated oceanographic observations, Vol. 12 [1973 Mid-Ocean Dynamical Experiment (MODE)]. Tech. Rep. WHOI-76-101, Woods Hole Oceanographic Institution.
- Owens, W. B., 1975: A numerical study of mid-ocean mesoscale eddies. Ph.D. thesis, The Johns Hopkins University, and NCAR Cooperative thesis No. 36, 105 pp.
- , and F. P. Bretherton, 1978: A numerical study of mid-ocean eddies. *Deep-Sea Res.*, **25**, 1-14.
- Rhines, P. B., 1977: The dynamics of unsteady currents. *The Sea*, Vol. 6, E. D. Goldberg, Ed., Wiley, 189-318.
- Schmitz, W. J., Jr., 1978: Observations of the vertical structure of low frequency fluctuations in the western North Atlantic. *J. Mar. Res.*, **36**, 295-310.
- Tarbell, S., and A. Spencer, 1978: A compilation of moored current data and associated observations, Vol. 16 [MODE-Site, 1971-75]. Tech. Rep. WHOI-78-5, Woods Hole Oceanographic Institution.
- Thompson, R. O. R. Y., 1971: Spectral estimation from irregularly spaced data. *IEEE Trans., Geosci. Electron.*, **GE-9**, 107-110.

MANDATORY DISTRIBUTION LIST

FOR UNCLASSIFIED TECHNICAL REPORTS, REPRINTS, AND FINAL REPORTS
PUBLISHED BY OCEANOGRAPHIC CONTRACTORS
OF THE OCEAN SCIENCE AND TECHNOLOGY DIVISION
OF THE OFFICE OF NAVAL RESEARCH

(REVISED NOVEMBER 1978)

1	Deputy Under Secretary of Defense (Research and Advanced Technology) Military Assistant for Environmental Science Room 3D129 Washington, D.C. 20301	12	Defense Documentation Center Cameron Station Alexandria, VA 22314 ATTN: DCA
	Office of Naval Research 800 North Quincy Street Arlington, VA 22217		Commander Naval Oceanographic Office NSTL Station Bay St. Louis, MS 39522
3	ATTN: Code 483	1	ATTN: Code 8100
1	ATTN: Code 460	1	ATTN: Code 6000
2	ATTN: 102B	1	ATTN: Code 3300
1	CDR J. C. Harlett, (USN) ONR Representative Woods Hole Oceanographic Inst. Woods Hole, MA 02543	1	NODC/NOAA Code D781 Wisconsin Avenue, N.W. Washington, D.C. 20235
	Commanding Officer Naval Research Laboratory Washington, D.C. 20375		
6	ATTN: Library, Code 2627		

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER WHOI-80- 56	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES FOR MODE EDDIES		5. TYPE OF REPORT & PERIOD COVERED Technical
7. AUTHOR(s) William J. Schmitz, Jr. and W. Brechner Owens		6. PERFORMING ORG. REPORT NUMBER WHOI CON. NO. 4341
9. PERFORMING ORGANIZATION NAME AND ADDRESS Woods Hole Oceanographic Institution Woods Hole, MA 02543		8. CONTRACT OR GRANT NUMBER(s) N00014-76-C-0197; OCE 75-03962
11. CONTROLLING OFFICE NAME AND ADDRESS NORDA National Space Technology Laboratory Bay St. Louis, MS 39529		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE November 1980
		13. NUMBER OF PAGES
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Reprinted from "Journal of Physical Oceanography, Vol. 9, No. 6, November 1979, pp. 1294-1297".		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) 1. Vertical structure 2. Model-data intercomparison 3. MODE eddy kinetic energy		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the comparison is		

indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.

<p>Woods Hole Oceanographic Institution WHOI-80-56</p> <p>OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES FOR MODE EDDIES by William J. Schmitz, Jr. and M. Brechner Owens. Reprinted from "Journal of Physical Oceanography", Vol. 9, No. 6, November 1979, pp. 1294-1297". Nov. 1980. Prepared for the Office of Naval Research under Contract N00014-76-C-0197; NR 083-400 and for the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE 75-03962.</p> <p>It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the agreement is indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.</p>	<p>1. Vertical structure</p> <p>2. Model-data inter-comparison</p> <p>3. MODE eddy kinetic energy</p> <p>I. Schmitz, William J., Jr.</p> <p>II. Owens, M. Brechner</p> <p>III. N00014-76-C-0197; NR 083-400</p> <p>IV. OCE 75-03962</p> <p>This card is UNCLASSIFIED</p>	<p>Woods Hole Oceanographic Institution WHOI-80-56</p> <p>OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES FOR MODE EDDIES by William J. Schmitz, Jr. and M. Brechner Owens. Reprinted from "Journal of Physical Oceanography", Vol. 9, No. 6, November 1979, pp. 1294-1297". Nov. 1980. Prepared for the Office of Naval Research under Contract N00014-76-C-0197; NR 083-400 and for the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE 75-03962.</p> <p>It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the agreement is indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.</p>	<p>1. Vertical structure</p> <p>2. Model-data inter-comparison</p> <p>3. MODE eddy kinetic energy</p> <p>I. Schmitz, William J., Jr.</p> <p>II. Owens, M. Brechner</p> <p>III. N00014-76-C-0197; NR 083-400</p> <p>IV. OCE 75-03962</p> <p>This card is UNCLASSIFIED</p>	<p>Woods Hole Oceanographic Institution WHOI-80-56</p> <p>OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES FOR MODE EDDIES by William J. Schmitz, Jr. and M. Brechner Owens. Reprinted from "Journal of Physical Oceanography", Vol. 9, No. 6, November 1979, pp. 1294-1297". Nov. 1980. Prepared for the Office of Naval Research under Contract N00014-76-C-0197; NR 083-400 and for the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE 75-03962.</p> <p>It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the agreement is indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.</p>	<p>1. Vertical structure</p> <p>2. Model-data inter-comparison</p> <p>3. MODE eddy kinetic energy</p> <p>I. Schmitz, William J., Jr.</p> <p>II. Owens, M. Brechner</p> <p>III. N00014-76-C-0197; NR 083-400</p> <p>IV. OCE 75-03962</p> <p>This card is UNCLASSIFIED</p>	<p>Woods Hole Oceanographic Institution WHOI-80-56</p> <p>OBSERVED AND NUMERICALLY SIMULATED KINETIC ENERGIES FOR MODE EDDIES by William J. Schmitz, Jr. and M. Brechner Owens. Reprinted from "Journal of Physical Oceanography", Vol. 9, No. 6, November 1979, pp. 1294-1297". Nov. 1980. Prepared for the Office of Naval Research under Contract N00014-76-C-0197; NR 083-400 and for the International Decade of Ocean Exploration Office of the National Science Foundation under Grant OCE 75-03962.</p> <p>It is demonstrated that the outcome of an intercomparison between data and the vertical distribution of eddy kinetic energy predicted by a previously developed numerical model of the MODE area is frequency dependent. In the range of periods from 50 to 150 or even to 400 days (one definition of the temporal mesoscale, the scale that the model was designed to simulate), the comparison is quite good. For periods in the range of 5 to 50 days, the agreement is poor. For periods longer than 400 days, the agreement is indeterminate. Earlier conclusions concerning the relation of model results to the MODE data should be qualified by stipulating frequency range, and future intercomparisons for any model in all regions should be conscious of the desirability of doing so across common frequencies.</p>	<p>1. Vertical structure</p> <p>2. Model-data inter-comparison</p> <p>3. MODE eddy kinetic energy</p> <p>I. Schmitz, William J., Jr.</p> <p>II. Owens, M. Brechner</p> <p>III. N00014-76-C-0197; NR 083-400</p> <p>IV. OCE 75-03962</p> <p>This card is UNCLASSIFIED</p>
--	--	--	--	--	--	--	--