

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40

**217,000 year-old DNA sequences of green sulfur bacteria in Mediterranean sapropels and their implications for the reconstruction of the paleoenvironment**

Marco J.L. Coolen<sup>1</sup> and Jörg Overmann<sup>2\*</sup>

<sup>1</sup> Woods Hole Oceanographic Institution, Department of Marine Chemistry and Geochemistry, Woods Hole, MA 02543, USA

<sup>2</sup> Section Microbiology, Ludwig-Maximilians-Universität München, Maria-Ward-Str. 1a, D-80638 München, Germany

*Running title:* Fossil DNA sequences of green sulfur bacteria

*Keywords:* ancient DNA, oceanic anoxic events, *Chlorobiaceae*, fossil DNA, green sulfur bacteria, sapropels

\*For correspondence. Email [j.overmann@lrz.uni-muenchen.de](mailto:j.overmann@lrz.uni-muenchen.de); Tel. +49-(0)89-2180-6123; Fax +49-(0)89-2180-6125

## 1 SUMMARY

2 Deep-sea sediments of the eastern Mediterranean harbor a series of dark, organic carbon-rich  
3 layers, so-called sapropels. Within these layers, the carotenoid isorenieratene was detected. Since  
4 it is specific for the obligately anaerobic phototrophic green sulfur bacteria, the presence of  
5 isorenieratene may suggest that extended water column anoxia occurred in the ancient  
6 Mediterranean Sea during periods of sapropel formation. Only three carotenoids (isorenieratene,  
7  $\beta$ -isorenieratene and chlorobactene) are typical for green sulfur bacteria and thus do not permit  
8 to differentiate between the ~80 known phylotypes. In order to reconstruct the paleoecological  
9 conditions in more detail, we searched for fossil 16S rRNA gene sequences of green sulfur  
10 bacteria employing ancient DNA methodology. 540 bp-long fossil sequences could indeed be  
11 amplified from up to 217,000-year-old sapropels. In addition, such sequences were also  
12 recovered from carbon-lean intermediate sediment layers deposited during times of an entirely  
13 oxic water column. Unexpectedly, however, all the recovered 16S rRNA gene sequences  
14 grouped with freshwater or brackish, rather than truly marine, types of green sulfur bacteria. It is  
15 therefore feasible that the molecular remains of green sulfur bacteria originated from populations  
16 which thrived in adjacent freshwater or estuarine coastal environments rather than from an  
17 indigenous pelagic population.

18

## 1 INTRODUCTION

2 All known green sulfur bacteria and about half of the species of purple sulfur bacteria are  
3 obligate anaerobic photolithoautotrophs which can only grow in the presence of light and  
4 reduced sulfur compounds as photosynthetic electron donors (Pfennig and Trüper, 1989;  
5 Overmann, 2001). Their specific physiological requirements render these bacteria suitable  
6 indicator organisms for past environmental conditions (Passier *et al.*, 1999; Menzel *et al.*, 2002).

7 Eastern Mediterranean sediments harbor more than 1 cm-thick, organic carbon-rich layers,  
8 so-called sapropels. Sapropels contain >2 % and up to 30% per weight of organic carbon and are  
9 embedded in hemipelagic carbonate oozes (< 0.5 wt% organic C) (Kidd *et al.*, 1978). The  
10 enhanced organic carbon preservation during sapropel formation has been explained by anoxia  
11 of deep Mediterranean bottom water (Oceanic Anoxic Events; Rossignol-Strick, 1985; Rohling  
12 and Hilgen, 1991; Passier *et al.*, 1999) or by increased marine primary production (Calvert,  
13 1983; Calvert *et al.*, 1992). 1.8 to 3.0 million-year-old Pliocene sapropels were shown to contain  
14 the carotenoid isorenieratene and its diagenetic derivatives (Passier *et al.*, 1999; Menzel *et al.*,  
15 2002). Since isorenieratene occurs almost exclusively in green sulfur bacteria, it was concluded  
16 that sulfidic bottom waters extended into the photic zone during sapropel formation and that the  
17 Mediterranean Sea ecosystem underwent repeated and major oxic-anoxic shifts during the past 3  
18 million years.

19 However, isorenieratene has also been detected in the actinobacteria *Streptomyces griseus*  
20 and *Brevibacterium linens* (Krubasik and Sandmann, 2000; Krügel *et al.*, 1999). On the other  
21 hand, certain strains of green sulfur bacteria do not contain any detectable amounts of those  
22 carotenoids which were originally thought to be typical for this group (Glaeser *et al.* 2002).  
23 Since green sulfur bacteria form a distinct and coherent phylogenetic lineage (Overmann and  
24 Tuschak, 1997; Imhoff, 2003), 16S rRNA gene sequences provide an alternative means to trace  
25 their occurrence and species composition in the environment (Overmann *et al.*, 1999). While  
26 green sulfur bacteria contain only three specific carotenoids (isorenieratene,  $\beta$ -isorenieratene and  
27 chlorobactene) (Overmann, 2001), 80 different 16S rRNA gene sequence types are recognized to

1 date (A. Manske and J. Overmann, submitted). The three carotenoids occur across all different  
2 subgroups of the green sulfur bacteria (Imhoff, 2003), whereas species isolated from the marine  
3 environment form a single well-separated sequence cluster (marine group 1) (Imhoff, 2003;  
4 Manske *et al.*, 2005). Therefore, analyses of their fossil 16S rRNA gene sequences in subsurface  
5 sediments would not only provide independent evidence for the occurrence of green sulfur  
6 bacteria but also permit a more detailed reconstruction of their paleoenvironment. To date,  
7 however, such a paleomicrobiological analysis has not been performed in the marine  
8 environment.

9 Fully hydrated DNA spontaneously decays over only hundreds of years, mainly through  
10 hydrolysis and oxidation (Shapiro, 1981; Lindahl, 1993; Hofreiter *et al.*, 2001). However, low  
11 temperatures, high ionic strength, anoxic conditions and protection from enzymatic degradation  
12 by adsorption extend the half-life of intact DNA by one or two orders of magnitude (Lindahl,  
13 1993; Poinar *et al.*, 1996; Willerslev *et al.*, 2004a). So far, intact DNA of anoxygenic  
14 phototrophic bacteria could be extracted from up to 9,100 year-old holocene lake sediments and  
15 was analyzed by PCR amplification and sequencing (Coolen and Overmann, 1998). Recently,  
16 amplification of fossil chloroplast DNA has been found to be reliable for samples which are  
17 hundreds of thousands of years old (Willerslev *et al.*, 2003). Under favorable conditions of  
18 preservation, genomic DNA of anoxygenic phototrophic bacteria thus may have persisted for  
19 similar time periods in Mediterranean sapropels.

20 In the present study, fossil DNA sequences and carotenoids of ancient green sulfur bacteria  
21 could be recovered from sapropels deposited between 8,000 and 217,000 years ago in the  
22 Eastern Mediterranean. Phylogenetic analyses were used to infer the most likely origin of the  
23 respective bacteria and provide a more detailed picture of the development of this marine  
24 ecosystem.

25

## 1 RESULTS

### 2 *Vertical structure and age of sediment layers*

3 A four meter-long sediment core obtained at a water depth of 2155 m from the Eastern  
4 Mediterranean southeast of Crete contained four different sapropels which were identified as  
5 sapropels S<sub>1</sub>, S<sub>6</sub>, S<sub>7</sub>, and S<sub>8</sub> based on geochemical evidence (Coolen *et al.*, 2002; Fig. 1). The  
6 ages of these sapropels are 8,000 (S<sub>1</sub>), 172,000 (S<sub>6</sub>), 195,000 (S<sub>7</sub>) and 217,000 (S<sub>8</sub>) years (Emeis  
7 *et al.*, 2000). The organic carbon content of the sapropels ranged between 2.3 and 8.5% (w/w),  
8 whereas intermediate layers contained only up to 0.16 % (Coolen *et al.*, 2002). The intermediate  
9 layer Z<sub>1</sub> is likely to represent a turbidite (K.-C. Emeis; pers. comm.).

10 For subsequent analyses, the top of the sediment (Z<sub>0</sub>, 4-6 cm below surface), each sapropel  
11 layer (S<sub>1</sub>, 13-17 cm; S<sub>6</sub>, 262-266 cm; S<sub>7</sub>, 314-318 cm; and S<sub>8</sub>, 364-368 cm), and three  
12 intermediate hemipelagic layers between the sapropels (denoted Z<sub>1</sub>, 62-66 cm; Z<sub>6</sub>, 285-289 cm;  
13 and Z<sub>7</sub>, 333-337 cm) were selected (Fig. 1).

### 14 *Green sulfur bacterial carotenoids in late Pleistocene and Holocene sapropels*

15 Isorenieratene and  $\beta$ -isorenieratene could be detected in all four sapropel layers where they  
16 reached concentrations between 56.4 and 1280 ng·(g dry weight sediment)<sup>-1</sup> (Fig. 1C). Initially,  
17 both carotenoids were identified by HPLC based on their characteristic retention time and  
18 absorption spectra (peaks at 454 and 482 nm and a shoulder at 428 nm). Analysis of these  
19 compounds by LC-MS in the APCI positive ion mode yielded major signals at m/z values of 529  
20 and 133. These masses match those of the protonated isorenieratene (i.e., [M+H]<sup>+</sup>) and of  
21 1,2,3,4-tetramethylbenzene, respectively, the latter representing a fragment typical of  
22 isorenieratene and  $\beta$ -isorenieratene. Our mass spectrometric analysis thus provides the first  
23 evidence for the presence of carotenoids of green sulfur bacteria in late Pleistocene and  
24 Holocene sapropels.

25 In contrast, only traces of carotenoid compounds with the retention times and absorption  
26 spectra of isorenieratene and  $\beta$ -isorenieratene were detected in intermediate layers.

1 Concentrations ranged from  $(1.4 \pm 0.1)$  to  $(5.9 \pm 0.3)$   $\text{ng}\cdot(\text{g dry weight sediment})^{-1}$  (Fig. 1C).  
2 Although above the detection limit, these trace amounts did not allow further analyses by LC-  
3 MS. Therefore, it cannot be completely ruled out that other carotenoids were present, though  
4 carotenoids with retention times and absorption spectra identical to isorenieratene/ $\beta$ -  
5 isorenieratene are currently not known.

#### 6 *Detection and quantification of DNA of green sulfur bacteria*

7 After optimizing our extraction protocol, genomic DNA could be isolated from all eight  
8 sediment layers (Fig. 1A). Between 0.7 and 5.0  $\mu\text{g}$  of DNA were extracted per gram dry weight  
9 of sediment from the four sapropels. These amounts significantly exceeded those obtained from  
10 the intermediate layers ( $0.01$  to  $0.12$   $\mu\text{g}\cdot(\text{g dry wt})^{-1}$ ) and were only matched by the value from  
11 the sediment surface ( $1.04$   $\mu\text{g}\cdot(\text{g dry wt})^{-1}$ ). Analysis by gel electrophoresis revealed that the  
12 DNA mostly consisted of up to 23 - 30 kb long fragments (data not shown). The shortest  
13 fragments detected measured 500 bp. This length distribution was even observed in extracts from  
14 the 217,000-year-old Mediterranean sapropel S<sub>8</sub>.

15 The specific primer combination and PCR conditions used in the present study permit the  
16 selective amplification of 16S rRNA gene fragments of green sulfur bacteria (Overmann *et al.*,  
17 1999). With the exception of the intermediate layer Z<sub>6</sub>, all samples yielded amplification  
18 products. The DNA extract of sapropel S<sub>7</sub> yielded only a faint amount of DNA of the correct  
19 molecular size, but also some unspecific amplification products. Of all sapropel layers  
20 investigated, S<sub>7</sub> contains the by far highest amount of kerogen (total organic carbon content of  
21 8.5% as compared to  $\leq 4.4\%$  in other sapropels; Coolen *et al.*, 2002). Similarly, Z<sub>6</sub> contained the  
22 highest amount of total organic carbon of all intermediate layers (0.16 versus  $\leq 0.01\%$ ). Most  
23 likely, the failure to obtain appropriate amplification products from these two layers resulted  
24 from the unfavourable ratio of fossil DNA to organic kerogen.

25 Each amplification reaction was rigorously checked for contamination with extraneous DNA  
26 by routinely including a set of procedural blanks and PCR controls (as detailed in the Materials

1 and Methods section). Yet, neither the controls for a contamination of PCR reagents, nor the  
2 controls for a contamination during DNA extraction ever yielded amplification products (Fig. 2).

3 A specific dot blot hybridization protocol was used to quantify the amount of green sulfur  
4 bacterial DNA in the different sediment layers after amplification (Fig. 3). Our original  
5 amplification method was found to be highly sensitive but not suitable for quantification because  
6 it did not yield a continuously increasing response with increasing concentrations of target DNA  
7 used in the PCR. Calibration was possible, however, when an altered amplification PCR protocol  
8 was employed (Fig. 3, standards). This technique permitted a detection of green sulfur bacteria  
9 DNA if the latter constituted as little as 0.0065% of the total community DNA. Green sulfur  
10 bacterial 16S rRNA gene sequences could be quantified in six of the sediment layers (Fig. 1C,  
11 Fig.3). DNA extracted from the intermediate layers  $Z_6$  and  $Z_7$  did not yield detectable signals,  
12 although the latter yielded a product with the original, more sensitive, PCR method.

13 Although most sapropel layers contained the highest absolute amounts of green sulfur  
14 bacterial DNA (Fig. 1C), maxima of the relative amounts did not always coincide with the  
15 presence or absence of sapropels (Fig. 1B). The highest percentage of green sulfur bacterial  
16 DNA ( $0.55 \pm 0.02\%$  of total community DNA) was determined for sapropel  $S_8$ , whereas the  
17 second largest fraction of 16S rRNA genes of green sulfur bacteria ( $0.088\%$  of total community  
18 DNA) was determined for intermediate layer  $Z_1$  (Fig. 1B).

### 19 *Phylogenetic identification of the fossil green sulfur bacteria*

20 In order to identify individual sequences, the amplified 16S rRNA gene fragments were analyzed  
21 further by denaturing gradient gel electrophoresis and sequencing. In order to be able to analyze  
22 sequences from intermediate layer  $Z_7$ , PCR products obtained with the more sensitive  
23 amplification method were used. However, only PCR reactions which yielded exclusively  
24 specific products were analyzed, thereby excluding the amplification products from  $S_7$ .

25 A total of six different melting types could be recognized by DGGE fingerprinting (Fig. 4).  
26 Nine of the DNA bands were excised and sequenced. The results confirm that sequences of

1 green sulfur bacteria had been selectively amplified from the extracted DNA. Seven of the  
2 bands (A through G in Fig. 4) yielded unambiguous sequences. Of the latter, bands A, B, E and F  
3 not only exhibited the same melting behavior during DGGE, but also contained the same  
4 sequence. This phylotype was found to be identical to those of 11 other strains or environmental  
5 sequences (Fig. 5) and fell into Group 3 among the green sulfur bacteria. Sequence type C was a  
6 member of Group 4a and only present in the intermediate layer  $Z_1$ , yet could not be detected in  
7 the adjacent sapropel  $S_1$  or the surface layer  $Z_0$ . Two other sequences (D and G) fell into Group 3  
8 (Fig. 5). Each represented a distinct, so far unknown sequence type (Fig. 5) and was found in  
9 only a single sapropel layer but was missing in all other samples (Fig. 4).

#### 10 *Adsorptive binding of DNA to the sediments*

11 Sediment material of the sapropel  $S_6$  and the intermediate layer  $Z_6$  was chosen for adsorption  
12 assays since they contained comparatively low amounts of indigenous DNA compared to the  
13 other layers (Fig. 1A). Since the concentration of indigenous DNA determined ( $1.44 \mu\text{g}\cdot(\text{g dry}$   
14  $\text{weight sediment})^{-1}$  in  $S_6$ ) amounted to only a very minor fraction (0.003%) of the total  
15 adsorption capacity, the DNA already present in the samples did not interfere with these  
16 adsorption assays.

17 The sapropel exhibited an extraordinarily high maximum adsorption capacity  $S_{max}$  for DNA  
18 ( $(79.3 \pm 2.8) \mu\text{mol double-stranded DNA (g dry wt)}^{-1}$ ; corresponding to  $52.8 \text{ mg (g dry wt)}^{-1}$ )  
19 (Fig. 6, Table 1). This value is comparable to that of pure montmorillonite and is only surpassed  
20 by purified humic acids (Table 1). Albeit lower, the adsorption capacity of  $Z_6$  for double  
21 stranded DNA was still significantly higher than that of soil (Table 1).

22

23

## 1 DISCUSSION

### 2 *Authenticity of the green sulfur bacterial DNA*

3 Analyses of ancient DNA are usually based on minute amounts of highly degraded template  
4 molecules and thus are subject to a considerable risk of contamination (Cooper and Poinar,  
5 2000). In order to avoid sources of extraneous DNA in the present study, subsampling of the  
6 sediment cores was performed in a separate, PCR-product-free laboratory in which green sulfur  
7 bacteria had never been worked with before, and a security level 2 laminar flow chamber  
8 dedicated to work with low DNA template number samples was used for subsequent DNA  
9 extraction and to set up amplification reactions. Each set of DNA extractions included controls  
10 for contamination of the chemicals and vessels employed. None of these extraction controls  
11 yielded an amplification product, indicating that no extraneous DNA from green sulfur bacteria  
12 had been introduced during DNA isolation. Furthermore, each amplification run comprised two  
13 reactions devoid of DNA template in order to control separately for a potential contamination of  
14 PCR reagents and tubes. These negative controls also did not yield any PCR product. Finally,  
15 quantification of fossil green sulfur bacterial DNA was performed in three independent parallels  
16 to test reproducibility of the results.

17 Whereas parallel extraction trials for the identical sediment layer were not feasible due to the  
18 restricted availability of deep sea sediment samples, a comparison of the results from the  
19 consecutive sediment horizons further confirms that the DNA genuinely originated from these  
20 samples. Sequences of green sulfur bacteria were not only detected on a single occasion, but  
21 occurred in seven out of eight different sediment layers. Furthermore, amplification of green  
22 sulfur bacterial sequences was reproducible for the same extract (compare standard deviations  
23 for quantification of green sulfur bacterial amplification products in Fig. 1C).

24 Three of the four sequence types obtained (Z1-C, S6-D, S8-G) were detected in only a single  
25 sample. It appears unlikely that each of the three sediment samples Z1, S6 and S8 was  
26 contaminated with yet another, different type of green sulfur bacterium. Two of these sequence  
27 types (S6-D, S8-G) have neither been isolated as a clone nor from a culture before. It is also

1 extremely unlikely that two samples were contaminated individually, yet by two different  
2 sequences which are not available in any laboratory world wide. Our phylogenetic analysis of the  
3 recovered sequences thus provides independent evidence for the conclusion that the green sulfur  
4 bacterial DNA is indigenous to the Mediterranean sediments.

#### 5 *Fossil origin and mechanisms of persistence of green sulfur bacterial DNA*

6 Compared to the multitude of studies targeting higher organisms, studies of ancient bacteria have  
7 been limited to a few pathogenic species (*Mycobacterium tuberculosis*, *M. leprae*, *Yersinia pestis*  
8 and *Treponema pallidum*) or to intestinal bacteria, and cover a time period of only the last 5,400  
9 years (Rollo, 1998). Information on past bacterial communities in the environment is very sparse  
10 (Coolen and Overmann, 1998; Coolen et al., 2006; Willerslev *et al.*, 2004b) which can be  
11 attributed to the difficulty to distinguish ancient from modern bacterial DNA.

12 All cultured representatives of the green sulfur bacteria are obligately anaerobic  
13 photolithoautotrophs and, accordingly, require the simultaneous presence of light and sulfide for  
14 growth. The family *Chlorobiaceae* comprises all known green sulfur bacteria plus numerous  
15 environmental sequences (Overmann and Tuschak, 1997; Overmann *et al.*, 1999; A. Manske and  
16 J. Overmann, submitted). Similar to their cultured relatives, all so-far-uncultured members of  
17 this group have been exclusively detected in illuminated sulfidic environments like the  
18 chemocline of lakes, lagoons or benthic microbial mats. As the sequences recovered in the  
19 present study clearly fall within this group, the available evidence indicates that they originated  
20 from obligately anaerobic photolithoautotrophs. All our attempts to enrich green sulfur bacteria  
21 by different cultivation methods (Overmann, 2001; Manske *et al.*, 2005) failed, supporting the  
22 view that viable green sulfur bacteria do not exist in these sediments. In conclusion, the green  
23 sulfur bacterial sequences obtained in the present study are highly unlikely to originate from  
24 bacteria growing within the sediments. Rather, the cells grew outside the sediments and the  
25 aphotic portion of the water column and were subsequently deposited in Mediterranean deep sea  
26 sediments.

1 DNA spontaneously decays into 100 - 600 bp short fragments after its deposition in  
2 lacustrine sediments (Coolen and Overmann, 1998). Based on our data, green sulfur bacterial  
3 genome fragments at least 540-bp in size persisted in Mediterranean sapropels over 217,000  
4 years. Survival of ancient DNA in a lake sediment could be demonstrated for a time period of  
5 9,100 years (Coolen and Overmann, 1998) due to the short lifespan of this ecosystem. The  
6 partial 16S rRNA gene fragments obtained in this study are among the oldest authenticated  
7 ancient bacterial sequences available to date. For animals and plants, the oldest authenticated  
8 records come from 60,000-year-old remains of brown bears (Barnes *et al.*, 2002), and from  
9 ~400,000-year-old chloroplast DNA (Willerslev *et al.*, 2003), respectively, both recovered from  
10 Alaskan permafrost. The age of the fossil 16S rRNA gene sequences of green sulfur bacteria  
11 detected in the present study falls well within this time frame. Based on the relatively high *in situ*  
12 temperature of 14°C, however, the efficient preservation of fossil DNA in eastern Mediterranean  
13 sediments must be attributed to factors other than low temperature.

14 Besides low temperatures, the persistence of fossil DNA is significantly extended by high  
15 ionic strength, anoxic conditions, and rapid dehydration and adsorption of DNA (Lindahl, 1993;  
16 Poinar *et al.*, 1996; 2003; Willerslev *et al.*, 2004a). The ionic strength and anoxic conditions in  
17 eastern Mediterranean sediments are comparable to other sediment environments. Adsorption to  
18 clay and other mineral surfaces significantly decreases the degradation rates for organic  
19 compounds in soils (Jones and Edwards, 1998), and marine sediments (Keil *et al.*, 1994) by up to  
20 five orders of magnitude. Adsorptive binding to hydroxyapatite retards the spontaneous decay of  
21 DNA (Lindahl, 1993). In addition to spontaneous hydrolysis, microbial DNase activity leads to  
22 rapid degradation of free DNA in water and sediments (Lorenz and Wackernagel, 1994), but is  
23 also effectively prevented by adsorption even in sandy sediments which bind DNA much less  
24 efficiently than the sapropels (Romanowski *et al.*, 1991; Crecchio and Stotzky, 1998). Based on  
25 our results, the outstanding adsorption capacity of eastern Mediterranean sediments, in particular  
26 of the sapropel matrix, represents the major reason for the efficient preservation of fossil DNA of

1 green sulfur bacteria. Potentially, sapropels may also represent important archives of fossil DNA  
2 from other microorganisms.

### 3 *Implications for the reconstruction of past ecosystems*

4 To date, isorenieratene and its degradation products have been used as an indicator of past water  
5 column anoxia (Passier *et al.*, 1999; Menzel *et al.*, 2002). Because of their very limited diversity,  
6 carotenoids do not permit a differentiation between species with contrasting physiology and  
7 ecology. In the eastern Mediterranean, isorenieratene has previously been shown to occur in  $\geq 1.8$   
8 million-year-old Pliocene sapropels (Passier *et al.*, 1999; Menzel *et al.*, 2002). However, only  
9 four of the 83 currently recognized sapropels (Emeis *et al.*, 2000) have been investigated so far.  
10 It was therefore unknown whether isorenieratene represents a typical constituent of the  
11 sapropels. The present study provides the first instance of occurrence of isorenieratene in marine  
12 deposits which are sufficiently young to also harbor 16S rRNA gene sequences of green sulfur  
13 bacteria. The latter are high resolution fossil biomarkers which permit an improved assessment  
14 of their origin and hence a more detailed reconstruction of the paleoenvironment.

15 All but one (Z1C) of the 16S rRNA gene sequences recovered were phylogenetically  
16 affiliated with typical freshwater or brackish water species, whereas no single representative of  
17 the marine group 1 could be detected. Secondly, the presence of green sulfur bacterial 16S rRNA  
18 genes in intermediate layers was unexpected since the latter were deposited under a fully  
19 oxygenated water column (Schmiedl *et al.*, 1998). Theoretically, DNA could have reached  
20 intermediate layers through vertical percolation after deposition in the sapropel layers. However,  
21 vertical migration of fossil DNA fragments should be independent of their actual base sequence  
22 and, as a result, sequence types present in sapropels should also be detectable in the adjacent  
23 intermediate layers. Yet, phylotype C was detected in intermediate layer Z<sub>1</sub>, but not in the  
24 adjacent S<sub>1</sub> (compare Fig. 4). This indicates that the strong adsorption to the sediment particles  
25 has effectively immobilized the fossil DNA of green sulfur bacteria. Thirdly, a long-distance  
26 transport of green sulfur bacterial biomarkers has been shown for North Atlantic deep-sea

1 sediments (Rosell-Melé *et al.*, 1997). Indeed, a turbidite origin was proposed for the intermediate  
2 layer Z1 (K.-C. Emeis, pers. comm.) which contained the by far highest fraction of green sulfur  
3 bacterial DNA among all organic carbon lean intermediate layers. It thus appears feasible that at  
4 least the DNA of green sulfur bacterial phylotypes affiliated with typical freshwater species may  
5 have originated from environments like coastal lagoons where these bacteria frequently form  
6 dense blooms.

7 In conclusion, our results suggest that not all green sulfur bacteria deposited in the ancient  
8 Mediterranean Sea during the last 217,000 years were autochthonous but instead are more likely  
9 to have originated from Mediterranean coastal environments. Our cumulative evidence casts  
10 some doubts on the hypothesis of the presence of green sulfur bacteria in the open waters of the  
11 eastern Mediterranean and, by inference, questions the alleged past photic zone anoxia. A future  
12 isolation and physiological characterization of the two previously unknown phylotypes detected  
13 in the present study will help to more precisely assign a certain habitat, hence origin, to the fossil  
14 remains of green sulfur bacteria present in the sapropel layers.

15

## 1 **EXPERIMENTAL PROCEDURES**

### 2 *Sampling and sample preparation*

3 A gravity core (#69-2SL) was obtained during cruise 40 leg 4 of the R/V Meteor on January 30,  
4 1998, at position 33°51,53'N and 24°54,46'E southeast of Crete. The core was collected at a  
5 depth of 2155 m and was subsampled employing the aseptic techniques developed for the  
6 recovery of fossil DNA from sediment samples (Coolen and Overmann, 1998; 2000). It was cut  
7 longitudinally which left behind a potentially contaminated surface. This surface was rapidly  
8 frozen with powdered dry ice and subsequently lifted off. Through the freshly exposed surface, 5  
9 cm<sup>3</sup> subsamples were retrieved aseptically using sterile plastic syringes which had their ends cut  
10 off and were immediately stored in sterile vials at -80<sup>0</sup>C until extraction.

### 11 *Precautions and controls to prevent contamination*

12 Preparation of sediment samples was performed directly after retrieval of sediment cores in the  
13 laboratory on board of the R/V Meteor, employing the aseptic DNA techniques established  
14 previously (Coolen and Overmann, 1998; Coolen *et al.*, 2006). Green sulfur bacteria had never  
15 been introduced or worked with in these premises. Further precautions against contamination of  
16 the samples with foreign DNA included the use of a laminar flow hood dedicated to low  
17 template number samples. Prior to each use, the hood was UV-sterilized for 4 hours and all  
18 surfaces were subsequently sterilized with sodium hypochlorite. Nucleic acids-free disposable  
19 plastics were used throughout and autoclaved before use. All solutions were prepared in fresh  
20 double quartz-distilled water, sterile filtered, and autoclaved. As a control for contamination  
21 during DNA-extraction, two procedural blanks without sediment were subjected to the whole  
22 extraction and purification procedure along with the sediment samples. One µl of each of these  
23 extraction controls (corresponding to the average volume of sediment extracts used for  
24 amplifications) was included in subsequent PCR amplifications. As additional controls, each  
25 amplification included reactions without DNA template to independently control for  
26 contamination of PCR reagents.

## 1 *Pigment analysis*

2 Pigments were extracted in the dark from 10 grams of freeze-dried and finely ground sediment  
3 samples. Acetone/methanol (7/2, v/v), acetone, and dichloromethane were used for successive  
4 extractions, followed by alkaline hydrolysis of the sediment with 6% KOH in methanol (Glaeser  
5 *et al.*, 2002). To the latter, distilled water and dichloromethane were added and after phase  
6 separation, the organic phases were combined and concentrated by rotary evaporation. The  
7 extracts were subsequently dried under a flow of nitrogen. Individual pigments were quantified  
8 by reverse-phase HPLC (Glaeser *et al.*, 2002) employing a NovaPak C<sub>18</sub> end-capped 60Å 4 µm  
9 4.6 × 250 mm column (Waters) with a Spherisorb 5 ODS 2 4.6 × 10 mm guard column (Waters).  
10 The detection limit of this method is 1.0 ng isorenieratene·(g dry weight sediment)<sup>-1</sup>.

11 For mass spectroscopy, the residue was redissolved in dichloromethane and applied to a  
12 silica column. The apolar carotenoids were eluted with dichloromethane, dried under nitrogen  
13 and the residue dissolved in acetone. This carotenoid fraction was then immediately analyzed on  
14 a HP 1100 series LC/MS equipped with an auto-injector and photodiode array detector.  
15 Separation was achieved on a ZORBAX Eclipse XDB-C<sub>18</sub> column (2.1 × 150 mm, 5 µm; Agilent  
16 Technologies, USA), maintained at 25°C, with a linear gradient from 100% solvent A to 100%  
17 solvent B in 50 min at a flow rate of 0.6 ml·min<sup>-1</sup>. Solvent A was methanol/water (4:1, v/v) and  
18 solvent B acetone/methanol/water (19:1:1, v/v/v). Total run time was 60 minutes. Detection was  
19 achieved by in-line UV-detection (250–700 nm) and positive ion APCI (Atmospheric Pressure  
20 Chemical Ionization) of the eluent in either scanning or Single Ion Monitoring (SIM) mode.  
21 Conditions for APCI-MS were as follows: nebulizer pressure 60 psi, vaporizer temperature  
22 325°C, drying gas (N<sub>2</sub>) flow 7 L·min<sup>-1</sup> and temperature 350°C, capillary voltage 3000V, corona 4  
23 µA. Positive ion spectra were generated by scanning *m/z* 100-1000. In SIM mode *m/z* 529  
24 (protonated molecule of isorenieratene) was monitored. Isorenieratene was quantified by  
25 comparing its UV response at 454 nm (the  $\lambda_{\max}$  of isorenieratene in the mobile phase) to known  
26 amounts of an authentic  $\beta$ -carotene standard (Aldrich) and correcting for the difference in  
27 extinction coefficients (Britton, 1995).

## 1 *Extraction of genomic DNA*

2 Each sediment sample was aliquoted in ten 2 ml bead-beat vials and 0.7 g of glass beads  
3 (0.1 mm diameter) and 0.9 ml lysis buffer (100 mM Tris-HCl, 500 mM sodium-EDTA and  
4 1 wt% SDS; pH 8.0) were added to each vial. The sediment samples were pre-heated for 5  
5 minutes at 70°C in a water bath. Cell-lysis was accomplished by bead-beating at 5,000 rpm for  
6 80 sec (Biospec Mini Bead-Beater; Bartlesville, Oklahoma, USA) followed by another  
7 incubation at 70°C for 30 min. Following centrifugation of the samples in a microfuge (2 min;  
8 14,000 rpm), the supernatants were transferred to a 45 ml teflon centrifuge tube. For extraction  
9 of extracellular, adsorbed DNA, each pellet was resuspended in 0.8 ml of 0.12 M sodium  
10 phosphate buffer (pH 8.0) containing 1% wt/vol SDS, followed by a second round of preheating,  
11 bead-beating and heating. Following centrifugation, the supernatants were recovered. Finally,  
12 each subsample was washed three times with 0.8 ml of 0.12 M sodium phosphate buffer (without  
13 SDS). For each sediment, the supernatants of all subsamples were pooled, yielding a total  
14 volume of about 40 ml.

15 Organic carbon in Mediterranean sapropels contains long chains of polymethylenic carbon  
16 (Petsch *et al.* 2001), which copurified with the genomic DNA and inhibited subsequent PCR. For  
17 removal of the kerogen, several consecutive purification steps were required. Most of the  
18 kerogen could be removed from the crude DNA extracts by adding 3 grams of autoclaved, acid  
19 washed polyvinylpolypyrrolidone (PVPP; Zhou *et al.*, 1996) to each sample. After an incubation  
20 for 20 min on a rotary shaker, the samples were centrifuged for 20 min at 20,400 x g (Beckman  
21 J2-HS, München, Germany), and the supernatants were transferred to a new sterile centrifuge  
22 tube. Extractions with phenol, phenol/chloroform/isoamylalcohol, and chloroform followed  
23 (Sambrook *et al.*, 1989) and the DNA was recovered by standard ethanol precipitation. Finally,  
24 the resulting pellet was washed with sterile double distilled water using a Centricon 50  
25 ultrafiltration unit (Amicon; Witten, Germany) and purified with the Wizard PCR-preps DNA-  
26 purification kit (Promega, Mannheim, Germany). The DNA concentrations in the extracts were  
27 quantified by fluorescent dye binding with PicoGreen (MoBiTec, Göttingen, Germany).

## 1 *Amplification of 16S rRNA genes of green sulfur bacteria*

2 16S rRNA sequences of green sulfur bacteria were selectively amplified with eubacterial primer  
3 341f and the group-specific primer GSB822r, using the previously published cycling conditions  
4 (Overmann *et al.*, 1999). The phylum green sulfur bacteria consists of a crown group of closely  
5 related green sulfur bacteria *sensu strictu* (the family *Chlorobiaceae*) as well as an increasing  
6 number of deep-branching phylotypes which are all uncultured and hence of unknown  
7 physiology. However, all members of the crown group investigated to date are typical  
8 unicellular, obligate photolithoautotrophs. We reassessed the specificity of primer GSB822 by  
9 also including the novel database entries for 16S rRNA gene sequences. Of a total of 86  
10 sequences of *Chlorobiaceae* tested, 11 showed a base substitution (A for T) at the 3'-end of the  
11 target sequence: 5'-AATACTAGATGTTGG(A instead of T)CAT-3'. In three other phylotypes,  
12 different base substitutions were found. All but one of these sequences fell into group 4a (Fig. 5),  
13 while the remaining was a member of group 3. In contrast, all truly marine phylotypes contained  
14 the probe target sequence. Based on this analysis, the PCR method was therefore found to be  
15 suitable to detect in particular the marine members of the green sulfur bacteria.

16 For separation by DGGE, a 40 bp-long GC-clamp has to be added at one end of the PCR  
17 product during amplification, employing primer GC 341f which carries the GC clamp (5'-  
18 CGCCCGCCGCGCCCCGCGCCCGGCCCGCCGCCCCCGCCCC-3') at its 5'-end (Muyzer *et*  
19 *al.*, 1998)). Initial experiments revealed a significantly reduced detection limit of green sulfur  
20 bacterial 16S rRNA genes when GC341f was employed directly for amplification. Consequently,  
21 the GC clamp was added by performing 15 additional cycles with 1 µl of the primary  
22 amplification and with 0.5 pmoles each of primers GC341f and GSB822r.

23 In order to control for the specificity of the amplification conditions, genomic DNA of  
24 *Chlorobium phaeobacteroides* strain MN1 (Overmann *et al.*, 1992) and *Chl. phaeovibrioides*  
25 DSMZ 269<sup>T</sup> were used as positive controls. Since the *Bacteroidetes* represent the sister group of  
26 the phylum green sulfur bacteria, DNA of *Cytophaga* sp. strain 2b served as a negative control.  
27 Purity of the PCR reagents was checked by including two reactions without DNA template in

1 each amplification trial. Possible contaminations during DNA extraction were checked by  
2 including reactions spiked with one  $\mu$ l of each of the extraction controls (see above).

### 3 *Denaturing gradient gel electrophoresis (DGGE)*

4 PCR-products were separated by DGGE (Muyzer *et al.*, 1998) in a Bio-Rad D Gene system  
5 (Biorad, München, Germany). PCR samples were applied directly onto 6% (wt/vol)  
6 polyacrylamide gels (acrylamide/N,N'-methylene bisacrylamide ratio, 37:1 [w/w]) in 1 x TAE  
7 buffer (pH 7.4) which had been prepared from sterile solutions and were cast aseptically between  
8 sterilized glass plates. The gels contained a linear gradient of 30% to 70% denaturant (100%  
9 denaturant correspond to 7 M urea plus 40% [v/v] formamide). Electrophoresis proceeded for 5 h  
10 at 200 V and 60°C. Afterwards, gels were stained for 20 min with sterile ethidium bromide  
11 solution and photographed. Finally, individual fragments were excised with a sterile scalpel,  
12 the DNA was eluted in sterile 1 x TAE (pH 8.0) by electrophoresis (3 h, 200 V) in Centricon 50  
13 concentrators inserted into a Centrilutor Micro Electroelutor (Amicon, Witten, Germany).

### 14 *Sequencing and phylogenetic analysis*

15 One  $\mu$ l of the eluted DNA was reamplified with primers 341f and GSB840r. Primers and  
16 deoxyribonucleoside triphosphates were removed using the QIAquick PCR Purification Spin Kit  
17 (Qiagen, Hilden, Germany). After cycle sequencing with the SequiTherm EXCEL Long-Read  
18 Sequencing Kit-LC (Biozym, Hess. Oldendorf, Germany) and employing primers 341f and  
19 GSB840r, sequence data were collected with a LiCor-4000 automated sequencer (Lincoln,  
20 Nebraska, USA).

21 Each sequence was checked for chimeras employing the CHECK\_CHIMERA option of the  
22 ribosomal database project (RDP). The 16S rRNA gene sequences were then analyzed using the  
23 ARB phylogeny software package (Ludwig *et al.*, 2004). The Fast Aligner V1.03 tool was used  
24 for automatic alignment and the resulting alignments were corrected based on the 16S rRNA  
25 secondary structure information for *Chlorobium vibrioforme* DSMZ 260<sup>T</sup>, as available through  
26 The Comparative RNA Web (CRW) Site ([www.rna.icmb.utexas.edu](http://www.rna.icmb.utexas.edu); Cannone *et al.*, 2002).

1 Phylogenetic trees were constructed including 16S rRNA gene sequences of available strains and  
2 environmental sequences. First, sequences longer than 1100 bp were used for the calculation,  
3 employing the MAXIMUM LIKELIHOOD algorithm (Fast DNA\_ML). The shorter  
4 environmental sequences were inserted afterwards without changing overall tree topology  
5 employing the PARSIMONY INTERACTIVE tool implemented in the ARB software package.

6 The 16S rRNA gene sequences obtained during the present study have been deposited in  
7 GenBank under accession numbers AF298531-AF298537.

### 8 *Quantification of DNA of green sulfur bacteria by dot blot hybridization*

9 For dot blot quantification, 16S rRNA gene sequences of green sulfur bacteria were amplified  
10 with an altered PCR protocol. Instead of a step down, primer annealing was performed at a  
11 constant temperature of 52°C for 40 s and 31 cycles. Samples were denatured for 7 min at 100°C  
12 and vacuum blotted onto positively charged nylon membranes (Boehringer Mannheim,  
13 Germany). The membrane was baked (25 min at 120°C) and pre-hybridized in 10 ml of DIG  
14 Easy Hyb buffer (Boehringer). Afterwards, 150 pmole of a highly specific probe for green sulfur  
15 bacteria (5'- TGCCACCCCTGTATC-3'; *E. coli* positions 532 to 546; Tuschak *et al.*, 1999), 5'  
16 labeled with dig-11-dUTP (MWG-Biotech, Germany), was added and hybridization was carried  
17 out for 12 h. After hybridization, the blot was washed twice for 5 min in 2 x SSC (150 mM  
18 NaCl, 15 mM Na-citrate, pH 7.0) plus 0.1% SDS, followed by two stringent washing steps  
19 (15 min in 0.1xSSC/0.1% SDS). The hybridization signal was detected by chemiluminescence  
20 with the DIG Luminescent Detection Kit (Boehringer). Lumi-Film (Boehringer) was exposed for  
21 30 min, developed and the image digitized with a flatbed scanner. For quantification of the  
22 individual dots, the ZERO-Dscan software (Scanalytics, Billerica, USA) was employed.  
23 Genomic DNA of *Chlorobium phaeovibrioides* DSMZ 269<sup>T</sup> was used for calibration. The  
24 homology of the 16S rRNA gene sequences of other Bacteria at the target site of probe 532 is  
25 low, which reflects the large phylogenetic distance of green sulfur bacteria to other bacteria  
26 (Overmann and Tuschak, 1997). Therefore we chose DNA of *Cytophaga* strain 2b (which,

1 however shows 5 mismatches) as a negative control for the specificity of the hybridization. For  
 2 each sample, quantification was conducted in three independent parallels. The quantification was  
 3 based on the assumption that genome sizes of green sulfur bacteria and the number of rRNA  
 4 operons are similar. Based on the currently available information, these assumptions appear to be  
 5 largely valid since genome sizes of green sulfur bacteria show comparatively little variation in  
 6 length (1.97 to 3.13 Mb) and typically comprise one (33% of the genomes) or two (66% of the  
 7 genomes) *rrn* operons ([http://genome.jgi-psf.org/draft\\_microbes/](http://genome.jgi-psf.org/draft_microbes/)).

#### 8 *Adsorption capacity for DNA of the sapropels*

9 In order to study the potential protection of fossil DNA by adsorption, equilibrium adsorption  
 10 isotherms were determined. Herring sperm DNA (Boehringer Mannheim) at concentrations  
 11 between 1 and 10  $\mu\text{g}\cdot\text{ml}^{-1}$  was incubated for 12 hours with aliquots (5 mg dry weight $\cdot\text{ml}^{-1}$ ) of  
 12 sapropel S<sub>6</sub> and the intermediate layer Z<sub>6</sub>. Adsorption of DNA was observed to reach a  
 13 maximum and therefore yielded the best fit with the Langmuir equation (Ogram *et al.*, 1987)  
 14 according to:

$$15 \quad S = S_{\max} \cdot \frac{K \cdot C_e}{(1 + K \cdot C_e)} \quad (1)$$

16 Equation (1) was fitted to the data points of the amount  $S$  of DNA adsorbed ( $\mu\text{mol}\cdot(\text{g dry wt})^{-1}$ )  
 17 and the concentration  $C_e$  of the substance remaining in solution (in  $\mu\text{mol}\cdot\text{ml}^{-1}$ ). This yielded an  
 18 estimate for the maximum adsorption capacity  $S_{\max}$  ( $\mu\text{mol}\cdot(\text{g dry wt})^{-1}$ ) and the Langmuir affinity  
 19 coefficient  $K$  ( $\text{ml}\cdot\mu\text{mol}^{-1}$ ). For comparison, equilibrium adsorption isotherms of DNA for  
 20 montmorillonite and cellulose were also measured.

21

**1 ACKNOWLEDGEMENTS**

2 We thank Daniel Dotschkal for help with the extraction of carotenoids and Jens Glaeser and  
3 Ellen Hopmans for assistance during HPLC determinations. Andrea Smock, Henrik Sass, and  
4 Heribert Cypionka are gratefully acknowledged for their support during the cruise and helpful  
5 discussions, and the master and the crew of the R/V *Meteor* for their help during collection of the  
6 sediment cores. Thanks to Ann Manske for help with the phylogenetic analyses. This work was  
7 funded by the Deutsche Forschungsgemeinschaft (grants Ov 20/3-2 and Ov 20/8-1 to 8-3).

8

9

1 **REFERENCES**

- 2 Barnes, I., Matheus, P. Shapiro, B., Jensen, D., and Cooper, A. (2002) Dynamics of Pleistocene  
3 population extinctions in Beringian brown bears. *Science* **295**: 2267-2270
- 4 Britton, G. (1995) UV/Visible Spectroscopy. In: Britton, G., Liaaen-Jensen, S., Pfande, H. (eds.)  
5 *Carotenoids, vol. 1B: Spectroscopy*. Birkhäuser Verlag, Basel.
- 6 Calvert, S.E. (1983) Geochemistry of the Pleistocene sapropels and associated sediments from  
7 the eastern Mediterranean. *Oceanol Acta* **6**: 225-267
- 8 Calvert, S.E., B. Nielsen, and Fontugne, M.R. (1992) The role of mat-forming diatoms in the  
9 formation of Mediterranean sapropels. *Nature* **359**: 223-225
- 10 Cannone, J. J., Subramanian, S., Schnare, M. N., Collett, J. R., D'Souza, L. M., Du, Y., *et al.*  
11 (2002) The Comparative RNA Web (CRW) Site: An Online Database of Comparative  
12 Sequence and Structure Information for Ribosomal, Intron, and other RNAs. *BioMed*  
13 *Central Bioinformatics* **3**: 2 [Correction: *BioMed Central Bioinformatics* **3**: 15]
- 14 Coolen, M.J.L. and Overmann, J. (1998) Analysis of subfossil molecular remains of purple  
15 sulfur bacteria in a lake sediment. *Appl Environ Microbiol* **64**: 4513-4521
- 16 Coolen, M.J.L., and Overmann, J. (2000) Functional exoenzymes as indicators of metabolically  
17 active bacteria in up to 124,000-year-old sapropel layers of the eastern Mediterranean Sea.  
18 *Appl Environ Microbiol* **66**: 2589-2598
- 19 Coolen, M.J.L., Cypionka, H., Smock, A., Sass, H., and Overmann, J. (2002) Ongoing  
20 modification of Mediterranean Pleistocene sapropels mediated by prokaryotes. *Science* **296**:  
21 2407-2410
- 22 Coolen, M. J. L., Boere, A., Abbas, B., Baas, M., Wakeham, S. G., and Sinninghe Damsté, J. S. ,  
23 (2006) Fossil DNA derived from alkenone-biosynthesizing haptophytes and other algae in  
24 Holocene sediment from the Black Sea, *Paleoceanography* **21**: PA1005-  
25 doi:10.1029/2005PA001188.
- 26 Cooper, A., and Poinar, H.N. (2000) Ancient DNA: do it right or not at all. *Science* **289**: 1
- 27 Crecchio, C., and Stotzky, G. (1998) Binding of DNA on humic acids: effect on transformation  
28 of *Bacillus subtilis* and resistance to Dnase. *Soil. Biol. Biochem.* **30**, 1061-1067
- 29 Emeis, K.-C., Sakamoto, T., Wehausen, R., and Brumsack, H.-J. (2000) The sapropel record of  
30 the eastern Mediterranean Sea: Results of Ocean Drilling Program Leg 160. *Palaeogeogr*  
31 *Palaeoclimatol Palaeoecol* **158**: 371-395

- 1 Glaeser, J., Baneras, L., Rütters, H., and Overmann, J. (2002) Novel bacteriochlorophyll *e*  
2 structures and species-specific variability of pigment composition in green sulfur bacteria.  
3 *Arch Microbiol* **177**: 475-485
- 4 Hofreiter, M., Serre, D., Poinar, H.N., Kuch, M., and Pääbo, S. (2001) Ancient DNA. *Nature Rev*  
5 *Gen* **2**: 353-359
- 6 Imhoff, J.F. (2003) Phylogenetic taxonomy of the family *Chlorobiaceae* on the basis of 16S  
7 rRNA and *fmo* (Fenna-Matthews-Olson protein) gene sequences. *Int J Syst Evol Microbiol*  
8 **53**: 941
- 9 Jones, D.L., and Edwards, A.C. (1998) Influence of sorption on the biological utilization of two  
10 simple carbon substrates. *Soil Biol Biochem* **30**: 1895-1902
- 11 Keil, R.G., Montluçon, D.B., Prahl, F.G., and Hedges, J.I. (1994) Sorptive preservation of labile  
12 organic matter in marine sediments. *Nature* **370**: 549-552
- 13 Kidd, R.B., Cita, M.B., and Ryan, W.B.F. (1978) Stratigraphy of eastern Mediterranean sapropel  
14 sequences recovered during DSDP Leg 42A and their paleoenvironmental significance. *Init*  
15 *Reports DSDP* **42**: 421-443
- 16 Krubasik, P., and Sandmann, G. (2000) A carotenogenic gene cluster from *Brevibacterium linens*  
17 with novel lycopene cyclase genes involved in the synthesis of aromatic carotenoids. *Mol*  
18 *Gen Genet* **263**: 423-432
- 19 Krügel, H., Krubasik, P., Weber, K., Saluz, H.P., and Sandmann, G. (1999) Functional analysis  
20 of genes from *Streptomyces griseus* involved in the synthesis of isorenieratene, a carotenoid  
21 with aromatic end groups, revealed a novel type of carotenoid desaturase. *Biochim Biophys*  
22 *Acta* **1439**: 57-64
- 23 Lindahl, T. (1993) Instability and decay of the primary structure of DNA. *Nature* **362**: 709-715
- 24 Lorenz, M.G., and Wackernagel, W. (1994) Bacterial gene transfer by natural genetic  
25 transformation in the environment. *Microbiol Rev* **58**: 563-602
- 26 Ludwig, W., Strunk, O., Westram, R., Richter, L., Meier, H., Yadhukumar, A. *et al.* (2004)  
27 ARB: a software environment for sequence data. *Nucleic Acids Res* **32**:1363-1371.
- 28 Manske, A.K., Glaeser, J., Kuypers, M.M.M., and Overmann, J. (2005) Physiology and  
29 phylogeny of green sulfur bacteria forming a monospecific phototrophic assemblage at 100  
30 m depth in the Black Sea. *Appl Environ Microbiol* **71**: 8049-8060

- 1 Menzel, D., Hopmans, E.C., van Bergen, P.F., de Leeuw, J.W., and Sinninghe Damsté, J.S.  
2 (2002) Development of photic zone euxinia in the eastern Mediterranean Basin during  
3 deposition of Pliocene sapropels. *Mar. Geol.* **189**: 215-226
- 4 Muyzer, G., Brinkhoff, T., Nübel, U., Santegoeds, C., Schäfer, H., and Wawer, C. (1998)  
5 Denaturing gradient gel electrophoresis (DGGE) in microbial ecology. In Akkermans,  
6 A.D.L., van Elsas, J.D., and de Bruijn, F.J. (eds.), *Molecular microbial ecology manual*, vol.  
7 3.4.4. Kluwer Academic Publishers, Dordrecht, The Netherlands. p. 1 - 27
- 8 Ogram, A., Sayler, G.S., and Barkay, T. (1987) The extraction and purification of microbial  
9 DNA from sediments. *J Microbiol Methods* **7**: 57-66
- 10 Overmann, J. (2001) Green Sulfur Bacteria. In: Boone, D.R., Castenholz, R.W. (eds) *Bergey's*  
11 *Manual of Systematic Bacteriology*, 2<sup>nd</sup> ed., Vol. 1. Williams and Wilkins, Baltimore, pp.  
12 601-630
- 13 Overmann, J., Coolen, M.J.L., and Tuschak, C. (1999) Specific detection of different groups of  
14 chemocline bacteria based on PCR and denaturing gradient gel electrophoresis of 16S rRNA  
15 gene fragments. *Arch Microbiol* **172**: 83-94
- 16 Overmann, J., Cypionka, H., and Pfennig, N. (1992) An extremely low-light adapted  
17 phototrophic sulfur bacterium from the Black Sea. *Limnol Oceanogr* **37**: 150-155
- 18 Overmann, J., and Tuschak, C. (1997) Phylogeny and molecular fingerprinting of green sulfur  
19 bacteria. *Arch Microbiol* **167**: 302-309
- 20 Paget, E., Monrozier, L.J., and Simonet, P. (1992) Adsorption of DNA on clay minerals:  
21 protection against DNase I and influence on gene transfer. *FEMS Microbiol Lett* **97**: 31-40
- 22 Passier, H.F., Bosch, H.J., Nijenhuis, I.A., Lourens, L.J., Böttcher, M.E., Leenders, A. et al.  
23 (1999) Sulphidic Mediterranean surface waters during Pliocene sapropel formation. *Nature*  
24 **397**: 146-149
- 25 Petsch, S. T., Eglinton, T. I., and Edwards, K. J. (2001) <sup>14</sup>C-dead living biomass: evidence for  
26 microbial assimilation of ancient organic carbon during shale weathering. *Science* **292**: 1127
- 27 Pfennig, N., and Trüper, H.G. (1989) Anoxygenic phototrophic bacteria. In: Staley J.T., M.P.  
28 Bryant, N. Pfennig and J.G. Holt (eds.): *Bergey's Manual of Systematic Bacteriology*, Vol.  
29 III, Williams and Wilkins, Baltimore, pp. 1635-1709
- 30 Poinar, H.N., Höss, M., Bada, J.L., and Pääbo, S. (1996) Amino acid racemization and the  
31 preservation of ancient DNA. *Science* **272**:864-866
- 32 Poinar, H., Kuch, M., McDonald, G., Martin, P., and Pääbo, S. (2003) Nuclear gene sequences  
33 from a late Pleistocene sloth coprolite. *Curr Biol* **13**: 1150-1152

- 1 Rohling, E.J., and Hilgen, F.J. (1991) The eastern Mediterranean climate at times of sapropel  
2 formation: a review. *Geol Mijnbouw* **70**: 253-264
- 3 Rollo, F. (1998) Ancient DNA: Problems and perspectives for molecular microbial  
4 palaeoecology. In: Carvalho, C.R. (ed.) *Advances in Molecular Ecology*, IOS Press,  
5 Amsterdam. p. 133-149
- 6 Romanowski, G., Lorenz, M.G., and Wackernagel, W. (1991) Adsorption of plasmid DNA to  
7 mineral surfaces and protection against Dnase I. *Appl Environ Microbiol* **57**: 1057-1061
- 8 Rosell-Melé, A., Maslin, M.A., Maxwell, J.R., and Schaefer, P. (1997) Biomarker evidence for  
9 “Heinrich” events. *Geochim Cosmochim Acta* **61**: 1671-1678
- 10 Rossignol-Strick, M. (1985) Mediterranean Quaternary sapropels, an immediate response of the  
11 African Monsoon to variation of insolation. *Paleogeogr Paleoclimatol Paleocol* **49**: 237-  
12 263
- 13 Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989) *Molecular cloning: A laboratory manual*.  
14 2nd edition, Cold Spring Harbor Laboratory Press, New York
- 15 Schmiedl, G. Hemleben, C., Keller, J., and Segl, M. (1998) Impact of climatic changes on the  
16 benthic foraminiferal fauna in the Ionian Sea during the last 330,000 year.  
17 *Paleoceanography* **13**: 447-458
- 18 Shapiro, R. (1981) Damage to DNA caused by hydrolysis. In: Seeberg, E., and Kleppe, K. (eds)  
19 *Chromosome Damage and Repair*, pp. 3-18, Plenum Press
- 20 Tuschak, C., Glaeser, J., and Overmann, J. (1999) Specific detection of green sulfur bacteria by  
21 *in situ* hybridization with a fluorescently labeled oligonucleotide probe. *Arch Microbiol* **171**:  
22 265-272
- 23 Willerslev, E., Hansen, A.J., Binladen, J., Brand, T.B., Gilbert, M.T.P., Shapiro, B., Bunce, M.,  
24 Wiuf, C., Gilichinsky, D.A., and Cooper, A. (2003) Diverse plant and animal genetic  
25 records from Holocene and Pleistocene sediments. *Science* **300**: 791-795
- 26 Willerslev, E., Hansen, A.J., Poinar, H.N. (2004a) Isolation of nucleic acids and cultures from  
27 fossil ice and permafrost. *TRENDS Ecol Evol* **19**: 141-147
- 28 Willerslev, E., Hansen, A.J., Rønn, R., Brand T.B., Barnes, I., Wiuf, C., Gilichinsky, D.,  
29 Mitchell, D., Cooper, A. (2004b) Long-term persistence of bacterial DNA. *Curr Biol* **14**:  
30 R9-R10
- 31 Zhou, J., Bruns, M.A., and Tiedje, J.M. (1996) DNA recovery from soils of diverse composition.  
32 *Appl Environm Microbiol* **62**: 316-322

1 **FIGURE LEGENDS**

2 **Fig. 1. A.** Total DNA content (●) at eight consecutive depths of core #69-2SL. The vertical  
 3 positions of the sapropel layers (black rectangles) and of sampling depths are denoted on  
 4 the right. **B.** DNA of green sulfur bacteria as fraction of total DNA (■). **C.** Absolute  
 5 amounts of green sulfur bacterial DNA as determined by dot blot quantification (▲) and  
 6 of the carotenoid isorenieratene (○). Horizontal bars indicate one standard deviation.

7 **Fig. 2.** Amplification products generated with primers GC341f and GSB822. Controls for  
 8 contamination of PCR reagents (no template), for contamination with extraneous DNA  
 9 (extraction controls, see Experimental procedures) and for primer specificity (*Cytophaga*  
 10 sp. strain 2b) were included in each PCR run. *Chlorobium phaeovibrioides* DSMZ 269<sup>T</sup>  
 11 and *Chlorobium phaeobacteroides* strain MN1 served as positive controls. A negative  
 12 image of an ethidium bromide-stained agarose gel is shown.

13 **Fig. 3.** Dot blot quantification of fossil DNA after amplification of 2 ng each of sample DNA.  
 14 The standard curve was obtained by using different amounts (given in pg) of genomic  
 15 DNA of *Chlorobium phaeovibrioides* DSMZ 269<sup>T</sup> as template. As negative controls, two  
 16 independent PCR amplifications were conducted either without DNA template (n.t.), with  
 17 1 µl of the extraction controls (Extr., compare Experimental Procedures), or with 2 ng of  
 18 genomic DNA of *Cytophaga* strain 2b (*Cyt*).

19 **Fig. 4.** Separation of green sulfur bacterial 16S rRNA gene fragments by DGGE. For  
 20 comparison, amplification products from the two brown-colored marine strains  
 21 *Chlorobium phaeovibrioides* DSMZ 269<sup>T</sup> and *Chl. phaeobacteroides* MN1 (Overmann *et*  
 22 *al.*, 1992) were included. DNA bands used in the subsequent phylogenetic analyses are  
 23 denoted by labels A through G. Asterisks denote sequences which did not yield  
 24 unambiguous sequences. Six different melting types are marked with arrows.

25 **Fig. 5.** Phylogenetic position of the seven fossil partial 16S rRNA gene sequences of green  
 26 sulfur bacteria (given in bold face; labeling refers to layer of origin plus the band no.  
 27 according to Fig. 4) recovered from sapropels and intermediate layers. Currently  
 28 recognized groups are indicated (Imhoff, 2003; denoted in rectangles). Groups comprising  
 29 salt tolerant or salt-requiring strains marked by black rectangles. Bar indicates 0.1 fixed  
 30 point mutations per nucleotide. Numbers at nodes give bootstrap values out of 100  
 31 resamplings for phylogenetic trees calculated by Maximum Likelihood / Maximum  
 32 Parsimony / Neighbor Joining methods. *C.*, *Chlorochromatium*; *Cba.*, *Chlorobaculum*,  
 33 *Chl.*, *Chlorobium*; *Chp.*; *P.* *Pelochromatium*; *Pld.*, *Pelodictyon*.

34

1 **Fig. 6.** Equilibrium adsorption isotherms of double-stranded herring sperm DNA bound to  
2 sapropel S6 (●) and the intermediate layer Z6 (○). Bars indicate one standard deviation.  
3 Lines indicate the curves fitted to the data according to the Langmuir equation (eq. 1, see  
4 text).

5

1 **Table 1.** Parameters of the Langmuir equilibrium adsorption isotherms for DNA binding to  
 2 different adsorbers  
 3

Adsorber	$S_{\max}^1$ $\mu\text{mol}\cdot(\text{g dry wt})^{-1}$	K $(\text{ml}\cdot\mu\text{mol}^{-1})$	Reference
Humic acids (pH 4)	$250.9 \pm 72.4$	$7.2 \pm 3.8$	Crecchio and Stotzky, 1998
Humic acids (pH 3)	$112.1 \pm 19.1$	$9.2 \pm 4.4$	Crecchio and Stotzky, 1998
Sapropel S <sub>6</sub> (50 mM MgCl <sub>2</sub> )	$79.3 \pm 2.8$	$447 \pm 63$	this work
Intermediate layer Z <sub>6</sub> (50 mM Tris-HCl) <sup>2</sup>	$18.0 \pm 1.9$	$209 \pm 116$	this work
Ca <sup>2+</sup> -montmorillonite (0.5 mM CaCl <sub>2</sub> )	$64.0 \pm 12.1$	$286 \pm 129$	Paget <i>et al.</i> , 1992
Mg <sup>2+</sup> -montmorillonite (50 mM MgCl <sub>2</sub> )	$59.6 \pm 6.3$	$41.8 \pm 16.6$	Barghorn and Overmann, unpubl.
Mg <sup>2+</sup> -montmorillonite (2.0 mM MgCl <sub>2</sub> )	$14.2 \pm 0.7$	$4099 \pm 1063$	Paget <i>et al.</i> , 1992
Mg <sup>2+</sup> -montmorillonite (0.5 mM MgCl <sub>2</sub> )	$12.6 \pm 0.7$	$293 \pm 48$	Paget <i>et al.</i> , 1992
Na <sup>+</sup> -montmorillonite (25 mM NaCl)	$1.76 \pm 1.18$	$451 \pm 706$	Paget <i>et al.</i> , 1992
Mg <sup>2+</sup> -cellulose (50 mM MgCl <sub>2</sub> )	$0.755 \pm 0.103$	$74.6 \pm 23.1$	Barghorn and Overmann, unpubl.
Memphis-soil	0.023	673	Ogram <i>et al.</i> , 1987

4  
 5 <sup>1</sup> Molarity of DNA calculated as molarity of nucleotides.

6 <sup>2</sup> Due to precipitation in the presence of Mg<sup>++</sup>, adsorption was determined in 50 mM Tris-HCl.

7




Fig.1  
Coolen and Overmann (2006)




Fig. 2  
Coolen and Overmann (2006)




Fig. 3  
Coolen and Overmann (2006)




Fig. 4  
Coolen and Overmann (2006)




Fig. 5  
Coolen and Overmann (2006)




Fig. 6  
Coolen and Overmann (2006)