

SHORT COMMUNICATION

Carbon content of *Mnemiopsis leidyi* eggs and specific egg production rates in northern Europe

Cornelia Jaspers^{1*}, John H. Costello^{2,3} and Sean P. Colin^{2,4}

¹DTU AQUA, TECHNICAL UNIVERSITY OF DENMARK, KAVALERGÅRDEN 6, 2920 CHARLOTTENLUND, DENMARK

²MARINE BIOLOGICAL LABORATORY, 7 MBL STREET, WOODSHOLE, MA 02543, USA

³BIOLOGY DEPARTMENT, PROVIDENCE COLLEGE, PROVIDENCE, RI 02918, USA

⁴DEPARTMENT OF MARINE BIOLOGY AND ENVIRONMENTAL SCIENCES, ROGER WILLIAMS UNIVERSITY, BRISTOL, RI 02809, USA

*CORRESPONDING AUTHOR: coja@aqua.dtu.dk

Current address: Helmholtz Centre for Ocean Research, Kiel, GEOMAR, Düsternbrooker Weg 20, 24105 Kiel, Germany

KEYWORDS: Comb jelly, ctenophore, invasive species

RUNNING HEAD: *M. leidyi* egg carbon and specific egg production rates

The comb jelly *Mnemiopsis leidyi* is regarded to be a successful invasive species, partly due to its high reproduction potential. However, due to the absence of direct carbon measurements of eggs, specific reproduction rates remain uncertain. We show that egg-carbon is $0.22 \pm 0.02 \mu\text{gC}$ and up to 21 times higher than previously extrapolated. With maximum rates of $11,232 \text{ eggs ind}^{-1} \text{ d}^{-1}$, largest animals in northern Europe invest c. $10\% \text{ d}^{-1}$ of their body carbon into reproduction.

The comb jelly *Mnemiopsis leidyi* attained large public and scientific attention during the last decades due to its commonly observed formation of bloom abundances in native and invaded areas (e.g. Costello *et al.*, 2012; Riisgård *et al.*, 2012). One of the traits suggested to be responsible for *M. leidyi*'s invasion success is its high fecundity. At the northern end of its distribution range in native areas, *M. leidyi* has been shown to produce up to 9,380 and $14,233 \text{ eggs ind}^{-1} \text{ d}^{-1}$ (Kremer, 1976a; Graham *et al.*, 2009), with similar rates of $9,910 \text{ eggs ind}^{-1} \text{ d}^{-1}$ for the native southern population in Biscayne Bay, Florida (Baker and Reeve, 1974). Within invaded European waters, rates of up to

3,000 and 12,000 eggs $\text{ind}^{-1} \text{ d}^{-1}$ have been recorded for northern and southern populations, respectively (Zaika and Revkov, 1994; Javidpour *et al.*, 2009). Since *M. leidyi* is a simultaneous hermaphrodite and fertilized eggs are produced on a daily basis during favourable conditions (Jaspers, 2012), *M. leidyi* can circumvent the Allee effect and efficiently seed new populations even from few founding individuals. Although the documented reproduction potential is large, carbon investment into reproduction remains speculative due to the lack of direct carbon measurements of eggs. Literature values of carbon concentration of eggs vary by a factor of 21, from 0.012 $\mu\text{gC egg}^{-1}$ (Reeve *et al.*, 1978) to 0.25 $\mu\text{gC egg}^{-1}$ (Anninsky *et al.*, 2007) but have so far never been measured directly. This leads to the discrepancy between high feeding rates (Colin *et al.*, 2010) and negligible investment into reproduction with 0.03 to <2% of *M. leidyi*'s body carbon per day, which is several fold less than its respiratory carbon demand (Reeve *et al.*, 1989). The aim of this study is to present direct carbon and nitrogen measurements of *M. leidyi* eggs to clarify carbon specific reproduction rates along with its maximum reproduction capacity within invaded, northern European waters. This information is crucial for understanding population dynamics and assessing carbon budgets of the comb jelly *M. leidyi* throughout its distribution range.

Carbon and nitrogen concentration of freshly spawned eggs ($n = 2800$) were measured from wild caught *Mnemiopsis leidyi* ($n = 5$) originating from Woods Hole, MA, NE USA (position: Latitude 41.525 N, Longitude -70.674 E). Animals were incubated overnight in GFF filtered seawater. Eggs were individually picked using a micropipette and washed 3 times in GFF filtered seawater before being placed, with as little water as possible, onto pre-combusted, pre-weighted GFF filters. To assure sufficient carbon and nitrogen content for the analyses, eggs were pooled into batches of 200, 300, 400 and 500 eggs per GFF filter. Filters with eggs were dried at 60°C for 3 days, stored in a desiccator and analyzed within 3 days at the Marine Biological Laboratory, USA using a FLASH 2000 NC Analyzer (ThermoFisher Scientific, Cambridge, UK). Blank filters to correct for contribution of filter and seawater were prepared at the same time using equivalent amount of water from the final washing step which was used as background correction. The measurements were repeated on 2 days with independently caught animals ($n = 5$) with replicates for each egg batch of 200 to 500 eggs, respectively.

Egg production rates of *M. leidyi* were measured during late August and early September 2010 in intermediate saline waters of northern Europe (position: Latitude 58.250 N, Longitude 11.447 E - Skagerrak, Gullmar Fjord, Sweden), representing high saline waters of the Baltic Sea region. Freshly caught, >50 mm sized animals (oral-aboral length, $n = 5$) were individually incubated in 7.5 L GFF filtered seawater at ambient salinity (22.5) and temperature (16.5°C) following natural light conditions. After 24 hours of incubation, *M. leidyi* were removed, total and oral-aboral lengths measured and eggs concentrated via reverse filtration. Eggs were

preserved in acidified Lugol solution at a final concentration of 2% for later enumeration under a dissecting microscope. To calculate specific egg production, the regression from total length (TL) to dry weight (DW) from Baker (Baker, 1973), as cited in Kremer et al. (Kremer *et al.*, 1986) was used (1) and the dry weight (DW) to carbon regression based on analyses of table I in Kremer et al. (Kremer *et al.*, 1986) leading to regression (2) with the following regression parameters ($F_{1,7} = 1698$, $p < 0.0001$, $R^2 = 0.997$, $n = 8$).

$$\text{Dry weight (mg)} = 0.038 \times \text{total length (mm)}^{2.42} \quad (1)$$

$$Mnemiopsis \text{ carbon (mg)} = 0.0018 \times \text{DW (mg)}^{1.318} \quad (2)$$

Direct measurement of early cleavage stages of *Mnemiopsis leidyi* eggs show that the carbon and nitrogen concentration is $0.22 \pm 0.02 \mu\text{gC egg}^{-1}$ and $0.07 \pm 0.01 \mu\text{gN egg}^{-1}$, respectively (Fig. 1). The average *M. leidyi* egg size is $503 \pm 58 \mu\text{m}$ ($n = 25$), similar to egg sizes observed in invaded northern Europe of $565 \pm 66 \mu\text{m}$ (Jaspers *et al.*, 2013). Early cleavage stages are depicted in Fig. 2. Largest sized animals found during August/September 2010 were 70 to 78 mm in total length, with an average reproduction rate of $8,432 \pm 1,884 \text{ eggs ind}^{-1} \text{ d}^{-1}$. The highest egg production rate of 11,232 eggs d^{-1} was attained by the largest sized animal (Table I).

Calculation of carbon specific egg production (SEP) shows that ca. 8.5% d^{-1} of the *M. leidyi* body carbon is channeled into reproduction (Table I).

We present the first direct carbon and nitrogen measurements of *Mnemiopsis leidyi* eggs. Previous studies have estimated carbon concentrations from extrapolation of larval carbon measurements or dry weight to carbon conversions of larvae (e.g. Reeve *et al.*, 1978; Reeve *et al.*, 1989; Anninsky *et al.*, 2007). Since no direct measurements of eggs are available, estimates of carbon concentrations used for carbon and energy budgets of *M. leidyi* vary by a factor of 21. The only direct carbon measurement of ctenophore eggs available is $0.56 \mu\text{gC egg}^{-1}$ for 200 μm larger *Bolinopsis mikado* eggs (Kasuya *et al.*, 2008). *M. leidyi* eggs have 37% of the volume of *B. mikado* eggs; however, the volume specific carbon and nitrogen concentrations are similar with 1 and 1.2 times of the *B. mikado* carbon and nitrogen concentrations, respectively. The commonly used carbon concentration for *M. leidyi* of $0.1 \mu\text{gC egg}^{-1}$ (Reeve *et al.*, 1989) is thus factor 2.2 less than expected if compared to the measured carbon content of *B. mikado* eggs (Kasuya *et al.*, 2008). Reeve *et al.* (Reeve *et al.*, 1989) estimated the carbon concentration from the assumption that the dry weight of eggs is 0.5 μg and used extrapolations of carbon measurements of $>1.1 \text{ mm}$ sized larvae to estimate egg carbon using a carbon ratio to dry weight of 20%. The lowest carbon concentration used for *M. leidyi* eggs is based on *Pleurobrachia* carbon and nitrogen measurements of $3.28 \pm 0.35\%$ and $0.87 \pm 0.09\%$ ($\pm \text{SD}$) of dry weights, assuming a *M. leidyi* dry weight of 0.35

$\mu\text{g egg}^{-1}$ (Reeve *et al.*, 1978). Thus, leading to an estimated carbon and nitrogen concentration per *M. leidyi* egg of $0.012 \mu\text{gC egg}^{-1}$ and $0.003 \mu\text{gN egg}^{-1}$. This is ca. 5% of the C and N concentration measured directly in this study. Our measured carbon content of eggs is similar to direct carbon measurements of $0.26 \mu\text{gC}$ for $500\mu\text{m}$ *M. leidyi* larvae from NE USA (Sullivan and Gifford, 2004). In an ecological perspective, *M. leidyi* eggs have the same nutritional value and comparable size range as e.g. copepod nauplii, echinoderm larvae, gastropod veliger and bivalve larvae (Martinussen and Båmstedt, 1995) and might therefore be an important food source for e.g. pelagic filter feeders. Therefore, predation could contribute to high mortality rates of *M. leidyi* eggs observed in intermediate saline waters of northern Europe (Jaspers *et al.*, 2013). However, predator prey investigations are necessary to enlighten direct interactions.

The carbon:nitrogen ratio for eggs measured in this study is 3.1, while the ratio for a large range of *M. leidyi* size classes is consistently around 4 (Kremer, 1976a). This indicates that although the carbon per unit weight of *M. leidyi* changes with size, hence throughout its life (Reeve *et al.*, 1989), the carbon to nitrogen ratio remains constant apart from the egg phase, where the C:N ratio is lower.

Due to the previously underestimated egg carbon concentrations of *M. leidyi*, egg production has so far been suggested to be a small and negligible fraction of the daily carbon demand (Kremer, 1976b; Kremer, 1982; Reeve *et al.*, 1989). Even though several thousand eggs may be produced over a few days, they have been estimated to represent <2% of the carbon biomass of the ctenophores per day (Kremer, 1976b; Reeve *et al.*, 1989), which is less than one third of the respiratory carbon demand (Kremer, 1982). We show that large sized animals produce up to $11,232 \text{ eggs ind}^{-1} \text{ d}^{-1}$ in invaded northern European waters, which is nearly four times more than previously shown for northern Europe (Javidpour *et al.*, 2009) and in the same range as for the Black Sea and other native habitats (Baker and Reeve, 1974; Kremer, 1976a; Graham *et al.*, 2009). Therefore, specific egg production in northern Europe for largest sized animals is around 7 to 10% d^{-1} of the body carbon. This is four times higher than previously documented, due to the underestimation of carbon concentrations of *M. leidyi* eggs. This demonstrates that egg production should be taken into account in energy and carbon budgets of *M. leidyi* in native and invaded habitats.

ACKNOWLEDGEMENTS

We would like to thank Matthew Erickson for running CN analyzes, the Marine Biological Laboratory, Woods Hole, MA for the use of their laboratory facilities for this study and the reviewers for their valuable comments. This work was supported by a grant from the German

Exchange Service (DAAD) to CJ; NSF –grant OCE1061353 to JHC and SPC. Egg production rates have been measured at the Sven Lovén Centre, Sweden and been supported by the ASSEMBLE grant agreement no. 227799 to CJ under support of the European Community - Research Infrastructure Action under the FP7 "Capacities" Specific Programme.

REFERENCES

- Anninsky, B. E., Finenko, G. A., Abolmasova, G. I. and Romanova, Z. A. (2007) Somatic organic content of the ctenophores *Mnemiopsis leidyi* (Ctenophora: Lobata) and *Beroe ovata* (Ctenophora: Beroida) in early ontogenetic stages. *Russ. J. Mar. Biol.*, **33**, 417-424.
- Baker, L. D. (1973) *The ecology of the ctenophore Mnemiopsis mccradyi (Mayer)*, in *Biscayne Bay, Florida*. University of Miami, Miami, Florida, pp. 1-131.
- Baker, L. D. and Reeve, M. R. (1974) Laboratory culture of the lobate ctenophore *Mnemiopsis mccradyi* with notes on feeding and fecundity. *Mar. Biol.*, **26**, 57-62.
- Colin, S. P., Costello, J. H., Hansson, L. J., Titelman, J. and Dabiri, J. O. (2010) Stealth predation and the predatory success of the invasive ctenophore *Mnemiopsis leidyi*. *Proc. Natl. Acad. Sci. USA*, **107**, 17223-17227.
- Costello, J. H., Bayha, K. M., Mianzan, H. W., Shiganova, T. A. and Purcell, J. E. (2012) Transitions of *Mnemiopsis leidyi* (Ctenophora: Lobata) from a native to an exotic species: a review. *Hydrobiologia*, **690**, 21-46.
- Graham, E. S., Tuzzolino, D. M., Burrell, R. B. and Breitburg, D. L. (2009) Interannual variation in gelatinous zooplankton and their prey in the Rhode River, Maryland. *Smiths. Contrib. Mar. Sci.*, **38**, 369-377.
- Jaspers, C. (2012) *Ecology of gelatious plankton: With emphasis on feeding interactions, distribution pattern and reproduction biology of Mnemiopsis leidyi in the Baltic Sea*. PhD Thesis. Technical University of Denmark, Charlottenlund, pp. 1-130.
- Jaspers, C., Haraldsson, M., Lombard, F., Bolte, S. and Kiørboe, T. (2013) Seasonal dynamics of early life stages of invasive and native ctenophores give clues to invasion and bloom potential in the Baltic Sea. *J. Plankton Res.*, **35**, 582-594.
- Javidpour, J., Molinero, J. C. and Sommer, U. (2009) Reconsidering evidence for *Mnemiopsis* invasion in European waters: reply. *J. Plankton Res.*, **32**, 97-98.
- Kasuya, T., Ishimaru, T. and Murano, M. (2008) Reproductive characteristics of the lobate ctenophore *Bolinopsis mikado* (Moser). *Plankton Benthos Res.*, **3**, 72-77.
- Kremer, P. (1976a) *The ecology of the ctenophore Mnemiopsis leidyi in Narragansett Bay*. PhD Thesis. University of Rhode Island, pp. 1-311.
- Kremer, P. (1976b) Population dynamics and ecological energetics of a pulsed zooplankton predator, the ctenophore *Mnemiopsis leidyi*. In M. Wiley (ed) *Estuarine Processes*. Vol. 1. Academic Press, New York, USA, pp. 197-215.
- Kremer, P. (1982) Effect of food availability on the metabolism of the ctenophore *Mnemiopsis mccradyi*. *Mar. Biol.*, **71**, 149-156.
- Kremer, P., Reeve, M. R. and Syms, M. A. (1986) The nutritional ecology of the ctenophore *Bolinopsis vitrea*: comparisons with *Mnemiopsis mccradyi* from the same region. *J. Plankton Res.*, **8**, 1197-1208.
- Martinussen, M. B. and Båmstedt, U. (1995) Diet, estimated daily food ration and predator impact by the scyphozoan jellyfishes *Aurelia aurita* and *Cyanea capillata*. In H. R. Skjoldal, C. Hopkins, K. E. Erikstad and H. P. Leinaas (eds), *Ecology of Fjords and Coastal Waters*. Vol. 1. Elsevier Science Publishers, Amsterdam, pp. 127-145.

- Reeve, M. R., Syms, M. A. and Kremer, P. (1989) Growth dynamics of a ctenophore (*Mnemiopsis*) in relation to variable food supply , I . Carbon biomass, feeding, egg production, growth and assimilation efficiency. *J. Plankton Res.*, **11**, 535-552.
- Reeve, M. R., Walter, M. A. and Ikeda, T. (1978) Laboratory studies of ingestion and food utilization in lobate and tentaculate ctenophores. *Limnol. Oceanogr.*, **23**, 740-751.
- Riisgård, H. U., Jaspers, C., Serre, S. and Lundgreen, K. (2012) Occurrence, inter-annual variability and zooplankton-predation impact of the invasive ctenophore *Mnemiopsis leidyi* and the native jellyfish *Aurelia aurita* in Limfjorden (Denmark) in 2010 and 2011. *BioInvasions Rec.*, **1**, 145-159.
- Sullivan, L. J. and Gifford, D. J. (2004) Diet of the larval ctenophore *Mnemiopsis leidyi* A. Agassiz (Ctenophora, Lobata). *J. Plankton Res.*, **26**, 417-431.
- Zaika, V. E. and Revkov, N. K. (1994) Anatomy of gonads and regime of spawning of ctenophore *Mnemiopsis* sp. in the Black Sea. *Zool. J.*, **73**, 5-10 (in Russian).

TABLE AND FIGURE LEGENDS

Table I: *Mnemiopsis leidyi* (n=5) reproduction rates in northern Europe (Aug/Sept 2010 during two sampling events) at *in situ* salinity (22.5) and temperature (16.5°C), with highest egg production (11,232 eggs d⁻¹, 78 mm total length) recorded in northern Europe so far. Carbon specific egg production (SEP) is presented based on given length to carbon conversions (see text) and egg carbon of 0.22 µgC egg⁻¹.

Fig. 1: Carbon and nitrogen content of *Mnemiopsis leidyi* eggs. The average is $0.22 \pm 0.016 \mu\text{gC egg}^{-1}$ and $0.07 \pm 0.005 \mu\text{gN egg}^{-1}$, respectively.

Fig. 2: Pictures of *Mnemiopsis leidyi* eggs ($503 \pm 58 \mu\text{m}$) of different, early cleavage stages including one late developmental stage (picture, right) with a fully developed larvae (300 µm) inside the egg shell. Egg sizes have a similar size range in native and invaded, northern European populations ranging between 440 to 625 µm and 480 to 630 µm diameter, respectively.

Table I:

	Maximum	Minimum	Average	SD
Total length (mm)	78	70	73.8	3.5
Oral-aboral length (mm)	57	51	54.3	2.5
Eggs ($\text{ind}^{-1} \text{ d}^{-1}$)	11,232	6,432	8,432	1,884
SEP	10.4	7	8.4%	1.4%

Fig. 1




Fig. 2

