

RESEARCH ARTICLE

10.1002/2016JF003975

Key Points:

- Shorelines fractality is an indicator of salt marsh vulnerability
- Salt marsh erosion rates maintain similar trend in time
- Salt marsh erosion depends on wave energy

Supporting Information:

- Supporting Information S1

Correspondence to:

N. Leonardi,
nicleona@liverpool.ac.uk

Citation:

Leonardi, N., Z. Defne, N. K. Ganju, and S. Fagherazzi (2016), Salt marsh erosion rates and boundary features in a shallow Bay, *J. Geophys. Res. Earth Surf.*, 121, 1861–1875, doi:10.1002/2016JF003975.

Received 30 MAY 2016

Accepted 10 SEP 2016

Accepted article online 13 SEP 2016

Published online 22 OCT 2016

Salt marsh erosion rates and boundary features in a shallow Bay

Nicoletta Leonardi^{1,2}, Zafer Defne³, Neil K. Ganju³, and Sergio Fagherazzi²

¹Department of Geography and Planning, University of Liverpool, Liverpool, UK, ²Department of Earth and Environment, Boston University, Boston, Massachusetts, USA, ³United States Geological Survey, Woods Hole, Massachusetts, USA

Abstract Herein, we investigate the relationship between wind waves, salt marsh erosion rates, and the planar shape of marsh boundaries by using aerial images and the numerical model Coupled-Ocean-Atmosphere-Wave-Sediment-Transport Modeling System (COAWST). Using Barnegat Bay, New Jersey, as a test site, we found that salt marsh erosion rates maintain a similar trend in time. We also found a significant relationship between salt marsh erosion rates and the shape of marsh boundaries which could be used as a geomorphic indicator of the degradation level of the marsh. Slowly eroding salt marshes are irregularly shaped with fractal dimension higher than rapidly deteriorating marshes. Moreover, for low-wave energy conditions, there is a high probability of isolated and significantly larger than average failures of marsh portions causing a long-tailed distribution of localized erosion rates. Finally, we confirm the existence of a significant relationship between salt marsh erosion rate and wind waves exposure. Results suggest that variations in time in the morphology of salt marsh boundaries could be used to infer changes in frequency and magnitude of external agents.

1. Introduction

Located at the interface between marine and terrestrial environments, salt marshes are ecosystem-based flood defenses that help reduce the impact of storms and hurricanes on coastal communities [e.g., Temmerman *et al.*, 2013; Möller *et al.*, 1999; Möller, 2006; Zhao and Chen, 2013]. In recent years, these vegetated surfaces have been at the center of many restoration projects based on the concept of “living shorelines” [e.g., Temmerman *et al.*, 2013; Fagherazzi, 2014]. The value of their storm protection services has been estimated up to 5 million USD per km² in the United States [e.g., Costanza *et al.*, 2008], and 786 million GBP per year for the UK marshes [UK National Ecosystem assessment, 2011; Foster *et al.*, 2013; Moller *et al.*, 2014]. Salt marshes also provide other important ecosystem services such as nutrient removal, habitat provision, and carbon sequestration over decennial to millennial time scales [e.g., Plater *et al.*, 1999; Zedler and Kercher, 2005; Mudd *et al.*, 2009]. Large salt marsh losses have been documented worldwide; as an example, in England and Wales salt marsh areal loss has been estimated to be around 83 ha yr⁻¹ by the UK Environmental Agency [Foster *et al.*, 2013; Environment Agency, 2011], 105 ha yr⁻¹ for the period in between 1993 and 2013 [Pye and French, 1993], and 349 ha yr⁻¹ for the period between 1998 and 2048 [Lee, 2001]. In the Solent (UK) 40% of the total area present in 1971 was eroded between 1971 and 2001 [Cope *et al.*, 2008], while for the Greater Thames the erosion was estimated to be around 25% of the total area present in 1973 [Cooper *et al.*, 2009]. For areas in the south west of the Netherlands and the Wadden Sea, cliff erosion up to 4 m/yr has been recorded, despite positive vertical accretion rates [e.g., Bakker *et al.*, 1993].

For the southern part of the Venice Lagoon and for an area of 50 × 50 m, marsh edge retreat has been estimated around 1.2–2.2 m/yr for the period from 1993 to 1995. Highly dynamic patterns in marsh change have been identified, with accretion rates varying from 1.1 to 2.3 cm/yr [Day *et al.*, 1998]. Highest accretion rates corresponded to the occurrence of strong storms as a consequence of sediment resuspension from nearby tidal flats. Day *et al.* [1998] suggested that this area was rapidly deteriorating due to lateral migration exceeding vertical accretion.

Erosion rates with similar order of magnitude and up to 80 cm/yr have been recently measured in the northern part of the Venice Lagoon [e.g., Bendoni *et al.*, 2016], where it has been also shown that failures of large marsh blocks do not necessarily correlate with instantaneous wave forcing, and that such large failures can also occur during calm periods. For the East Coast of the United States, long-term field measurements of salt marsh erosion in Plum Island Sound and Virginia Coast Reserve [Leonardi and Fagherazzi, 2014, 2015] also demonstrated

that failures of large marsh blocks can happen independently from high-wave energy conditions. High-resolution field data also suggest that salt marsh erosion is more spatially variable in case of low-wave energy conditions, rather than under high wave energy, and that under low-wave energy conditions erosion events follow a long-tailed frequency-magnitude distribution [Leonardi and Fagherazzi, 2014, 2015].

Salt marsh resistance to external agents is linked to sediment dynamics as well. For instance, measurements in the Wadden Sea [Schuerch *et al.*, 2014], as well as numerical and field experiments in the Scheldt estuary [Temmerman *et al.*, 2003] indicate that salt marsh erosion or accretion depends on the exchange of sediments between salt marshes, tidal flats, and tidal channels. Vegetated surface also influence salt marsh erosion by indirectly modifying soil parameters [e.g., Feagin *et al.*, 2009].

In terms of vertical dynamics, salt marsh accretion rates have been found to be spatially and temporally variable [e.g., Day *et al.*, 1998; Temmerman *et al.*, 2005; Pethick, 1993]. Variability in accretion rates has been identified as largely depending on the proximity of tidal channels, erosion rates of nearby areas, as well as tidal flat characteristics [Day *et al.*, 1998; Temmerman *et al.*, 2005]. As an example, salt marshes in the southeast England have been found to more rapidly accrete right behind the erosional front [e.g., Day *et al.*, 1998]. In spite of accretion rates being spatially variable, numerous studies confirm that feedbacks between sediment deposition, organic matter accretion, and biomass production generally allow salt marshes to be stable on the vertical direction and keep pace with sea level rise [e.g., Cundy and Croudace, 1996; van der Wal and Pye, 2004; Temmerman *et al.*, 2003, 2005; Temmerman and Kirwan, 2015; Kirwan *et al.*, 2016; Priestas *et al.*, 2015].

While resistant to sea level rise, salt marshes are inherently weak and vulnerable to lateral erosion induced by wave action because wind waves impact bare sediments below the vegetated surface where material is easily mobilized [e.g., Schwimmer, 2001; van de Koppel *et al.*, 2005; Marani *et al.*, 2011; McLoughlin *et al.*, 2015; Fagherazzi, 2013; Leonardi *et al.*, 2016].

Because wind waves are one of the main agents contributing to marsh lateral erosion, understanding their action is critical for restoration projects aimed at the maintenance and improvement of wetland ecosystem services.

Salt marsh erosion by wind waves is a complex phenomenon that includes several mechanisms such as the continuous removal of small particle aggregates from the marsh platform and marsh edges or the sudden and discontinuous detachment of marsh portions having size comparable to the bank height [e.g., Bendoni *et al.*, 2014]. For example, waves can undercut the root mat with the formation of a marsh overhang that eventually breaks off and topple [e.g., Schwimmer, 2001; Allen, 1989]. For salt marshes, downward shearing, bending followed by tensile fractures, and subhorizontal tensile fractures are common failures mechanisms as well [e.g., Allen, 1989].

Herein, we will use the term “erosion event” to refer to the erosion rate (m/yr) along relatively short marsh portions (i.e., stretches of shoreline of the order of 3 m) and that might be caused by continuous particle removal, mass failures, or by a combination of both.

A direct relationship between wave energy and salt marsh lateral retreat has been recognized in prior studies [e.g., Leonardi *et al.*, 2016; Schwimmer, 2001; McLoughlin *et al.*, 2015]. However, many existing studies only focus on small spatial scales (shoreline lengths of the order of hundreds of meters), and less attention has been paid to the uniformity of salt marsh erosion along stretches of shorelines at a bay or estuary scale, as well as to the possible influence of different wind wave exposures and erosion rates on the shape of marsh boundaries.

Studies of erosion rate uniformity along marsh boundaries are relevant to the risk assessment of coastal wetlands, as they provide information about the occurrence of erosion events far from the average. In fact, the occurrence of unexpected erosion episodes raises questions about the effectiveness of risk assessment plans and coastal protection schemes. There is, thus, a need to further understand the occurrence of infrequent episodes, and whether they might be different from average, long-term erosion trends [Malamud, 2004; Bernardara *et al.*, 2008; Leonardi and Fagherazzi, 2014, 2015]. The occurrence of isolated large erosion events could lead to inaccurate evaluations of the erosion trends in the area if detailed data sets for the morphological behavior of salt marshes or detailed surveys of the region are not present. It is thus appropriate to distinguish erosion rates averaged over large spatial scales with respect to localized erosion rates.

Many studies use wave power as a proxy for wind wave exposure [e.g., *Schwimmer, 2001; Leonardi et al., 2016*]. While this approach has been successful, wave thrust (integral along the vertical of the dynamic pressure) could be another useful indicator for the action of wind waves on wetland boundaries because it represents the actual force contributing to the dynamic equilibrium of marsh shorelines [e.g., *Tonelli et al., 2010; Bendoni et al., 2014; Francalanci et al., 2013*] and allows taking into account the presence of different water levels and marsh boundary features such as platform elevation with respect to mean sea level, and orientation of the boundary with respect to wave direction.

Water levels have been found to influence the action of wind waves [e.g., *Tonelli et al., 2010; Möller et al., 1999; Moller et al., 2014; Möller, 2006*]; laboratory, numerical, and field experiments [e.g., *Tonelli et al., 2010; Bendoni et al., 2014; Francalanci et al., 2013*] confirm that the erosive action of wind waves at the edge of salt marshes is more severe when water levels are slightly below the marsh platform. Specifically, wave action has been found to increase with water depth, up to the point when the marsh is submerged, and then it rapidly decreases. When the wave peak is lower than the marsh platform, waves are partially reflected, interfering with incident waves. Once the bank is submerged, wind waves are affected by the water depth on the marsh platform. In these conditions waves rapidly break, and a large amount of energy is dissipated due to the presence of vegetation [e.g., *Le Hir et al., 2000; Möller et al., 1999; Möller, 2006; Temmerman et al., 2005*]. For very high water levels skimming flow may take place across a completely submerged canopy reducing wave energy dissipation. However, at these high water depths the wave thrust on the marsh boundary is minimal, leading to limited marsh erosion [*Tonelli et al., 2010*]. Wave thrust calculation is also fundamental to physically based models aimed at describing specific failure mechanisms such as block toppling. For the latter, marsh blocks are schematized as rigid and subject to hydrodynamic forces, gravity, and soil cohesion [e.g., *Bendon et al., 2014, 2016; Francalanci et al., 2013*]. It is therefore appropriate to test the suitability of more physically based parameters to describe marsh erosion, as the base to gradually incorporate specific failure mechanisms into numerical studies.

In this manuscript we focus on erosion processes and wave action at the boundary of salt marshes in the Barnegat Bay-Little Egg Harbor system using aerial images and a numerical model. We focus on the spatial distribution of lateral erosion rates and on the relationship between the shape of marsh boundaries and erosion rates. Fractal dimension is used as an indicator for the shape of marsh boundaries. We explore whether marsh boundary geometry can be used to infer whether salt marshes are slowly or rapidly degrading. To this end, an erosion analysis has been carried out for extensive stretches of shoreline (order of hundreds of kilometers). We further explore the long-term relationship between erosion rates and wave thrust at the marsh boundary. The analysis of marsh boundary geometry builds on previous work conducted on other salt marshes in the United States Atlantic Coast, and according to which a relationship exists between marsh boundary roughness and wave energy [*Leonardi and Fagherazzi, 2014, 2015*]. The second part of the work builds on numerous studies according to which there is a strict relationship between salt marsh erosion rates and wave energy [*Schwimmer, 2001; Marani et al., 2011; Leonardi et al., 2016*].

2. Study Site

The Barnegat Bay-Little Egg Harbor system is located along the central New Jersey coastline in Ocean County on the eastern seaboard of the USA (Figure 1) and consists of three shallow coastal bays: Barnegat Bay, Manahawkin Bay, and Little Egg Harbor. The system is composed of a shallow, lagoon type estuary, separated from the Atlantic Ocean by a narrow barrier island complex approximately 70 km long.

The width of the lagoons ranges from 2 to 6 km, and the water depths range from 1 to 5 m, with an average value of 1.5 m. Most of the western side of Barnegat Bay is deeper (~1–3 m depth) than the eastern side (~1 m depth). The deepest area runs along the Intracoastal Waterway and spans the bay length [*Kennish, 2001; Defne and Ganju, 2014; Aretxabaleta et al., 2014*]. The barrier is breached at Barnegat Inlet and Little Egg Inlet. Barnegat Inlet has a mean width of 400 m, and a maximum water depth of 15 m. Little Egg Inlet has a width of approximately 2 km, and average water depth of 10 m. Figure 1 shows the bathymetry of the area and long-term statistics of hourly mean wind speed and direction (period from 1991 to 2013).

During the fall and winter seasons winds predominantly come from northwest to west directions; during spring and summer the jet stream retreats northward and winds are from the south to the southwest; in summer subtropical high pressures generally produce warm and humid southerly breezes and average wind




Figure 1. (a) Bathymetry of the Barnegat Bay-Little Egg Harbor system. Yellow lines indicate the boundaries of the computational domain. (b) Wind rose for the area for the period 1991–2013 (wind station, station ID LLNR 830, 40°15'3"N, 73°9'52"W).

speed are less than 15 km/h. The area is also characterized by northeasterly winds coming from coastal low-pressure storms (Nor'easters) which move toward the north along the coast. These extratropical storms occur from September to March. In summer, thunderstorms are frequent. Late summer and fall can be characterized by the occurrence of hurricanes and tropical storms. Generally, the frequency with which major hurricanes impact the area is lower with respect to more southerly Atlantic states. For Barnegat Bay and nearby areas, on average, one tropical cyclone passes within 100 km from the coast every 4.5 years [Kennish, 2001; Elsner and Kara, 1999]. As an example, between 1935 and 1967 nine hurricanes occurred within 160 km from the Bay [Kennish, 2001]. Extratropical coastal storms (Nor'easters) are more frequent in the area. Because Nor'easters winds have usually wind velocity lower than hurricanes, their damage is generally restricted to the coast, while hurricanes damage can extend inland significantly. On the other hand, since Nor'easters are large, their damage can cover extensive coastal areas [Elsner and Kara, 1999].

Wind measurements were used to reconstruct the wave climate in the area. In enclosed basins like the one studied herein where waves are locally generated, it is realistic to assume that waves instantaneously adjust their direction to the wind direction [Mariotti *et al.*, 2010]. Tides are semidiurnal, the tidal range outside the Bay is over 1 m, but it attenuates within the estuary to less than 20 cm in some locations [Aretxabaleta *et al.*, 2014]. In the past, the lagoon was almost completely fringed by salt marshes. In 1888, the salt marsh area was approximately 14,850 ha, while in 1995 the marsh extent was estimated to be 9940 ha; this decline has been attributed to both natural and human disturbances, such as land use change and dredging operations [Lathrop and Bogner, 2001]. Lower marsh areas are covered with the short form of *Spartina alterniflora*, while the high marsh is covered with *Spartina patens* [Suk *et al.*, 1999]. Vegetation characteristics can be important when dealing with erosion, as vegetation can trap sediments through leaves and promote sedimentation rates, slow the flow, increase soil resistance through the root mat, promote certain type of marsh failures (e.g., root mat toppling) with respect to other failure mechanisms, and influence soil properties [e.g., Feagin *et al.*, 2009; Möller, 2006]. Among others, species zonation, vegetation density, stem height, and Young's




Figure 2. (a) Example of digitized shoreline (detail at latitude 39.64°, longitude -74.22°). Fractions of total digitized shoreline subject to different erosion rates for the periods (b) 1930–2007 and (c) 2007–2013.

modulus can significantly influence frictional resistance [e.g., *Allen and Pye, 1992*]. To characterize vegetation friction, many studies have focused on the overall roughness coefficient of vegetation by using a variety of laboratory, field, and numerical methods [e.g., *Tsujiimoto et al., 1996; Järvelä, 2002*]. Other studies have been focusing on the effect of vegetation on flow resistance with artificial plants or by schematizing stems as rigid cylinders. These studies have used a variety of laboratory and numerical techniques [e.g., *Nepf, 1999*]. Real plants have been also used to directly investigate frictional resistance trough field and laboratory experiments [e.g., *Järvelä, 2002; Armanini et al., 2005; Allen and Pye, 1992*].

3. Methods

3.1. Aerial Photography

To determine the geometry of salt marsh boundaries and related erosion rates at a large spatial scale, we used a suite of aerial images (1930, 1995, 2002, 2007, 2010, and 2013). For the 1930, 2007, and 2013 aerial photographs, we digitized more than 100 km of marsh shoreline, corresponding to the majority of the interior marsh boundary. Using the digitized aerial images, we then computed salt marsh erosion from 1930 to 2013 and from 2007 to 2013 (Figures 2 and 3). The three aerial images were selected among the available ones based on their quality, and with the aim to investigate long-term erosion rates (1930–2013), as well as more recent and shorter-term rates of change (2007–2013). Specifically, images for 1930 and 2013 were the farthest available in time, and the period from 2007 to 2013 was the shortest interval at which changes in marsh erosion profile were still detectable for the majority of the shoreline.

For the 1995, 2002, and 2010 images we digitized a series of 10 small marsh shoreline portions of the order of 1 km to retrieve information about erosion at a higher temporal resolution (Figure 3a). The 1930 image is a black and white aerial photograph produced by scanning 261 mosaic tile prints at 400 dpi, with an average pixel size of 1.98 m, and 24.3 m RMS. The 2002 and 2007 images are respectively infrared and natural color orthophotographs produced at a scale of 1:2400 with a 0.3 m pixel resolution. The average horizontal accuracy is ±1.2 m with a 95% confidence level. The 1995, 2010, and 2013 natural color orthophotographs were obtained by using 1 m ground sample distance source images and were rectified with an average horizontal accuracy of ±5 m with a 95% confidence level. Images for 1995, 2002, 2007, 2010, and 2013 were obtained in their orthorectified form and did not require particular postprocessing. The 1930 photograph was rectified using georeferencing tools within ArcGIS®, and by considering as ground control points within different tiles, corner of buildings and road intersections. The mean root-mean-square error of the 1930 aerial photograph rectifications was ±5 m. The wetland boundary was digitized by contouring the shoreline through ArcGIS® points features placed at an average distance of 2.5 m. The total position uncertainty, U_t , was estimated following established methodologies [e.g., *Cowart et al., 2011*], and by accounting for both rectification (E_r) and digitalization (E_d) errors.

$$U_t = \pm \sqrt{\sum E_r^2 + \sum E_d^2} \tag{1}$$

where the sum refers to all used images. A digitalization error, E_d , of 0.5 m was calculated by digitalizing same stretches of shorelines multiple times and finding the maximum distance between multiple digitalizations.




Figure 3. (a) Spatial distribution of average erosion rates from 1930 to 2007. Numbers indicate portions of the shoreline that were also digitized for the years 1995, 2002, and 2010. White arrows, as well as numbering order follow the alongshore coordinate plotted on the horizontal axis of Figure 3b. (b) Erosion rates as a function of the alongshore coordinate. Red and blue lines correspond to erosion rates for the periods 1930–2007 and 2007–2013 respectively. Yellow and cyan lines are moving averages of the same variables over a 1 km distance. The alongshore coordinate follows numbering in Figure 3a. Vertical dashed lines separate portion of the alongshore coordinate covered by different arrows.

Maximum position uncertainties (U_t) are 24.8m for the comparison of 1930–2007 and 25.33 m for the 1930–2013 image comparison, which correspond to a maximum uncertainty of 0.32 m/yr, and 0.30 m/yr, respectively. Given two tracked shorelines, at every point along the first boundary, erosion was estimated as the minimum distance of that point from points belonging to the shoreline of the other year.

To compute shoreline fractal dimension, we followed the Minkowski-Bouligand method [e.g., Dubuc *et al.*, 1989]. Fractal dimension allows evaluating the degree of marsh boundary complexity by computing how fast length measurements change with the scale at which they are measured. Generally speaking, given a topological set S of dimension n , for any small value $\epsilon > 0$, let $N_\epsilon(S)$ be the minimum number of n dimensional cubes of side-length ϵ needed to cover S . The fractal dimension, D , is such that

$$N_\epsilon(S) \sim 1/\epsilon^D \text{ as } \epsilon \rightarrow 0 \tag{2}$$

and

$$D = -\lim_{\epsilon \rightarrow 0} \frac{\log N_\epsilon(S)}{\log(\epsilon)} \tag{3}$$

When approaching the limit, if the surface is smooth and differentiable, its fractal dimension, D , equals the topological dimension, d . For a rough and nondifferentiable curve the fractal dimension may exceed the topological dimension d .

We are also interested in evaluating the frequency-magnitude distribution of erosion events. In particular, it is useful to have information about the possible occurrence of unexpected and above average erosion rates that belong to the tail of the erosion events distribution. To gain more information about these tail events, we approximate the frequency-magnitude distribution of erosion events by means of a logarithmic frequency-magnitude distribution. The main advantages of using a distribution that is normal in the logarithm of a parameter are as follows: (i) the distribution is positive-definite, and it is thus suitable to represent

quantities that cannot be negative, such as erosion rate values; and (ii) for many natural processes, where the possibility of events falling far from the average exists, a logarithmic distribution often provides a better fit with respect to Gaussian-like functions [e.g., Limpert *et al.*, 2001].

For a lognormal distribution, the probability density function, f_x , is

$$f_x(x, \mu, \sigma) = \frac{1}{x\sigma\sqrt{2\pi}} e^{-\frac{(\ln x - \mu)^2}{2\sigma^2}} \quad (4)$$

where μ is the mean and σ is the variance. With increasing variance, the tail of the distribution of erosion events gets longer, distribution skewness increases, and the possibility of erosion events far from the mean increases as well.

3.2. Numerical Model

In this study we used the Coupled-Ocean-Atmosphere-Wave-Sediment-Transport (COAWST) Modeling System [Warner *et al.*, 2008, 2010]. In COAWST, the ocean model Regional Oceanic Modeling System (ROMS), atmospheric model WRF, wave model Simulating Waves Nearshore (SWAN), and modules of the Community Sediment Transport Model are fully coupled by means of the Model Coupling Toolkit. The ocean model ROMS is a three-dimensional, free-surface, terrain-following model solving finite difference approximations of the Reynolds Averaged Navier-Stokes equations, using hydrostatic and Boussinesq assumptions [Chassignet *et al.*, 2000; Shchepetkin and McWilliams, 2005; Haidvogel *et al.*, 2008; Kumar *et al.*, 2011]. To include the effect of surface waves, the momentum equations require, as input, information for wave height, wave energy dissipation, propagation direction, and wavelength. These information are obtained from the numerical model SWAN (Simulating Waves Nearshore), which solves the transport equations for waves action density, and accounts for shoaling, refraction, wind waves generation, wave breaking, bottom dissipation, and nonlinear wave interactions [Booij *et al.*, 1999; Ris *et al.*, 1999].

The boundaries of the model domain are presented in Figure 1a (yellow line) and include Barnegat Bay, Little Egg Harbor, part of Great Bay to the South, and Manasquan Inlet to the north. The computational grid consists of 160 east-west and 800 north-south grid points with seven vertical layers having equal depth. Cell sizes vary from 40 to 200 m, with the grid being refined at the inlets and around complex morphological features. The bathymetry is based on the National Ocean Service Hydrographic Survey data [NOAA National Ocean Survey, 2012; Defne and Ganju, 2014]. At the seaward boundaries, we prescribed water level variations typical of a tidal cycle based on the ADCIRC tidal constituents database for the North Atlantic Ocean and used a combination of Flather and Chapman boundary conditions. The Flather [1976] boundary condition allows radiating out deviations of the normal component of barotropic velocities at the speed of external gravity waves, and the Chapman boundary condition is the corresponding condition for surface elevations [Chapman, 1985]. At the westward boundary we prescribed a radiation boundary condition following Orlandi [1976] radiation scheme and allowing tidal energy to propagate landward. The model was calibrated by changing the bottom roughness coefficient to attain the best agreement with measurements from seven water level stations and three tidal discharge stations of the U.S. Geological Survey within the Barnegat Bay-Little Egg Harbor estuary and for a period comprising the first 2 weeks of March 2012. The Brier-Skill-Score [e.g., Murphy and Epstein, 1989] was used to assess model calibration. Skill assessments of the model ranged from very good to excellent [Defne and Ganju, 2014]. More specific information about the model as well as model calibration and setup can be found in Defne and Ganju [2014]. The model was run with constant wind speed and direction using 3 classes of wind (5 m/s, 10 m/s, and 25 m/s), 8 directions spanning from 0 to 360°, and 4 different storm surge values (± 0.3 m and $+ - 1$ m), for a total of 96 simulations. Results obtained from these simulations were combined with wind speed, wind direction, and water level data retrieved from a nearby NOAA station (station ID LLNR 830, 40°15'3"N, 73°9'52"W) to get a proxy for the long-term wave climate along the shoreline. We obtained the model outputs indicated above by using the phase-averaged model SWAN to calculate wave parameters along marsh boundaries and then followed results from a phase resolving model [Tonelli *et al.*, 2010] to evaluate the reduction in wave thrust occurring when waves overtop the marsh scarp. The model proposed by Tonelli *et al.* [2010] combines nonhydrostatic Boussinesq equations, nondispersive nonlinear shallow water equations, and has been widely validated. Results presented by Tonelli *et al.* [2010] and used in this manuscript focus on areas of the marsh platform close to the marsh edge (1–3 m) and incorporate the presence of vegetation through the use of friction factors typically derived for salt

marshes (0.07–0.47) [Department of the Army, Waterways Experiment Station, Corps of Engineers, and Coastal Engineering Research Center, 1984]. Wave thrust is calculated as the integral along the vertical axis of the dynamic pressure of waves. The dynamic pressure of wind waves can be calculated as

$$p_d(z, t) = \rho g K_p(z) \eta(t) \quad (5)$$

where $K_p(z)$ is the pressure-response factor accounting for the dynamic component due to water particle acceleration and can be calculated as

$$K_p(z) = \frac{\cosh k(h+z)}{\cos(kh)} \quad (6)$$

To take into account the dependence of wave thrust on water depth, we followed results from Tonelli *et al.* [2010] and prescribed an up to 60% linear reduction in wave thrust proportional to the water depth on the marsh platform up to 0.4 m above the surface, after which wave thrust is kept constant. As part of this project we also developed a subroutine for wave thrust calculation, which is now implemented in the COAWST modeling suite (see supporting information).

This is a simplified approach and presents several limitations particularly in regard to the dissipative effect of vegetation which might strongly change depending on both vegetation and wave characteristics. Vegetation properties can be especially important: trough wave flume experiments it has been shown that over a distance of 40 m, up to 60% of dissipation is attributable to the vegetation cover even under extreme storm conditions [Moller *et al.*, 2014; Möller, 2006]. Similar conclusions have been drawn from field measurements, according to which wave energy dissipation up to 63% was found to occur over the first 100 m [Möller *et al.*, 1999; Brampton, 1992]. Variable dissipation rates have been measured in the field [e.g., Koch, 1996; Prager and Halley, 1999] and an exponential energy decay with respect to the distance of the shoreline has been generally accepted with exponents of the order of 0.0187 [e.g., Brampton, 1992; Möller *et al.*, 1999]. This simplified approach is herein accepted in part for the lack of direct field measurements of waves over vegetated surfaces, and in part because we are focusing on the action of waves at the edges of the marsh boundary. Over such short distances (1–3 m) the action of vegetation is not fully developed even if an exponential formulation for wave energy dissipation over distance is assumed [Tonelli *et al.*, 2010; Möller *et al.*, 1999].

4. Results

4.1. Boundary Erosion Analysis

Figures 2b and 2c report fractions of the entire shoreline subject to different erosion rates and present an overview of the long term (1930–2007) erosion rate, with respect to more recent and short-term erosion rate values (2007–2013). Percentages refer to the entire digitized shoreline. For both time intervals, more than half of the digitized shoreline is eroding at rates ranging from 0.25 to 2 m/yr. Around 25% of the shoreline is eroding slower than 0.25 m/yr, or is not eroding at all. For both periods erosion rates are similar and reach a maximum near Little Egg Harbor. Despite small percentage differences among different erosion categories, no substantial variations can be noticed when comparing average long-term and short-term erosion rates at a large spatial scale.

Spatial variations in erosion rate for the period 1930–2007 are presented in Figure 3. The highest erosion rates occur in proximity of Little Egg Inlet, around points of greatest fetch in Little Egg Harbor, and in the western part of Barnegat Bay. Figure 3b shows variation in erosion rate with respect to the alongshore distance from the shoreline. The horizontal coordinate follows the white arrows of Figure 3a. Blue and red lines are long-term (1930–2007) and short-term (2007–2013) erosion rates, respectively. Yellow and cyan lines are corresponding averages over moving distances of 1 km. Despite the measurement noise, the erosion rates for the two periods maintain a similar trend for most of the marsh boundary, in terms of both magnitude and spatial distribution (Figure 4a). Strongest differences occur for points around Little Egg Inlet, where the erosion rate is one of the highest within the domain but has lowered in the period 2007–2013 with respect to the period 1930–2007. To better highlight possible temporal variations, erosion rates for shorter intervals of time are presented in Figure 4b for 10 different stretches of shoreline having an average length of 1 km. Apart from




Figure 4. (a) Comparison between average erosion rates from 2007 to 2013 and average erosion rates from 1930 to 2007 ($R^2 = 0.76$). (b) Average erosion rate in different time intervals for 10 stretches of shoreline having length of the order of 1 km, and whose locations are indicated in Figure 3a. Note that the period 2010–2013 includes hurricane Irene and hurricane Sandy.

locations in proximity of Little Egg Inlet, erosion rates remain similar in time and mainly oscillate around a constant value.

Fractal dimension was used to estimate shoreline roughness, and the existence of a potential relationship between erosion rate and morphology of salt marsh boundaries. Figures 5a and 5c show the relationship between the fractal dimension of the marsh boundary and erosion rates for the period from 1930 to 2007, and from 2007 to 2013, respectively. Fractal dimension is significantly ($p < 0.05$) and negatively correlated to erosion rate, suggesting that rapidly eroding areas have smoother marsh boundaries with respect to slowly eroding areas.

To gain further information about salt marsh erosion processes, it is useful to investigate the frequency-magnitude distribution of erosion events. The likelihood of an erosion event to deviate from average erosion




Figure 5. Relationship between erosion rate (horizontal axis) and fractal dimension (vertical axis) for the periods (a) 1930–2007 and (c) 2007–2013. Solid black circles indicate values obtained by averaging data over regular bins to emphasize the overall trend. Relationship between erosion rate (horizontal axis) and the log-normal standard deviation of the erosion events (vertical axis) for the periods (b) 1930–2007 and (d) 2007–2013. Solid black circles indicate values obtained by averaging data over regular bins to emphasize the overall trend; Figures 5c–5e are examples of frequency-magnitude distributions of erosion events for different stretches of shoreline, and corresponding shoreline images. For shoreline locations, see corresponding numbering in Figure 2c.




Figure 6. Map of average wave thrust values for the entire bay from 1991 to 2013.

rates was estimated by approximating the frequency-magnitude distribution of erosion events using a log-normal distribution, and by computing corresponding standard deviation values. The lognormal standard deviation is negatively correlated to average erosion rates (Figures 5b and 5d). This result indicates that unexpected erosion rates far from the mean occur more often in slowly eroding areas. On the contrary, low standard deviation values, corresponding to high average erosion rates, indicate that in rapidly degrading marshes the majority of erosion episodes fall around the mean. Figures 5e–5g provide examples of frequency-magnitude distributions of erosion events and marsh boundary profiles at three different sites. Site locations are indicated in Figure 3 (numbering in Figures 5c–5e corresponds to numbering in Figure 3a). It is possible to notice that a long tail of the frequency-magnitude distribution of erosion events corresponds to a slowly eroding marsh with a rough boundary.

4.2. Wind Wave Exposure Analysis

Figure 6 represents long-term (1991–2013) average wave thrust values calculated at marsh boundaries for the entire bay. Wave thrust values in Figure 6 correspond to values averaged throughout a tidal cycle, and then further averaged from 1991 to 2013. Herein results refer to the period from 1991 to 2013 because no wind data before 1991 are available from buoy stations. Wave thrust values along the boundary were obtained by associating at every point of the boundary and for every wind speed, direction, and water level, wave thrust values obtained from the numerical tests carried out to assess general system susceptibility to different winds and water levels. If a reduction in wave thrust for a submerged marsh is taken into account (Figure 6b), average wave thrust values for the period of interest strongly decline but maintain a similar spatial distribution. Some differences in the spatial distribution of the two different wave thrust calculations are present at the northeastern corner of Great Bay, and at the west side of Barnegat Bay. Generally speaking, the eastern part of Barnegat Bay, the southeast side of Great Bay, and points of greatest fetch in Little Egg Harbor appear to be the most exposed to the action of wind waves.

Figure 7a shows the relationship between wave thrust and salt marsh erosion rates. Wave thrust values refer to the 1991–2013 period, while erosion rates values refer to the 1930–2013 period. These time frames are meant to represent long-term averages for both wave thrust and erosion rates. The time frame limitation for wave thrust values is connected to wind data availability. We retain that the comparison is acceptable because it spans relatively long temporal intervals, and because it has been previously shown that for the majority of the shoreline erosion rates maintain a similar trend in time. Figures 7b–7d represent the same relationship but for three different salt marsh categories: mainland marshes, likely characterized by muddy substrates [Kennish, 2001]; back-barrier marshes, displaying thin sand deposits driven by storms for some distance into the marshes [Fischer, 1961]; and marshes with abundance of ponds [Elsey-Quirk and Adamowicz, 2016]. The latter category was chosen based on results suggesting the importance of marsh ponds for marsh




Figure 7. (a) Relationship between erosion rate and wave thrust. Grey lines are error bars. A linear fit of gives linear coefficient of $11.915 \text{ m}^2/\text{kN}/\text{yr}$. (b–d) Relationship between erosion rate and wave thrust for three different salt marsh categories corresponding to: mainland marshes, marshes with ponds, and back-barrier marshes. A linear fit of the points gives the following coefficients: 14.32, 8.30, and $7.11 \text{ m}^2/\text{kN}/\text{yr}$, respectively.

deterioration [e.g., *Mariotti and Fagherazzi, 2013*]. In some marshes of Barnegat Bay the number of ponds was artificially increased by Open Marsh Water Management (OMWM), a technique introduced for mosquito control. Recent studies show that marsh area with pools and ponds have less sediment deposition and less surface accretion, while being on average higher [*Elsay-Quirk and Adamowicz, 2016*]. Each cell of the grid used for the computation of wave thrust contains multiple digitization points and erosion rates. For this reason, when comparing wave thrust and erosion rates, data have been binned. Each point in the panel represents average wave thrust and erosion rate for a given bin. Each grey line is the standard deviation for that bin. Binning has been done every 40 points which is the average number of digitalization points per grid cell. When comparing wave thrust, and erosion rate for a specific marsh category (e.g., back-barrier marshes or marsh with ponds, Figures 7b–7d), we first isolated the data belonging to each marsh category, and then we binned the data using the same number of bins than for Figure 7a. Despite the noticeable scatter, wave thrust and salt marsh erosion rates are significantly correlated for each salt marsh category. The best fit occurs when all marshes are considered together. Scatter might arise from a variety of other sources. Among others, differences in vegetation density, bioturbation, and land use might significantly affect salt marsh erosion rates [*Feagin et al., 2009; Paramor and Hughes, 2004*]. Scatter might also arise from heterogeneities in sediment substrate caused by the intermittent presence of overwash sand layers likely causes a large scatter in soil resistance and erosion rates.

5. Discussion

From a long-term analysis of salt marsh lateral retreat in the Barnegat Bay and Little Egg Harbor system, we show that no drastic temporal variations in salt marsh erosion rates are present. Specifically, for the majority of the digitized shoreline, erosion rates evaluated for the period from 1930 to 2007 are similar and of the same order of magnitude of erosion rates that took place from 2007 to 2013. Same results are obtained when considering smaller stretches of marsh boundary and shorter time intervals. Erosion rate appears constant for

the majority of the locations taken into account. Points around Little Egg Inlet are the only locations where a different decrease in erosion rate has occurred. This difference might be due to changes in land use, and coastal restoration practices adopted in the area [Psuty and Ofiara, 2002]. As an example, before 1950 no buildings were present in close proximity to the shoreline, while nowadays areas between Barnegat Inlet and Little Egg inlet are highly developed. Many locations along the barrier island are interested by both soft and hard engineering structures such as groins and jetties exerting shadow effects on the longshore drift which is predominantly to the south [Psuty and Ofiara, 2002]. Beach nourishment also occurred in the area. For instance, the entire barrier island received beach nourishment in 1962–1963 after the Ash Wednesday Northeaster; during the 1990s beach nourishment projects were also carried out in areas south from Barnegat Inlet up to Brant Beach [Psuty and Ofiara, 2002]. Generally, the southern barrier island edges around Little Egg Inlet have significantly expanded; this could have helped protecting the interior part of the salt marsh peninsula where erosion rates have been declining.

According to Figure 4b, erosion rates for the 2007–2010 period are of the same order of magnitude of the rates for the 2010–2013 period. This is especially relevant considering that the second period (2010–2013) was characterized by the occurrence of Hurricane Irene and Hurricane Sandy. Differently from the storms in 2007–2010 that only passed near the area, both Irene and Sandy made landfall in New Jersey and are the only storms in record that caused hurricane force wind in our study area. Both storms caused damages, and the latter is considered as the most destructive in the history of New Jersey [e.g., Blake *et al.*, 2012] (National Hurricane Center; see also Figure S1 in the supporting information). Specifically, Sandy was ranked as 1/900 year event, and the return period for the storm surge has been estimated to be 1570 years, based on generalized extreme value return curves [Brandon *et al.*, 2014]. In fact, Hurricane Sandy crossed the New Jersey coastline at an angle closer to 90° than any other hurricane in record (Figure S1). This was caused by a combination of a blocking high pressure over the western North Atlantic and the interaction with an extratropical upper level disturbance [Hall and Sobel, 2013].

For the area of interest, extreme erosion of salt marshes did not happen in spite of the two large storms. This could be connected to the fact that once a salt marsh is submerged, wave action significantly decreases due to wave energy dissipation above marsh surfaces [e.g., Tonelli *et al.*, 2010; Moller *et al.*, 2014]. For instance, even if the storm surge associated with Hurricane Sandy caused severe damages to the open coast (for example, to sandy beaches, Figure S1), the presence of a high storm surge might have reduced wave impact on salt marshes which are more vulnerable when the water level is around mean sea level. This is in agreement with results presented in Leonard *et al.* [2016] showing in terms of geomorphic work that salt marsh erosion is dictated by average weather conditions (gentle to fresh breeze according to Beaufort wind scale) rather than strong storms. Our findings are also supported by recent results showing high temporal resolution field measurements of salt marsh erosion collected in the Venice Lagoon every month for 1.5 years, at six different locations [Bendoni *et al.*, 2016]. According to these surveys, erosion remains constant in time, and such that the cumulative erosion profile has constant gradient [Bendoni *et al.*, 2016].

For the Barnegat Bay area, the late Holocene relative sea level rise has been around 1.3 ± 0.2 mm/yr [Engelhart *et al.*, 2009]. While this could have affected erosion rates, there is extensive evidence that salt marshes are able to keep pace with sea level rise and maintain their elevation constant with respect to mean sea level thanks to their relatively high accretion rates [e.g., Day *et al.*, 1998; Fagherazzi, 2013; Fagherazzi *et al.*, 2013; Kirwan and Murray, 2008; Kirwan and Megonigal, 2013; Kirwan *et al.*, 2010, 2016]. Eley-Quirk and Adamowicz [2016] measured accretion rates of 1.8 ± 0.8 mm/yr in the Barnegat Bay marshes, which partly offset sea level rise. Herein we mainly focus on spatial variations in erosion rate within Barnegat Bay, and a deeper understanding of the role of sea level rise in salt marsh decline is outside the scope of this work, as it might be difficult to infer differential erosional rates induced by sea level variations at a Bay scale.

The emergent relationship between salt marsh erosion and shoreline roughness suggests that the shape of marsh boundaries is a possible geomorphic signature of the magnitude of erosion events and wetland-vulnerability to wind waves. Specifically, while rough shorelines (high fractal dimension) correspond to slowly eroding salt marshes, smooth shorelines (low fractal dimension) characterize rapidly degrading coastlines. The relationship between marsh boundary geometry and erosion rates is also recognizable from the frequency-magnitude distribution of erosion events. While rapidly eroding shorelines are

characterized by a Gaussian frequency-magnitude distribution, slowly eroding shorelines display a logarithmic distribution. Similar results have been reported in *Leonardi and Fagherazzi* [2014, 2015] for salt marshes at other five locations in Massachusetts and in the Virginia Coast Reserve. By using field data and a cellular automata model, it has been shown that different geomorphic features emerge under high and low wave energy when natural heterogeneities (e.g., different soil properties, bioturbation, seepage erosion, and many others) along marsh boundaries are taken into account. At low wave energy, the existence of a variability in erosional resistance causes differential erosion rates which in turn promote rough marsh boundaries and a long-tailed distribution of erosion events. On the contrary, high-wave energy conditions correspond to a smooth marsh boundary and a Gaussian-like distribution of erosion episodes. For areas normally subject to low-energy conditions, the occurrence of high energy events leads to boundary smoothing [Leonardi and Fagherazzi, 2015]. Variability in erosional resistance could be caused by several factors, such as vegetation cover, soil resistance, biological processes, that are inevitably present along marsh shorelines [e.g., Feagin et al., 2009]. However, it has been shown that such emergent properties can be reproduced by just assuming randomly distributed resistance values and different wave energy conditions [Leonardi and Fagherazzi, 2014, 2015]. The understanding of the role played by salt marsh properties on salt marsh erosion would have benefitted from detailed field measurements of marsh resistance. These field measurements are not present in this manuscript, and this is a limitation that might be addressed in future studies.

Under a climate change scenario, a relationship between the erosion rate and the shape of marsh boundaries could be potentially used to infer changes in external agents, such as changes in mean wave climate or a change in the frequency of extreme events. This relationship can be also used to identify more vulnerable areas by visually evaluating the roughness of marsh boundaries.

6. Conclusions

We digitized more than 100 km of marsh shoreline and calculated salt marsh erosion rates for the Barnegat Bay and Little Egg Harbor system. We further computed shoreline fractal dimension, and the standard deviation of the erosion rates. Finally, we correlated erosion rates to wave thrust values. Our analysis led to three main conclusions:

1. The magnitude of salt marsh erosion rate is recognizable from the morphology of the marsh boundary. Rapidly deteriorating salt marshes have relatively smooth marsh boundary profiles. On the contrary, slowly eroding salt marshes have been found to have rough marsh boundary profiles.
2. For the majority of the shoreline, no drastic temporal changes in erosion rate have been observed. The spatial and temporal distribution of salt marsh erosion rates remained mostly unchanged.
3. We confirm the existence of a relationship between salt marsh erosion rate and wave energy exposure. Specifically, we found a significant correlation between wave thrust and shoreline change.

Acknowledgments

This research was supported by the Department of the Interior Hurricane Sandy Recovery program (project GS2-2D) and by NSF DEB-0621014 (VCR-LTER program), OCE-1238212 (PIE-LTER program). We acknowledge the NJ Office of Information Technology (NJOIT), Office of Geographic Information Systems (OGIS) as the source of aerial images. The numerical model COAWST is freely available for download at <http://woodshole.er.usgs.gov/operations/modeling/COAWST/>. We also thank Amer Suvalic for his contribution to shoreline digitization. We thank the Editors and reviewers for their constructive and insightful comments.

References

- Allen, J. R. L. (1989), Evolution of salt-marsh cliffs in muddy and sandy systems: A qualitative comparison of British west-coast estuaries, *Earth Surf. Process. Landf.*, *14*(1), 85–92.
- Allen, J. R. L., and K. Pye (1992), *Coastal Saltmarshes: Their Nature and Importance*. Salt Marshes, Morphodynamics, Conservation and Engineering Significance, pp. 1–18, Cambridge Univ. Press, Cambridge.
- Aretxabaleta, A. L., B. Butman, and N. K. Ganju (2014), Water level response in back-barrier bays unchanged following Hurricane Sandy, *Geophys. Res. Lett.*, *41*, 3163–3171, doi:10.1002/2014GL059957.
- Armanini, A., M. Righetti, and P. Grisenti (2005), Direct measurement of vegetation resistance in prototype scale, *J. Hydraul. Res.*, *43*(5), 481–487.
- Bakker, J. P., J. de Leeuw, K. S. Dijkema, P. C. Leendertse, H. H. Prins, and J. Rozema (1993), Salt marshes along the coast of the Netherlands, *Hydrobiologia*, *265*, 73, doi:10.1007/BF00007263.
- Bondoni, M., S. Francalanci, L. Cappiotti, and L. Solari (2014), On salt marshes retreat: Experiments and modeling toppling failures induced by wind waves, *J. Geophys. Res. Earth Surf.*, *119*, 603–620, doi:10.1002/2013JF002967.
- Bondoni, M., R. Mel, L. Solari, S. Lanzoni, S. Francalanci, and H. Oumeraci (2016), Insights into lateral marsh retreat mechanism through localized field measurements, *Water Resour. Res.*, *52*, 1446–1464, doi:10.1002/2015WR017966.
- Bernardara, P., D. Schertzer, E. Sauquet, I. Tchiguirinskaia, and M. Lang (2008), The flood probability distribution tail: How heavy is it?, *Stoch. Environ. Res. Risk Assess.*, *22*(1), 107–122.
- Blake, E. S., T. B. Kimberlain, R. J. Berg, J. P. Cangialosi, and J. L. Beven (2012), National Hurricane Center. [Available at http://www.nhc.noaa.gov/data/tcr/AL182012_Sandy.pdf].
- Booij, N., R. C. Ris, and L. H. Holthuijsen (1999), A third-generation wave model for coastal regions: 1. Model description and validation, *J. Geophys. Res.*, *104*, 7649–7666, doi:10.1029/98JC02622.
- Brampton, A. H. (1992), Engineering significance of British saltmarshes, in *Saltmarshes: Morphodynamics, Conservation and Engineering Significance*, edited by J. R. L. Allen and K. Pye, pp. 115–122, Cambridge Univ. Press, Cambridge.

- Brandon, C. M., J. D. Woodruff, J. P. Donnelly, and R. M. Sullivan (2014), How unique was Hurricane Sandy? Sedimentary reconstructions of extreme flooding from New York Harbor, *Sci. Rep.*, *4*, doi:10.1038/srep07366.
- Chapman, D. C. (1985), Numerical treatment of cross-shelf open boundaries in a barotropic coastal ocean model, *J. Phys. Oceanogr.*, *15*, 1060–1075.
- Chassignet, E. P., H. Arango, D. Dietrich, T. Ezer, M. Ghil, D. B. Haidvogel, C.-C. Ma, A. Mehra, A. M. Paiva, and Z. Sirkes (2000), DAMÉE-NAB: The base experiments, *Dynam. Atmos. Oceans*, *32*(3–4), 155–183, doi:10.1016/S0377-0265(00)00046-4.
- Cooper, A., T. McCann, and D. Rogers (2009), Northern Ireland countryside survey 2007: Broad habitat change 1998–2007, Northern Ireland Environment Agency, Belfast.
- Cope, S. N., A. P. Bradbury, and M. Gorcznska (2008), Solent Dynamic Coast Project: Main Report, New Forest District Council/Channel Coast Observatory.
- Costanza, R., O. Pérez-Maqueo, M. L. Martinez, P. Sutton, S. J. Anderson, and K. Mulder (2008), The value of coastal wetlands for hurricane protection, *AMBIO, J. Hum. Environ.*, *37*(4), 241–248, doi:10.1579/0044-7447(2008)37[241:TVOCWF]2.0.CO;2.
- Cowart, L., D. R. Corbett, and J. P. Walsh (2011), Shoreline change along sheltered coastlines: Insights from the Neuse River estuary, NC, USA, *Rem. Sens.*, *3*(7), 1516–1534, doi:10.3390/rs3071516.
- Cundy, A. B., and I. W. Croudace (1996), Sediment accretion and recent sea-level rise in the Solent, southern England: Inferences from radiometric and geochemical studies, *Estuarine Coastal Shelf Sci.*, *43*, 449–467.
- Day, J. W., S. Francesco, R. Andrea, and D. Are (1998), Rapid deterioration of a Salt Marsh in Venice Lagoon, Italy, *J. Coastal Res.* *14*(2), 583–90. [Available at <http://www.jstor.org/stable/4298811>].
- Defne, Z., and N. Ganju (2014), Quantifying the residence time and flushing characteristics of a shallow, back-barrier estuary: Application of hydrodynamic and particle tracking models, *Estuaries Coasts*, 1–16, doi:10.1007/s12237-014-9885-3.
- Department of the Army, Waterways Experiment Station, Corps of Engineers, and Coastal Engineering Research Center (1984), Shore protection manual, vol. 1, 4th ed., 532 pp., Department of the Army, Waterways Experiment Station, Corps of Engineers, and Coastal Engineering Research Center, Washington, D. C.
- Dubuc, B., J. F. Quiniou, C. Roques-Carmes, C. Tricot, and S. W. Zucker (1989), Evaluating the fractal dimension of profiles, *Phys. Rev. A*, *39*(3), 1500–1512.
- Elsley-Quirk, T., and S. C. Adamowicz (2016), Influence of physical manipulations on short-term salt marsh morphodynamics: Examples from the North and Mid-Atlantic Coast, USA, *Estuaries Coasts*, *39*(2), 423–439.
- Elsner, J. B., and A. B. Kara (1999), *Hurricanes of the North Atlantic: Climate and Society*, Oxford Univ. Press, New York.
- Engelhart, S. E., B. P. Horton, B. C. Douglas, W. R. Peltier, and T. E. Törnqvist (2009), Spatial variability of late Holocene and 20th century sea-level rise along the Atlantic coast of the United States, *Geology*, *37*(12), 1115–1118.
- Environment Agency (2011), The Extent of Saltmarsh in England and Wales: 2006–2009, Environment Agency, Bristol.
- Fagherazzi, S. (2013), The ephemeral life of a salt marsh, *Geology*, *41*(8), 943–944, doi:10.1130/focus082013.1.
- Fagherazzi, S. (2014), Coastal processes: Storm-proofing with marshes, *Nat. Geosci.*, *7*(10), 701–702.
- Fagherazzi, S., G. Mariotti, P. L. Wiberg, and K. J. McGlathery (2013), Marsh collapse does not require sea level rise, *Oceanography*, *26*(3), 70–77, doi:10.5670/oceanog.2013.47.
- Feagin, R. A., S. M. Lozada-Bernard, T. M. Ravens, I. Möller, K. M. Yeager, and A. H. Baird (2009), Does vegetation prevent wave erosion of salt marsh edges?, *Proc. Natl. Acad. Sci.*, *106*(25), 10,109–10,113.
- Fischer, A. G. (1961), Stratigraphic record of transgressing seas in light of sedimentation on Atlantic coast of New Jersey, *AAPG Bull.*, *45*(10), 1656–1666.
- Flather, R. A. (1976), A tidal model of the northwest European continental shelf, *Mem. Soc. R. Sci. Liege*, *6*, 141–164.
- Foster, N. M., M. D. Hudson, S. Bray, and R. J. Nicholls (2013), Intertidal mudflat and saltmarsh conservation and sustainable use in the UK: A review, *J. Environ. Manage.*, *126*, 96–104.
- Francalanci, S., M. Bondoni, M. Rinaldi, and L. Solari (2013), Ecomorphodynamic evolution of salt marshes: Experimental observations of bank retreat processes, *Geomorphology*, *195*, 53–65, doi:10.1016/j.geomorph.2013.04.026.
- Haidvogel, D. B., et al. (2008), Ocean forecasting in terrain-following coordinates: Formulation and skill assessment of the Regional Ocean Modeling System, *J. Comput. Phys.*, *227*(7), 3595–3624, doi:10.1016/j.jcp.2007.06.016.
- Hall, T. M., and A. H. Sobel (2013), On the impact angle of Hurricane Sandy's New Jersey landfall, *Geophys. Res. Lett.*, *40*, 2312–2315, doi:10.1002/grl.50395.
- Järvelä, J. (2002), Flow resistance of flexible and stiff vegetation: A flume study with natural plants, *J. Hydrol.*, *269*(1), 44–54.
- Kennish, M. J. (2001), Physical description of the Barnegat Bay—Little Egg Harbor estuarine system, *J. Coastal Res.*, 13–27, doi:10.2307/25736225.
- Kirwan, M. L., and J. P. Megonigal (2013), Tidal wetland stability in the face of human impacts and sea-level rise, *Nature*, *504*(7478), 53–60.
- Kirwan, M. L., and A. B. Murray (2008), Ecological and morphological response of brackish tidal marshland to the next century of sea level rise: Westham Island, British Columbia, *Global Planet. Change*, *60*(3), 471–486.
- Kirwan, M. L., G. R. Guntenspergen, A. D'Alpaos, J. T. Morris, S. M. Mudd, and S. Temmerman (2010), Limits on the adaptability of coastal marshes to rising sea level, *Geophys. Res. Lett.*, *37*, L23401, doi:10.1029/2010GL045489.
- Kirwan, M. L., S. Temmerman, E. E. Skeehan, G. R. Guntenspergen, and S. Fagherazzi (2016), Overestimation of marsh vulnerability to sea level rise, *Nat. Clim. Change*, *6*(3), 253–260.
- Koch, E. W. (1996), Hydrodynamics of a shallow *Thalassia testudinum* bed in Florida, USA, in *Seagrass Biology—Proceedings of an International Workshop*, edited by J. Kuo, R. C. Phillips, D. I. Walker, and H. Kirkman, pp. 105–110, Western Australia Museum, Perth, Australia.
- Kumar, N., G. Voulgaris, and J. C. Warner (2011), Implementation and modification of a three-dimensional radiation stress formulation for surf zone and rip-current applications, *Coast. Eng.*, *58*(12), 1097–1117, doi:10.1016/j.coastaleng.2011.06.009.
- Lathrop, R. G., Jr., and J. A. Bognar (2001), Habitat loss and alteration in the Barnegat Bay Region, *J. Coastal Res.*, 212–228, doi:10.2307/25736235.
- Le Hir, P., W. Roberts, O. Cazaillet, M. Christie, P. Bassoullet, and C. Bacher (2000), Characterization of intertidal flat hydrodynamics, *Cont. Shelf Res.*, *20*(12), 1433–1459.
- Lee, M. (2001), Coastal defence and the Habitats Directive: Predictions of habitat change in England and Wales, *Geogr. J.*, *167*(1), 39–56.
- Leonardi, N., and S. Fagherazzi (2014), How waves shape salt marshes, *Geology*, doi:10.1130/G35751.1.
- Leonardi, N., and S. Fagherazzi (2015), Effect of local variability in erosional resistance affects large scale morphodynamic response of salt marshes to wind waves resistance affects large scale morphodynamic, *Geophys. Res. Lett.*, *42*, 5872–5879, doi:10.1002/2015GL064730.
- Leonardi, N., N. K. Ganju, and S. Fagherazzi (2016), A linear relationship between wave power and erosion determines salt-marsh resilience to violent storms and hurricanes, *Proc. Natl. Acad. Sci.*, *113*(1), 64–68.

- Limpert, E., W. A. Stahel, and M. Abbt (2001), Log-normal distributions across the Sciences: Keys and clues: On the charms of statistics, and how mechanical models resembling gambling machines offer a link to a handy way to characterize log-normal distributions, which can provide deeper insight into v , *BioScience*, 51(5), 341–352.
- Malamud, B. D. (2004), Tails of natural hazards, *Phys. World*, 17(8), 25.
- Marani, M., A. D'Alpaos, S. Lanzoni, and M. Santalucia (2011), Understanding and predicting wave erosion of marsh edges, *Geophys. Res. Lett.*, 38, L21401, doi:10.1029/2011GL048995.
- Mariotti, G., and S. Fagherazzi (2013), Critical width of tidal flats triggers marsh collapse in the absence of sea-level rise, *Proc. Natl. Acad. Sci.*, 110(14), 5353–5356.
- Mariotti, G., S. Fagherazzi, P. L. Wiberg, K. J. McGlathery, L. Carniello, and A. Defina (2010), Influence of storm surges and sea level on shallow tidal basin erosive processes, *J. Geophys. Res.*, 115, C11012, doi:10.1029/2009JC005892.
- McLoughlin, S., P. Wiberg, I. Safak, and K. McGlathery (2015), Rates and forcing of marsh edge erosion in a shallow coastal bay, *Estuaries Coasts*, 38(2), 620–638, doi:10.1007/s12237-014-9841-2.
- Möller, I. (2006), Quantifying saltmarsh vegetation and its effect on wave height dissipation: Results from a UK East coast saltmarsh, *Estuarine Coastal Shelf Sci.*, 69(3), 337–351.
- Möller, I., T. Spencer, J. R. French, D. J. Leggett, and M. Dixon (1999), Wave transformation over salt marshes: A field and numerical modelling study from North Norfolk, England, *Estuarine Coastal Shelf Sci.*, 49(3), 411–426.
- Moller, I., et al. (2014), Wave attenuation over coastal salt marshes under storm surge conditions, *Nat. Geosci.*, 7(10), 727–731.
- Mudd, S. M., S. M. Howell, and J. T. Morris (2009), Impact of dynamic feedbacks between sedimentation, sea-level rise, and biomass production on near-surface marsh stratigraphy and carbon accumulation, *Estuarine Coastal Shelf Sci.*, 82(3), 377–389.
- Murphy, A. H., and E. S. Epstein (1989), Skill scores and correlation coefficients in model verification, *Mon. Weather Rev.*, 117(3), 572–582, doi:10.1175/1520-0493(1989)117<0572:SSACI>2.0.CO;2.
- Nepf, H. M. (1999), Drag, turbulence, and diffusion in flow through emergent vegetation, *Water Resour. Res.*, 35, 479–489, doi:10.1029/1998WR900069.
- NOAA National Ocean Survey (2012), National Ocean Service Hydrographic Survey data, National Oceanic and Atmospheric Administration, [Available at <http://www.ngdc.noaa.gov/mgg/bathymetry/hydro.html>, Accessed 2012.]
- Orlanski, I. (1976), A simple boundary condition for unbounded hyperbolic flows, *J. Comp. Sci.*, 21(3), 251–269.
- Paramor, O. A. L., and R. G. Hughes (2004), The effects of bioturbation and herbivory by the polychaete *Nereis diversicolor* on loss of saltmarsh in south-east England, *J. Appl. Ecol.*, 41(3), 449–463.
- Pethick, J. (1993), Shoreline adjustments and coastal management: Physical and biological processes under accelerated sea-level rise, *Geogr. J.*, 159, 162–168.
- Plater, A. J., A. J. Long, C. D. Spencer, and R. A. Delacour (1999), The stratigraphic record of sea-level change and storms during the last 2000 years: Romney Marsh, southeast England, *Quaternary Int.*, 55(1), 17–27.
- Prager, E. J., and R. B. Halley (1999), The influence of seagrass on shell layers and Florida Bay mudbanks, *J. Coastal Res.*, 15, 1151–1162.
- Prietas, A. M., G. Mariotti, N. Leonardi, and S. Fagherazzi (2015), Coupled Wave Energy and Erosion Dynamics along a Salt Marsh Boundary, Hog Island Bay, Virginia, USA, *J. Mar. Sci. Eng.*, 3(3), 1041–1065.
- Psuty, N. P., and D. D. Ofiara (2002), *Coastal Hazard Management: Lessons and Future Directions From New Jersey*, Rutgers Univ. Press, New Brunswick.
- Pye, K., and P. W. French (1993), *Targets for Coastal Habitat Re-creation*, University Press, Cambridge.
- Ris, R. C., L. H. Holthuijsen, and N. Booij (1999), A third-generation wave model for coastal regions: 2. Verification, *J. Geophys. Res.*, 104, 7667–7681, doi:10.1029/1998JC900123.
- Schuerch, M., T. Dolch, K. Reise, and A. T. Vafeidis (2014), Unravelling interactions between salt marsh evolution and sedimentary processes in the Wadden Sea (southeastern North Sea), *Prog. Phys. Geogr.*, 38(6), 691–715.
- Schwimmer, R. A. (2001), Rates and processes of marsh shoreline erosion in Rehoboth Bay, Delaware, U.S.A., *J. Coastal Res.*, 17(3), 672–683, doi:10.2307/4300218.
- Shchepetkin, A. F., and J. C. McWilliams (2005), The Regional Oceanic Modeling System (ROMS): A split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Modell.*, 9(4), 347–404, doi:10.1016/j.ocemod.2004.08.002.
- Suk, N. S., Q. Guo, and N. P. Psuty (1999), Suspended solids flux between salt marsh and adjacent bay: A long-term continuous measurement, *Estuarine Coastal Shelf Sci.*, 49(1), 61–81.
- Temmerman, S., and M. L. Kirwan (2015), Building land with a rising sea, *Science*, 349(6248), 588–589.
- Temmerman, S., G. Govers, P. Meire, and S. Wartel (2003), Modelling long-term tidal marsh growth under changing tidal conditions and suspended sediment concentrations, Scheldt estuary, Belgium, *Mar. Geol.*, 193(1), 151–169.
- Temmerman, S., T. J. Bouma, G. Govers, Z. B. Wang, M. B. De Vries, and P. M. J. Herman (2005), Impact of vegetation on flow routing and sedimentation patterns: Three-dimensional modeling for a tidal marsh, *J. Geophys. Res.*, 110, F04019, doi:10.1029/2005JF000301.
- Temmerman, S., P. Meire, T. J. Bouma, P. M. J. Herman, T. Ysebaert, and H. J. De Vriend (2013), Ecosystem-based coastal defence in the face of global change, *Nature*, 504(7478), 79–83.
- Tonelli, M., S. Fagherazzi, and M. Petti (2010), Modeling wave impact on salt marsh boundaries, *J. Geophys. Res.*, 115, C09028, doi:10.1029/2009JC006026.
- Tsujiimoto, T., T. Kitamura, Y. Fujii, and H. Nakagawa (1996), Hydraulic resistance of flow with flexible vegetation in open channel, *J. Hydroscience Hydraul. Eng.*, 14(1), 47–56.
- UK National Ecosystem Assessment (2011), *UK National Ecosystem Assessment: Chapter 11 Coastal Margins*, UNEP/WCMC, Cambridge.
- van de Koppel, J., D. van der Wal, J. P. Bakker, and P. M. Herman (2005), Self-organization and vegetation collapse in salt marsh ecosystems, *Am. Nat.*, 165(1), E1–E12.
- van der Wal, D., and K. Pye (2004), Patterns, rates and possible causes of saltmarsh erosion in the Greater Thames area (UK), *Geomorphology*, 61(3), 373–391.
- Warner, J. C., C. R. Sherwood, R. P. Signell, C. K. Harris, and H. G. Arango (2008), Development of a three-dimensional, regional, coupled wave, current, and sediment-transport model, *C. R. Geosci.*, 34(10), 1284–1306, doi:10.1016/j.cageo.2008.02.012.
- Warner, J. C., B. Armstrong, R. He, and J. B. Zambon (2010), Development of a Coupled Ocean–Atmosphere–Wave–Sediment Transport (COAWST) Modeling System, *Ocean Modell.*, 35(3), 230–244, doi:10.1016/j.ocemod.2010.07.010.
- Zedler, J. B., and S. Kercher (2005), WETLAND RESOURCES: Status, trends, ecosystem services, and restorability, *Annu. Rev. Environ. Resour.*, 30(1), 39–74, doi:10.1146/annurev.energy.30.050504.144248.
- Zhao, H., and Q. Chen (2013), Modeling attenuation of storm surge over deformable vegetation: Methodology and verification, *J. Eng. Mech.*, 140(12), 4014090, doi:10.1061/(ASCE)EM.1943-7889.0000704.