

On-Line Supplemental Information for
**Impacts of Hurricanes Katrina and Rita on the Microbial Landscape of
the New Orleans Area**

C.D. Sinigalliano^{1,2}, M.L. Gidley¹, T. Shibata¹, D. Whitman³, T.H. Dixon¹, E. Laws^{4,5}, A. Hou⁴, D. Bachoon⁶, L. Brand¹, L. Amaral-Zettler^{7,8}, R. Gast^{7,9}, G. F. Steward⁵, O. D. Nigro⁵, R. Fujioka⁵, W.Q. Betancourt⁵, G. Vithanage⁵, J. Mathews¹, L.E. Fleming¹, H. M. Solo-Gabriele¹

¹ NSF-NIEHS Oceans and Human Health Center, Rosenstiel School of Marine and Atmospheric Science, University of Miami, Coral Gables, Florida

² Southeast Environmental Research Center, Florida International University, Miami, Florida

³ Department of Earth Sciences, Florida International University, Miami, Florida

⁴ Louisiana State University, School of the Coast and Environment, Baton Rouge, LA

⁵ Pacific Research Center for Marine Biomedicine, School of Ocean and Earth Science and Technology, University of Hawaii, Honolulu, Hawaii

⁶ Georgia College and State University, Milledgeville, Georgia

⁷ Woods Hole Center for Oceans and Human Health, Woods Hole, Massachusetts

⁸ Marine Biological Laboratory, Woods Hole, MA

⁹ Woods Hole Oceanographic Institution, Woods Hole, MA

This PDF file of supplemental information contains:

Materials and Methods

Supplemental Information Tables S-1 to S-5

Materials and Methods

Remote Sensing Image Analysis for Inundation Depths and Flood Volume

Computations: The flooding in New Orleans is clearly delineated on a pan-sharpened color-infrared SPOT satellite image taken 30 August 2005. A digital map of flood depth on the NE bank of the Mississippi River was constructed by combining observed areas of flooding on the SPOT image with high resolution digital elevation data (1). Elevations were extracted along both the edge of flooding and on exposed highway surfaces in the flooded interior to create a set of flood heights. These heights varied spatially throughout the study area, and were used to construct a model of flood height composed of several contiguous polygons of constant height. The boundaries between polygons generally corresponded to barriers to water flow such as levies, floodwalls, highways, or natural topographic ridges. Finally, a raster model of flood depth was calculated by subtracting a 10 m resolution digital elevation model extracted from the USGS National Elevation Database (<http://seamless.usgs.gov>) from the flood height model.

Sample Collection: Surface water samples from Lake Pontchartrain and from canals were collected in sterile 10L polypropylene carboys, polypropylene bottles, or acid-washed and ethanol-rinsed polyethylene cubitainers[®] (Hedwin Corporation) from the locations indicated in Fig. 1 and Table S-1. Water samples were kept in a cooler with ice packs maintained between 12 and 24 °C (*Vibrio* analyses) or on ice (all other analyses) until return to the laboratory within approximately 4-6 hours. Approximately 100g sediment samples, both from canal shorelines and from dried floodwater sediment

deposits in residential yards, were collected into sterile 250mL specimen cups from the top 3-5 cm of sediment using sterile plastic spatulas. In the case of shoreline samples from lakes and canals, the sediment was collected from just above the water line. Sediment samples were also kept on ice until return to the laboratory. Additional environmental sampling was also conducted at this same time that will be reported elsewhere. Upon return to the laboratory, water samples were used directly for processing and analysis by chromogenic substrate assay (IDEXX, Westbrook, ME), plate count enumeration, and extraction of nucleic acids. In the case of sediment samples, 30g of sediment were resuspended in 300mL of sterile 1X PBS solution in sterile Whirlpak[®] bags (Nasco, Ft. Atkinson, WI), then sealed and vigorously agitated and kneaded to resuspend the bacteria into the PBS. The resuspended samples were allowed to settle for 10 minutes, then the supernatant was processed for chromogenic substrate, and nucleic acid extraction in a similar fashion to the environmental water samples.

Total Microbial Community DNA Extractions: Two different collection and DNA extraction methods were employed: (1) Filtration through polyethersulfone membrane capsule filters (0.22 μm pore size Sterivex[™], Millipore, Billerica, MA) followed by DNA extraction and purification using a modified salt precipitation method (PUREGENE[®], Gentra Systems, Minneapolis, MN). This high-yield method was used for community sequence analysis, for pathogen screening, and DNA archiving. (2) Filtration through flat polycarbonate (PC) track-etched membranes (0.45 μm pore size, 47 mm diameter; GE Osmonics, Minnetonka, MN) followed by cell lysis using a bead-

based homogenizer (FastPrep[®], Qbiogene, Solon, OH) and DNA purification using a silica-based procedure (FastDNA[®] Spin Kit for Soil, Qbiogene). This method was used for samples to be analyzed by quantitative real-time PCR (qPCR).

For the first method, subsamples (1 L) of lake and canal water were filtered through the Sterivex[™] filter capsules using positive pressure with a peristaltic pump or by applying vacuum to the filter outlet. Filters were pumped to dryness then filled with 2mL of lysis buffer from the PUREGENE[®] Genomic DNA Isolation Kit, sealed with Luer end caps, wrapped in parafilm, and then stored frozen at -80°C until further processing. Filter capsules were subsequently thawed and 10µL of Lytic Enzyme solution (supplied with kit) was added. Filters were resealed and incubated at 37°C for 30 min on the rotating carousel of a hybridization oven. Ten microliters of Proteinase K solution (supplied with kit) was then added to the filters, which were resealed and incubated at 65°C for 1 h with rotation in a hybridization oven. The lysate from all the filters for a given sample was withdrawn using a 3cc syringe and combined into a 15 ml tube, cooled to room temperature and then DNA was purified from each pooled sample according to PUREGENE[®] kit manufacturer's instructions. Resulting DNA was quantified and archived at -80°C for subsequent analysis.

For the second method, bacteria from 100mL water samples were collected on flat PC filters, which were then transferred to the vials of "lysing matrix E" from the Fast DNA Spin Kit for Soil (QBiogene, Irvine, CA), and one *Lactococcus* Sample Prep Control SmartBead[™] (Cepheid) containing lyophilized *Lactococcus* control cells at a known concentration was added to each vial. The DNA from the PC filters (along with

DNA from the *Lactococcus* SPC bead) was then extracted in these bead-beating vials using the Fast DNA Spin Kit for Soil (QBiogene) according to manufacturer's instructions, except that the filter replaced the sediment pellet. Eluted DNA was stored at -80°C for subsequent analysis by qPCR. In the case of sediment samples, 100mL of the supernatant of resuspended sediment bacteria in 1X PBS was collected on PC filters and extracted with the Fast DNA Spin Kit for Soil as described above for water samples.

Enterococci and *E. coli* Using Membrane Filtration Method: Standardized methods were utilized (Method 1600 (2) and Method 1103.1 (3)) with the processing of water samples within 6 hours. In brief, the method requires the filtration of the water sample through a membrane which captures the bacteria. This membrane is then placed on the selective medium. For enterococci, mEI media (membrane-Enterococcus Indoxyl-b-D-Glucooside) and for *E. coli*, mTEC media (membrane-Thermotolerant *Escherichia coli*) were used. The enterococci plates were incubated at 41°C for 24h, and colonies with blue halos counted. The *E. coli* plates were incubated at 35°C for 2h and then incubated at 44.5°C for 22h. After incubation, the filter was transferred to a pad saturated with urea substrate and yellow, yellow-green and yellow brown colonies were counted after 15 min. Both *E. coli* and enterococci were reported as CFU/100ml.

Enterococci Using Chromogenic Substrate Method: Duplicate 10-fold, 100-fold, and 1000-fold dilutions of each water sample were prepared in sterile 1X PBS. Each dilution was then processed and assayed using the Enterolert™ enterococci test kit and Quanti-Tray/2000 (IDEXX, Westbrook, ME), according to manufacturer's directions. In the

case of sediment samples, duplicate dilutions in sterile 1X PBS were made of the initial sediment bacterial resuspension described above, processed, and assayed in the same manner as water samples with the chromogenic substrate method.

Enterococci and *E. coli* Using Quantitative PCR Method: Ten-fold and 100-fold dilutions of DNA extracts from water or sediment samples were analyzed for relative abundance of enterococci and *E. coli* using real-time quantitative PCR water quality assay kits from Cepheid (Sunnyvale, CA).

For enterococci enumeration, 5 μ L samples of each DNA dilution from the PC filter extracts was amplified using the “Total *Enterococcus* Scorpion® Primer and Probe Set and Positive Control DNA” beads and the “OmniMix-HS™ Lyophilized PCR Master Mix” beads according to manufacturer’s instructions. Each Total *Enterococcus* assay bead contains a proprietary mix of fluorescently labeled Scorpion primers targeting *Enterococcus*-specific portions of the 16s rDNA, internal amplification control DNA, and a second set of Scorpion primers that target the internal control DNA all lyophilized together in amounts sufficient for four 25 μ L reactions per bead. The Omnimix-HS™ beads (Cepheid) contain all other buffers, reagents, nucleotides, enzymes, (except for water and target DNA) lyophilized into a bead for two 25 μ L reactions. Amplifications were done in batches of 16 reactions on a SmartCycler™ real-time PCR thermocycler (Cepheid), by dissolving 4 enterococci primer beads with 8 OmniMix-HS™ beads, and 340 μ L of sterile molecular-grade water. This mixture was aliquotted in 20 μ L volumes into sterile Smartcycler-Tubes™ (Cepheid), 5 μ L of template DNA was added to each

tube, which was then sealed, centrifuged, and cycled using the following parameters: 45 cycles of 95°C for 5 sec and 62°C for 43 sec, with the FCTC25 dye set calibration, FAM threshold manually set at 7, and other settings at default. For each amplification set, negative controls of sterile water in place of DNA template were run, as well as positive controls of known amounts of DNA from *Enterococcus faecalis*. Standard curves for quantitation were run in duplicate and were generated by using 5 serial dilutions of “Enterococcus Postive Control DNA Beads” (Cepheid) which consisted of lyophilized beads containing *E. faecalis* DNA of known concentration that had been calibrated to CFU equivalents of enterococci. Quantitation of enterococci in unknown samples could thus be determined using these standard qPCR amplification curves.

In a similar fashion, *E. coli* abundance was enumerated using dilutions of template DNA from the PC filter extracts to be amplified by qPCR using the conditions described above for *Enterococcus*, but using “*E. coli* Species Scorpion Primer and Probe Set with Positive Control DNA” beads (Cepheid), and standard curves generated from “*E. coli* Postive Control DNA” beads (Cepheid). Assay and cycling conditions were the same as for the *Enterococcus*.

For the *Lactococcus* Sample Processing Control (SPC), dilutions of template DNA from the PC filter extracts were amplified in a similar manner as above, except that the primers beads used were from the “*Lactococcus* SPC SmartBeads: Scorpion Primer and Probe Set with Positive Internal Control DNA.”

Clostridium perfringens: Water samples for *C. perfringens* were assayed using membrane filtration on mCP agar (4). Sample aliquots of 100 ml were placed into a water bath and pasteurized for 15 minutes at 60°C. 25 ml aliquots of pasteurized samples were filtered thru 0.45 µm pore size membranes (GN-6 Metrice1; Pall Gelman, East Hills NY) and placed filter side up onto mCP agar. Agar plates were incubated anaerobically at 45°C for 24 hours. After 24 hours, plates containing yellow (sucrose positive) colonies were exposed to ammonium hydroxide fumes for 20 seconds and resulting pink or magenta colonies were counted.

FRNA coliphages: FRNA coliphages were assayed on *E. coli* HS(pFamp)R using the most probable number assay described by Sobsey *et al.*, (5). The assay involves enrichment of 100 mL of water which are subsequently divided into aliquots of 30 (3), 3 (3) and 0.3 (3) mL. These aliquots are seeded with *E. coli* HS(pFamp)R (0.5 mL) and enriched with 10X concentrated TSB (5 mL) with Ampicilin and Streptomycin (100X) including addition of MgCl₂ (4 M) (1.25 mL). Aliquots were incubated for 24 h at 37 °C. After the incubation period, subsamples are spun at high speed and aliquots of 5µL were spotted onto a lawn of *E. coli* HS(pFamp)R. Lysis zones were recorded and positive or negative results were entered into a MPN program to derive a most probable number of plaque-forming units (PFU) along with a 95% confidence interval (5).

Bacteroidales: Fecal *Bacteroidales* as a measurement of general fecal pollution was measured by standard PCR using reamplification or by real time PCR using the *Taq*

nuclease assay (TNA) described by Dick and Field (6) which targets a segment of the *Bacteroidales* 16S rRNA gene. The combination of primers and fluorogenic probe enables amplification of 16S rDNA of *Bacteroidales* from human, cow, dog, cat, pig, elk, deer and gull feces. Although the TNA was originally developed as a quantitative assay, the data derived from water samples reported here were based on presence/absence evaluations.

***Bifidobacterium*:** A nested PCR protocol to detect *B. adolescentis* was performed on the extracted DNA samples as described by King, et al. (7). The specificity of this assay has previously been tested against feces from human, cow, horse, pig, dog, chicken, and goose, as well as against sewage samples and environmental water samples known to be contaminated with either human or animal feces, and has only shown detection of human-source fecal contamination.

The first step consisted of an amplification using the universal 16S rRNA primer, 785R (CTACCAGGGTATCTAATCC) and the *Bifidobacterium* genus specific primer, IM26F (GATTCTGGCTCAGGATGAACG). Each PCR reaction contained a 30 μ L volume with 0.3 mM dNTP, 3 mM MgCl₂, 1 U Taq DNA polymerase, 1X PCR reaction buffer, and ~30 ng of template DNA. The samples were run on a thermocycler (Techne Model TC-312, Burlington, NJ) under the following conditions: initial denaturing at 94°C for 5 min; 45 cycles of 94°C for 30sec, 48°C for 30sec, and 72°C for 30sec; and final elongation at 72°C for 5 min. Product from this PCR reaction was then used as template for a second PCR mixture.

For this second PCR protocol, the template was amplified using *Bifidobacterium adolescentis* species-specific primers ADO1 (CTCCAGTTGGATGCATGT), and ADO2 (CGAAGGTTGCTCCCAGT) (8). One microliter of product from the first PCR was added to a 30 μ L reaction mixture containing the same concentrations of MgCl₂, 1X reaction buffer, dNTP, and Taq as above. These samples were run on the thermal cycler under the following conditions: initial denaturing at 94°C for 5 min; 45 cycles of 94°C for 30sec, 48°C for 20sec, 55°C for 30sec, and 72°C for 1 min; and final elongation at 72°C for 5 min. Products from both the first and seconds rounds of PCR were subjected to electrophoresis in a 1.5% agarose gel stained with ethidium bromide. The detection limit for this method was 200 *Bifidobacterium adolescentis* cells. This was determined using *B. adolescentis* genomic DNA ATCC ® number 15703D™ .

Vibrio Species: A portion of each water sample was diluted 1:100 into sterile solution of peptone (0.1% w:v) and NaCl (3% w:v) in water (PS; (9)). Twenty-five milliliters of undiluted or diluted sample were filtered in duplicate onto 0.45 μ m pore size mixed cellulose ester membrane filters (GN-6 Metrice1; Pall Gelman, East Hills NY). One of each of the duplicate filters was placed face up on Thiosulfate-Citrate-Bile salts-Sucrose (TCBS) agar (Accumedia Manufacturers, Inc. Lansing, MI) and the other on CHROMagar® Vibrio (DRG International Inc., Mountainside, NJ). Plates were incubated overnight at 37°C. Total colonies were enumerated and representative colonies from each medium were picked with sterile plastic loops and gridded onto fresh plates of the opposing medium (colonies from TCBS were gridded onto CHROMagar Vibrio and vice

versa) and incubated overnight at 37°C. Initial putative species identifications were based on color changes on both chromogenic media. Gridded colonies were picked with a sterile loop and resuspended in 1 ml of PS with glycerol (15% v:v) contained in sterile cryovials. The resuspended colonies were frozen on dry ice for shipment to Hawaii and then maintained at -80°C. Clonal populations were recovered from the glycerol stocks by serial streaking then maintained on T1N1 (10% NaCl, 10% Tryptone) agar slants overlaid with mineral oil.

For PCR assays, cells from clonal colonies were resuspended in 100µL TE buffer in thin-walled PCR tubes using sterile loops. Tubes were heated to 100°C for 10 min in a thermal cycler to release bacterial DNA then centrifuged to pellet debris. A portion of each supernatant was used as the template for subsequent PCR reactions. PCR-based assays were performed to determine presence or absence of intergenic transcribed spacer sequences diagnostic for *V. cholerae* (VcITS; (10)), a *V. parahaemolyticus*-specific thermolabile hemolysin gene (*tlh*)(11), or the *V. vulnificus*-specific hemolysin A gene (*vvhA*)(12). As a positive control, samples were amplified with general bacterial primers for the 16S rRNA gene (13) with minor modifications. Samples that failed to amplify with the 16S rRNA gene primers were cleaned using InstaGene™ Matrix (BioRad Laboratories, Hercules, CA) following the manufacturer's protocol for bacterial DNA isolation, except that the initial water wash steps were omitted. Instead, fifty microliters of cell suspension in TE buffer was added directly to 200 ul of InstaGene™ Matrix buffer. Samples were then heated for 15 minutes at 56°C, vortexed, then heated for 8 minutes at 100°C, vortexed, then centrifuged at 10,000 X g for 2 minutes. Twenty

microliters of the cleaned sample was used in the PCR analysis.

PCR assays were performed with the Expand PCR System (Roche Diagnostics, Alameda, CA) and contained 1X reaction buffer, 1 μ L template DNA extract, 0.25 μ M of each primer, and 1.5 mL MgCl₂. All PCR reactions were initiated with a 4 min hold at 95°C followed by 35 cycles of denaturation at 94°C for 1 min, annealing at a reactions-specific temperature for 1 min, and extension at 72°C for 1 min.

The primer sequences (all written 5'-3') and the annealing temperature used with each primer set are as follows: VcITS-F (TTA AGC STT TTC RCT GAG AAT G) and VcITS-R (AGT CAC TTA ACC ATA CAA CCC G) at 60°C annealing, TLH-F (AAA GCG GAT TAT GCA GAA GCA CTG) and TLH-R (GCT ACT TTC TAG CAT TTT CTC TGC) at 58°C, VVH-F (TTC CAA CTT CAA ACC GAA CTA TGA C) and VVH-R (ATT CCA GTC GAT GCG AAT ACG TTG) at 63°C annealing, and 27F-B (AGR GTT YGA TYM TGG CTC AG) and 1429R (GGY TAC CTT GTT ACG ACT T) at 50°C annealing.

Presence and size of amplification products was determined by agarose gel electrophoresis, staining with SYBRsafe fluorescent DNA stain (Invitrogen, Carlsbad, CA), and visualization and documentation using UV transillumination and digital image capture (VersaDoc 3000, BioRad Laboratories, Hercules, CA).

Legionella: A nested PCR amplification procedure using primers designed for the genus *Legionella* (p1.2 & cp3.2 (14), LEG225 & LEG858 (15)) was used to detect the presence of *Legionella* species in surface water samples from the transect samples taken in Lake

Pontchartrain. The primers used in this method are specific for members of the genus, and sequencing is necessary to determine whether the amplified fragments represent human pathogens. In addition, amplification using primers specific for a portion of the *mip* gene of *L. pneumophila* were carried out following the method of Templeton et al. (16). One microliter of DNA extract (diluted 1:100) was used as the template for the PCR reactions, and products recovered using the *L. pneumophila* specific primers were sequenced to confirm their origin and similarity to published *L. pneumophila* *mip* sequences. Sequencing was carried out on PCR products recovered by band isolation (Zymoclean Gel DNA Recovery Kit; Zymo Research, Orange, CA) using the amplification primers and the ABI Prism® BigDye™ Terminator Cycle Sequencing Ready Reaction Kit (Applied Biosystems, Foster City, CA). Reactions were run on an ABI377 and chromatograms were analyzed using Sequencher 4.6 (Gene Codes Corporation, Ann Arbor, MI).

***Cryptosporidium* and *Giardia* species:** On December 17, 2006, water samples for the enumeration of *Cryptosporidium* oocysts and *Giardia* cysts were collected from both the outfall of the 17th St. Canal at Lake Pontchartrain, and from the 17th St. Canal at the upstream (i.e. city) side of New Orleans Pump Station #6 (i.e. the Metairie Pump Station), while active pumping of water from the city interior was in progress. *Cryptosporidium* oocysts and *Giardia* cysts were harvested from the water and enumerated according to EPA Method 1623, using the Filta-Max™ filtration and elution system (IDEXX, Westbrook, ME). Filter processing, immuno-magnetic separation, and immuno-

fluorescent antibody staining (IFA) with subsequent fluorescent microscopy analysis of oocysts and cysts by EPA Method 1623 (17) was conducted by BCS Laboratories, Inc (Miami, FL), a NELAC-certified laboratory. Spiked control samples for each sample location were prepared as per EPA Method 1623 with EasySeed™ *Cryptosporidium* and *Giardia* control (BTF Precise Microbiology, Inc., Pittsburgh, PA). Sample volumes filtered for the canal outfall site were 196.30 liters and 146.90 liters for the natural sample and spiked control sample, respectively. Sample volumes filtered for the 17th St. Canal at Pump Station #6 were 110.35 liters and 71.74 liters for the natural sample and spiked control, respectively.

Soil samples (200g each) from sites Y2 and Y3 were again collected on this date in the same manner as described in the “Sample Collection” section above. Enumeration of *Cryptosporidium* oocysts and *Giardia* cysts in soil samples was conducted commercially by BCS Laboratories, Inc. There is currently no EPA standard method for *Cryptosporidium* and *Giardia* analysis in soils. Oocysts and cysts were extracted from 25g sub-samples of soil using the NaCl flotation method described by Kuczynska and Shelton (18). Soil sub-samples (25g) were dispersed in 100mL of Tris-Tween 80 dispersing solution (50mM Tris and 0.5% [vol/vol] Tween 80) with magnetic stirring for 15 min, then sieved, washed, centrifuged, and finally the oocysts and cysts harvested by flotation in NaCl (S.G. = 1.21) as previously described (18). After final resuspension of oocyst/cyst pellets in 100uL of distilled water, the (oo)cysts were then stained by IFA and enumerated by fluorescent microscopy as described above for EPA Method 1623.

Microbial Assemblage Analysis: PCR amplifications employed universal primers targeting the V4 –V8 region of the SSU rRNA gene of eukaryotes, archaea and bacteria corresponding to *E. coli* reference positions 517 through 1391: 517F (5' GCCAGCAGCCGCGGTAA3') and 1391R (5'-GACGGGCGGTGTGTRCA-3'). PCR reactions were carried out using an Eppendorf Mastercycler under the following conditions: 35 cycles at 94°C for 1 min, 50°C for 1 min, 72°C for 2 min; 72°C for 10 min. An Invitrogen Purelink Purification kit (Invitrogen, Carlsbad, CA) was used to purify PCR products for cloning and exclude any fragments less than 300 base pairs. An A-tailing reaction and an additional purification with the Purelink kit preceded cloning using the Invitrogen TOPO TA kit.

A RevPrep Orbit robotic template preparation instrument served to isolate and purify plasmid DNA (Genomic Solutions, Ann Arbor, MI). Sequencing of plasmid DNA was carried out using an ABI 3730 (Applied Biosystems, Foster City, CA) sequencing machine. A total of 192 clones were sequenced from each library in the forward and reverse directions using M13F (5' GTAAAACGACGGCCAGT 3') and M13R (5' AACAGCTATGACCATG 3') primers. We subjected individual sequence reads to quality control, vector trimming and assembly using phred, crossmatch and phrap software (19,20,21). We then ran BLAST searches against the nucleotide and environmental sequence databases to identify the closest relatives of the sequences observed to those in GenBank. Copies of the BLAST report summaries are available on the Woods Hole Centers for Oceans and Human Health website (<http://www.whoi.edu/science/cohh/whcohh/>). Using the BLAST results, we separated

out bacterial sequences from archaeal and eukaryotic sequences and further subjected these sequences to the Ribosomal Database Project (RDP) II's Classifier (22) to obtain phylogenetic information about our environmental sequence data. The final total numbers of bacterial clones varied between libraries. The data shown in pie charts (Fig. 4) represents an overview of the bacterial diversity with reference to the associated environments of the study sites from which top BLAST hits were obtained. We find this approach useful in making broad scale comparisons between clone libraries derived from different sites and different dates. The phylogenetic affinity of the potential pathogens or pathogen related sequences is derived from taxonomic information obtained from the RDP classifier using the default settings.

Chlorophyll-a Analysis: Water samples (200 ml) were filtered through GF/F glass fiber filters, which were then frozen at -80°C until extracted. Filters were extracted in two steps, first for 30 min with 10mL of dimethyl sulfoxide at 5°C, and then overnight with an additional 15mL of 90% acetone at -20°C. Extracts were measured fluorometrically before and after acidification for measurement of chlorophyll and phaeopigment concentration using a fluorometer (Turner Designs Model 10-000R, Sunnyvale, CA) equipped with an infrared-sensitive photomultiplier and calibrated using pure chlorophyll-*a*.

Supplemental Table S-1: GPS Coordinates of Sampling Sites

Site ID	latitude	longitude	sample site description
A1	30.0249	-90.1228	Lake Pontchartrain
A2	30.0247	-90.1336	Lake Pontchartrain
A3	30.0295	-90.1542	Lake Pontchartrain
A4	30.0329	-90.1743	Lake Pontchartrain
B1	30.0264	-90.1210	Lake Pontchartrain
B2	30.0355	-90.1209	Lake Pontchartrain
B3	30.0543	-90.1210	Lake Pontchartrain
B4	30.0719	-90.1209	Lake Pontchartrain
C1	30.0280	-90.1128	Lake Pontchartrain
C2	30.0319	-90.1038	Lake Pontchartrain
C3	30.0410	-90.0857	Lake Pontchartrain
C4	30.0500	-90.0681	Lake Pontchartrain
CS	30.0688	-90.1636	Lake Pontchartrain
D1	30.0323	-90.0771	Lake Pontchartrain
D2	30.0321	-90.0745	Lake Pontchartrain
D3	30.0412	-90.0753	Lake Pontchartrain
D4	30.0592	-90.0766	Lake Pontchartrain
E1	30.0340	-90.0377	Lake Pontchartrain
E2	30.0334	-90.0352	Lake Pontchartrain
E3	30.0422	-90.0374	Lake Pontchartrain
E4	30.0600	-90.0413	Lake Pontchartrain
R1	30.0255	-90.1203	Industrial Canal
R2	30.0257	-90.0829	Bayou St. John
R3	30.0296	-90.0741	London Ave. Canal
R4	30.0316	-90.0380	Industrial Canal
R5	30.0115	-90.0701	London Ave. Canal
R6	30.0119	-90.0860	Bayou St. John
S1	29.9868	-90.1251	17th Street Canal
S3	30.0250	-90.0316	Industrial Canal
Y1	30.0183	-90.0611	Flooded sediment
Y2	30.0151	-90.0694	Flooded sediment
Y3	30.0236	-90.0204	Flooded sediment
Y4	30.0089	-90.1151	Flooded sediment
Y5	30.0158	-90.1025	Flooded sediment
Y6	30.0338	-89.9635	Flooded sediment
Y7	30.0568	-89.9459	Flooded sediment
Y8	29.9905	-90.0475	Flooded sediment
Y9	29.9724	-90.0181	Flooded sediment
Y10	29.9640	-89.9872	Flooded sediment
Y11	29.9477	-90.1041	Flooded sediment
Z1	29.9572	-90.0630	non-flooded sediment
Z2	29.9360	-90.0741	non-flooded sediment
Z3	29.9233	-90.0923	non-flooded sediment
Z4	29.9259	-90.1276	non-flooded sediment
Z5	29.9911	-90.1301	non-flooded sediment
Z6	29.9547	-90.1661	non-flooded sediment
Z7	29.9988	-90.2139	non-flooded sediment
Z8	29.9893	-90.1718	non-flooded sediment
Z9	30.0044	-90.1406	non-flooded sediment
Z10	30.0127	-90.0098	non-flooded sediment

Supplemental Table S-2: Abundance of fecal indicators and total *Vibrio* and presence of *Bacteroidales*, *Bifidobacterium*, and *Legionella* spp. in Lake Pontchartrain waters. Units for enumerated samples include colony-forming units (CFU) for samples analyzed by membrane filtration and CFU equivalents for samples enumerated using qPCR. Samples enumerated using chromogenic substrate and coliphage culture method correspond to most probable numbers (MPN). *Bacteroidales*, *Bifidobacterium*, and *Legionella* spp. are reported as detected (+) or not detected (neg). “na” indicates no sample analyzed for corresponding microbe.

Table on following page.

Lake Sample Sites →		Transect # 1 – Number/100mL				Transect # 2 - Number/100mL				Lake Transect #3 – Number/100mL			
Assay	Date	A1	A2	A3	A4	B1	B2	B3	B4	C1	C2	C3	C4
Enterococci													
qPCR	11 Oct. 2005	21	<5	<5	<5	14	<5	<5	<5	6	<5	<5	9
MF ^a	11 Oct. 2005	8	<1	1	<1	28	2	<1	1.5	2	<1	<1	1
CS ^b	25 Oct. 2005	244	25	10	<1	340	20	26	5	80	10	5	5
qPCR	25 Oct. 2005	113	29	<5	<5	291	34	5	<5	61	<5	<5	<5
MF	25 Oct. 2005	38	1	1	5	37	3	1	2	12	6	4	3
CS	8 Nov. 2005	50	21	<1	5	60	5	5	1	31	10	<1	5
qPCR	8 Nov. 2005	36	11	<5	<5	105	8	<5	<5	19	<5	<5	<5
MF	8 Nov. 2005	12	4	3	10	50	3	1	1	13	19	1	<1
<i>E. coli</i>													
qPCR	25 Oct. 2005	297	30	18	neg	120	57	<5	10.5	12	37	<5	21
MF	25 Oct. 2005	150	10	<1	<1	200	15	<1	10	20	50	35	15
qPCR	8 Nov. 2005	227	75	59	50	71	<5	13	<5	159	77	39	16
MF	23 Nov. 2005	70	<1	5	2	95	13	7	10	28	4	4	3
FRNA coliphages													
culture	25 Oct. 2005	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
culture	8 Nov. 2005	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1	<1
<i>Clostridium perfringens</i>													
culture	25 Oct. 2005	16	4	4	<4	<4	<4	<4	<4	<4	<4	<4	<4
culture	8 Nov. 2005	na	<4	<4	<4	<4	<4	<4	<4	<4	<4	<4	<4
culture	12 Dec. 2005	40	34	<4	<4	4	<4	<4	<4	<4	<4	<4	<4
culture	27 Jan. 2006	<4	4	<4	4	4	8	<4	<4	<4	<4	<4	<4
Bacteroidales													
TNA ^c	25 Oct. 2005	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg
PCR	25 Oct. 2005	+	+	neg	neg	+	+	neg	neg	+	+	neg	neg
TNA	8 Nov. 2005	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg
PCR	8 Nov. 2005	+	+	neg	neg	+	+	neg	neg	+	+	neg	neg
<i>Bifidobacterium</i>													
PCR	25 Oct. 2005	+	+	neg	neg	+	+	neg	neg	+	neg	neg	neg
PCR	8 Nov. 2005	+	+	neg	neg	+	+	neg	neg	neg	neg	neg	neg
Total <i>Vibrio</i>													
MF	11 Oct. 2005	2690	532	524	352	2350	776	148	128	756	532	612	196
MF	27 Jan. 2006	676	na	na	na	412	408	36	na	na	na	na	na
<i>Legionella</i>													
genus PCR	11 Oct. 2005	+	+	+	+	+	+	+	neg	+	+	+	+
<i>mip</i> PCR	11 Oct. 2005	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg
genus PCR	25 Oct. 2005	+	+	neg	+	+	+	+	+	+	+	+	neg
<i>mip</i> PCR	25 Oct. 2005	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg
genus PCR	8 Nov. 2005	+	neg	neg	+	neg	+	+	+	neg	neg	neg	neg
<i>mip</i> PCR	8 Nov. 2005	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg	neg
genus PCR	12 Dec. 2005	na	na	na	na	+	+	neg	+	na	na	na	na
<i>mip</i> PCR	12 Dec. 2005	na	na	na	na	neg	+	neg	neg	na	na	na	na
genus PCR	27 Jan. 2006	na	na	na	na	+	neg	neg	+	na	na	na	na
<i>mip</i> PCR	27 Jan. 2006	na	na	na	na	neg	neg	neg	neg	na	na	na	na
genus PCR	29 Mar. 2006	na	na	na	na	+	+	+	+	na	na	na	na
<i>mip</i> PCR	29 Mar. 2006	na	na	na	na	neg	neg	neg	neg	na	na	na	na

^aMF = membrane filtration; ^bCS = chromogenic substrate; ^cTNA = *Taq* nuclease assay

Supplemental Table S-3: Abundance of fecal indicators and total *Vibrio* and presence of *Bacteroidales*, *Bifidobacterium*, and *Legionella* spp. in Canal waters.

Units for enumerated samples include colony-forming units (CFU) for samples analyzed by plate counts and CFU equivalents for samples enumerated by qPCR. Samples enumerated using chromogenic substrate and coliphage culture method correspond to most probable numbers (MPN). *Bacteroidales*, *Bifidobacterium*, and *Legionella* spp. are reported as detected (+) or not detected (neg). “na” indicates no sample analyzed for corresponding microbe.

Table on following page.

Panel A: Water from New Orleans Canals Discharging into Lake Pontchartrain - CFU/100mL

Date → Site/ Indicator	Assay	11 Oct. 2005	19 Oct. 2005	24 Oct. 2005	2 Nov. 2005	12 Nov. 2005	
						Before Pumping	After pumping
Enterococci							
17 th Street Canal (S1)	CS ^a	na	266	107	63	115	442
	qPCR	185	537	217	127	139	1480
	MF ^b	255	202	93	115	26	2150
Industrial Canal (S3)	CS	na	15	15	5	15	na
	qPCR	16	<5	5.7	<5	12	na
	MF	24	2	6	2	4	na
<i>E. coli</i>							
17 th Street Canal (S1)	qPCR	1,760	2,700	2,000	3,110	383	3,570
	MF	6,000	11,700	13,000	3,600	300	7,500
Industrial Canal (S3)	qPCR	85	176	623	418	10	na
	MF	30	46	10	20	10	na
FRNA coliphages							
17 th Street Canal (S1)	culture	na	na	8	na	na	na
Industrial Canal (S3)	culture	na	na	<1	na	na	na
<i>Clostridium perfringens</i>							
17 th Street Canal (S1)	culture	na	na	88	<4	na	na
Industrial Canal (S3)	culture	na	na	<4	<4	na	na
Bacteroidales							
17 th Street Canal (S1)	TNA ^c	neg	neg	neg	neg	neg	neg
	PCR	+	na	+	na	na	na
Industrial Canal (S3)	TNA	neg	neg	neg	neg	neg	neg
	PCR	+	na	neg	na	na	na
<i>Bifidobacterium</i>							
17 th Street Canal (S1)	PCR	na	+	+	+	+	+
Industrial Canal (S3)	PCR	na	+	+	+	+	na
Total <i>Vibrio</i>							
17 th Street Canal (S1)	MF	5,580	na	na	na	na	na
Industrial Canal (S3)	MF	1,780	na	na	na	na	na
<i>Legionella</i> spp.							
17 th Street Canal (S1)	Genus PCR	+	+	+	+	na	na
Industrial Canal (S3)	Genus PCR	+	+	neg	+	na	na
<i>L. pneumophila</i>							
17 th Street Canal (S1)	<i>mip</i> gene	neg	neg	neg	neg	na	na
Industrial Canal (S3)	<i>mip</i> gene	neg	neg	neg	neg	na	na

Panel B: Water from New Orleans Canal Shoreline – CFU/100mL

Site ID	12 Nov. 2005 – Indicator & Assay – CFU/100mL			
	Enterococci, CS	Enterococci, qPCR	<i>E. coli</i> , qPCR	<i>Bifidobacterium</i> PCR
Industrial Canal (R1)	42	17	350	+
Bayou St. John (R2)	387	345	9,610	-
London Ave. Canal (R3)	282	30	730	+
Industrial Canal (R4)	158	15	50	-
London Ave. Canal (R5)	193	208	332	+
Bayou St. John (R6)	152	53	130	+

^aCS = chromogenic substrate; ^bMF = membrane filtration; ^cTNA = *Taq* nuclease assay

Supplemental Table S-4: Vibrio abundance and percentage of pathogenic Vibrio isolates. Total colony counts on TCBS medium (CFU per 100 ml) and % of all purified isolates from either TCBS or CHROMagar® Vibrio media identified as *V. cholerae*, *V. vulnificus*, and *V. parahaemolyticus*, or other species for water samples collected from Lake Pontchartrain and from the 17th Street and Industrial Canals.

Panel A: 11 Oct 2005						
Station	Total Vibrio CFU/100mL	Number of Isolates	Positive Isolates (%)			
			<i>V. cholerae</i>	<i>V. vulnificus</i>	<i>V. parahaemolyticus</i>	Other
Lake, A1	2690	23	30	39	4	26
Lake, A2	532	11	18	55	0	27
Lake, A3	524	13	8	62	0	31
Lake, A4	352	25	16	36	4	44
Lake, B1	2350	15	13	60	7	20
Lake, B2	776	19	5	74	0	21
Lake, B3	148	11	36	27	9	27
Lake, B4	128	42	24	24	0	52
Lake, C1	756	7	29	57	0	14
Lake, C2	532	13	23	54	0	23
Lake, C3	612	5	0	40	0	60
Lake, C4	196	22	5	41	0	55
17 th St. Canal, S1	5580	29	79	3	0	17
Industrial Canal, S3	1780	29	10	72	3	14
Average	1150	19	21	46	1.8	31

Panel B: 27 Jan 2006						
Station	Total Vibrio CFU/100mL	Number of Isolates	Positive Isolates (%)			
			<i>V. cholerae</i>	<i>V. vulnificus</i>	<i>V. parahaemolyticus</i>	Other
Lake, A1	676	19	0	0	0	100
Lake, B1	412	13	0	0	0	100
Lake, B2	408	11	0	0	0	100
Lake, B3	36	12	8	0	0	92
Lake, D1	472	12	0	0	0	100
Lake, D2	596	15	0	0	0	100
Lake, D3	44	12	0	0	17	83
Lake, D4	32	12	8	0	0	92
Lake, E1	332	13	0	0	0	100
Lake, E2	268	14	0	0	0	100
Lake, E3	172	12	0	0	0	100
Lake, E4	60	12	0	0	0	100
17 th St. Canal, S1	288	32	0	0	0	100
Industrial Canal, S3	664	19	11	0	0	89
Average	298	14	1.8	0	1.1	97

Supplemental Table S-5: Abundance of Fecal Indicators in Shoreline Sediments of Lake Pontchartrain and New Orleans Canals, and in flooded and non-flooded residential sediments of the New Orleans area. All samples were enumerated as number per gram of dry sediment, except for *Bifidobacterium* which was reported as detected (+) or not detected (-). Results from % organic analysis of the sediments provided for reference.

Site ID ^b	12 Nov 05				25 Mar 06			20 Jun 06
	<i>E. coli</i> ^a	enterococci ^a		% organic	enterococci	<i>Bifidobacterium</i>	% organic	enterococci
	qPCR	qPCR	CS ^c		CS			CS
R1	39	26	25	7.3	20	+	1.9	27
R2	774	969	134	4.8	132	+	3.1	207
R3	69	200	489	5.6	166	+	7.1	316
R4	0.7	4	10	0.2	15	-	0.2	8
R5	167	870	467	4.6	240	+	7.3	680
R6	102	300	194	2.8	581	+	12	237
Y1	471	468		12.9	360	+	13	224
Y2	271	666		10.1	240	+	7.8	1,860
Y3	975	585		13.6	1,040	+	12	303
Y4								57
Y5								518
Y6								7
Y7								6,220
Y8								139
Y9								5
Y10								162
Y11								274
Z1								835
Z2								18
Z3								154
Z4								391
Z5								1
Z6								7,590
Z7								28
Z8								378
Z9								130
Z10								19

^aUnits for *E. coli* and enterococci by qPCR are in CFU equivalents per g dry sediment. Enterococci by CS in units of MPN per gram of dry sediment

^bSample types include shoreline sediment for samples which include an “R” in the name, flooded residential soils correspond to “Y”, and non-flooded residential soils correspond to “Z”.

^cCS = chromogenic substrate

References for On-Line Supplemental Information:

1. U.S. Army Corps of Engineers, Saint Louis District (2003) Raw LIDAR Elevation Data, New Orleans East quadrangle (and other quadrangles), Louisiana, UTM 15 NAD83, in *Louisiana Federal Emergency Management Agency Project - Phase I of Louisiana LIDAR Data Development: Lower Mississippi-Lafourche Parish, Louisiana* (U. S. Army Corps of Engineers, Saint Louis District contract number DACW43-00D-0511 0014, Task Order 05)
2. U. S. Environmental Protection Agency (2002) *Method 1600: Enterococci in Water by Membrane Filtration Using membrane-Enterococcus Indoxyl-beta-D-Glucoside Agar (mEI)* (EPA Publication 821-R-02-022).
3. U. S. Environmental Protection Agency (2002) *Method 1103.1: Escherichia coli (E. coli) in Water by Membrane Filtration Using membrane-Thermotolerant Escherichia coli Agar (mTEC)* (EPA Publication 821-R-02-020).
4. U.S. Environmental Protection Agency (1996) *ICR Microbial Laboratory Manual* (EPA Publication 600-R-95-178). US EPA Office of Water, Washington D.C.
5. Sobsey MD, Yates MV, Hsu FC, Lovelace G, Battigelli D, Margolin A, Pillai SD, Nwachuku N (2004). Development and evaluation of methods to detect coliphages in large volumes of water. *Wat. Sci Technol.* 50: 211-217.

6. Dick LK, Field KG (2004). Rapid estimation of numbers of fecal *Bacteroidetes* by use of a quantitative PCR assay for 16S rRNA genes. *Appl. Environ. Microbiol.* 70: 5695-5697.
7. King EL, Bachoon DS, Gates KW (2007). Rapid Detection of Human Fecal Contamination in Estuarine Environments by PCR targeting of *Bifidobacterium adolescentis*. *Journal of Microbial Methods*, 68(1): 76-81.
8. Bernhard AE, Field KG (2000) Identification of nonpoint sources of fecal pollution in coastal waters by using host-specific 16S ribosomal DNA genetic markers from fecal anaerobes. *Appl. Environ. Microbiol.* 66: 1587-1594.
9. Azanza PV, Buckle KA, Fleet GH (1996) Effect of diluents on the enumeration of *Vibrio vulnificus*. *Food Microbiology* 30: 385-390 (1996).
10. Chun J, Huq A, Colwell RR (1999). Analysis of 16S rRNA intergenic spacer regions of *Vibrio cholerae* and *Vibrio mimicus*. *Appl. Environ. Microbiol.* 65: 2202-2208.
11. Bej AK *et al.* (1999) Detection of total and hemolysin-producing *Vibrio parahaemolyticus* in shellfish using multiplex PCR amplification of *tl*, *tdh*, and *trh*. *J. Microbial Methods* 36: 215-225.
12. Brasher CW, De Paola A., Jones DD, Bej AK (1998). Detection of microbial pathogens in shellfish with multiplex PCR. *Current Microbiology* 37: 101-107.

13. Lane DJ (1991) 16S/23S rRNA sequencing. in *Nucleic Acid Techniques in Bacterial Systematics*, eds Stackebrandt E, Goodfellow M. (Wiley, New York) pp. 115-175.
14. Jonas D, Rosenbaum A, Weyrich S, Bhakdi S (1995) Enzyme-linked immunoassay for detection of PCR-amplified DNA of legionellae in brochoalveolar fluid. *Journal of Clinical Microbiology* 33: 1247-1252.
15. Miyamoto H, Yamamoto H, Arima K, Fujii J, Maruta K, Izu K (1997) Development of a new seminested PCR method for detection of *Legionella* species and its application to surveillance of legionellae in hospital cooling tower water. *Applied and Environmental Microbiology* 63: 2489-2494.
16. Templeton KE, Scheltinga SA, Sillekens P, Crielaard JW, van Dam AP, Goossens H, Claas ECJ (2003) Development and clinical evaluation of an internally controlled, single-tube multiplex real-time PCR assay for detection of *Legionella pneumophila* and other *Legionella* species. *Journal of Clinical Microbiology* 41: 4016-4021.
17. U. S. Environmental Protection Agency (2001) *Method 1623: Cryptosporidium and Giardia in Water by Filtration/IMS/FA* (EPA Publication 821-R-01-025).

18. Kuczynska E, Shelton DR (1999) Method for detection and enumeration of *Cryptosporidium parvum* oocysts in feces, manures, and soils. *Applied and Environmental Microbiology* 65: 2820-2826.
19. Gordon D, Abajian C, Green P (1998) CONSED: a graphical tool for sequence finishing. *Genome Research* 8: 195.
20. Ewing B, Hillier L, Wendl MC, Green P (1998) Base-calling of automated sequencer traces using phred. I. Accuracy assessment. *Genome Research* 8: 175.
21. Ewing B, Green P (1998) Base-calling of automated sequencer traces using phred. II. Error probabilities. *Genome Research* 8: 186.
22. Cole JR, Chai B, Farris RJ, Wang Q, Kulam SA, McGarrell DM, Garrity GM, Tiedje JM (2005) The ribosomal database project (RDP-11): sequences and tools for high throughput rRNA analysis. *Nucleic Acids Res* 33: D294-6.