

Oxygen Isotope data from 37 R/V Albatross IV, R/V Endeavor, R/V Oceanus, and R/V Parizeau (Canadian) cruises in the Gulf of Maine and Georges Bank, the Gulf of St. Lawrence and Scotian Shelf from 1994-1999 (GB project)

Website: <https://www.bco-dmo.org/dataset/2397>

Data Type: Cruise Results

Version: 1

Version Date: 2004-11-15

Project

» [U.S. GLOBEC Georges Bank](#) (GB)

Program

» [U.S. GLOBal ocean ECosystems dynamics](#) (U.S. GLOBEC)

Contributors	Affiliation	Role
Houghton, Robert W	Lamont-Doherty Earth Observatory (LDEO)	Principal Investigator
Allison, Dicky	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Oxygen Isotope data from 37 R/V Albatross IV, R/V Endeavor, R/V Oceanus, and R/V Parizeau (Canadian) cruises in the Gulf of Maine and Georges Bank, the Gulf of St. Lawrence and Scotian Shelf from 1994-1999 (GB project)

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Program Information](#)
- [Funding](#)

Coverage

Spatial Extent: N:50.08 E:-55.83 S:40.45 W:-69.28

Temporal Extent: 1994-05-31 - 1999-06-24

Dataset Description

Oxygen Isotope data, GLOBEC and Canadian Cruises 1994-1998

Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio $H_2^{18}O/H_2^{16}O$ is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO, and presented in delta¹⁸O notation where:

- Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033.
- The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file.
- Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography.
- The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.

Contributors:

Bob Houghton and Rick Fairbanks

Any questions, contact:

Bob Houghton

Lamont-Doherty Earth Observatory of Columbia University

Palisades, NY 10964

voice: 845 365 8328

fax: 845 365 8157

email: houghton@ldeo.columbia.edu

Updated: November 15, 2004; gfh

Acquisition Description

Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio $H^{218}O/H^{216}O$ is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.

Processing Description

Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033.

The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file.

Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography.

The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
month_begin	starting month of cruise	
day_begin	starting day of cruise	
month_end	ending month of cruise	
day_end	ending day of cruise	
si	name of contributing scientist(s)	
location	sampling program/area	
event	event number, combination of cruiseid and cast number	
station_std	standard station number	
cast	CTD/bottle cast number	
bottle	bottle number/identification	
lat	latitude, negative = South	decimal degrees
lon	longitude, negative = West	decimal degrees
delta_O_18	isotope ratio H ²¹⁸ O/H ²¹⁶ O, presented in delta18O notation	
temp	temperature	degrees centigrade
sal	salinity, PSS	
trans_v	light transmission	volts
cond_mM	conductivity	milliMhos/centimeter
depth_w	water depth	meters
depth	depth of sample	meters
diff_delta_O_18	difference range between replicates	
comments	identifies repeated analysis	
cruiseid	cruise identification	
year	Year of cruise, 4 digit year	
flvolt	fluorescence, from CTD mounted fluorometer	volts

Instruments

Dataset-specific Instrument Name	CTD Seabird 911
Generic Instrument Name	CTD Sea-Bird 911
Dataset-specific Description	CTD measurements taken by a SBE911 (SeaBird) CTD instrument package
Generic Instrument Description	<p>The Sea-Bird SBE 911 is a type of CTD instrument package. The SBE 911 includes the SBE 9 Underwater Unit and the SBE 11 Deck Unit (for real-time readout using conductive wire) for deployment from a vessel. The combination of the SBE 9 and SBE 11 is called a SBE 911. The SBE 9 uses Sea-Bird's standard modular temperature and conductivity sensors (SBE 3 and SBE 4). The SBE 9 CTD can be configured with auxiliary sensors to measure other parameters including dissolved oxygen, pH, turbidity, fluorescence, light (PAR), light transmission, etc.). More information from Sea-Bird Electronics.</p>

Dataset-specific Instrument Name	Mark 5 CTD
Generic Instrument Name	CTD Neil Brown Mark 5
Dataset-specific Description	Mark 5 CTD.
Generic Instrument Description	The Neil Brown Instrument Systems Mark 5 CTD is used to measure conductivity, temperature, and depth of sea water. The MK5 profiler has a higher sampling rate than the SeaBird SEACAT. (For the GLOBEC Georges Bank project the Mark 5 was instrumented with an expanded suite of sensors and deployed almost exclusively at GLOBEC Standard stations.)

Dataset-specific Instrument Name	VG Prism II Mass-Spectrometer
Generic Instrument Name	Isotope-ratio Mass Spectrometer
Dataset-specific Description	The isotope ratio H ₂ ¹⁸ O/H ₂ ¹⁶ O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.
Generic Instrument Description	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

[[table of contents](#) | [back to top](#)]

Deployments

Website	https://www.bco-dmo.org/deployment/57369
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9404/AL9404.htm
Start Date	1994-05-31
End Date	1994-06-10
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9505

Website	https://www.bco-dmo.org/deployment/57371
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9505/al9505rot.pdf
Start Date	1995-05-09
End Date	1995-05-18
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9506

Website	https://www.bco-dmo.org/deployment/57372
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9506/al9506new.html
Start Date	1995-06-05
End Date	1995-06-15
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9508

Website	https://www.bco-dmo.org/deployment/57373
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9508/a9508rp2.HTM
Start Date	1995-07-10
End Date	1995-07-20
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9605

Website	https://www.bco-dmo.org/deployment/57375
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9605/al9605.html
Start Date	1996-05-06
End Date	1996-05-17
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9607

Website	https://www.bco-dmo.org/deployment/57376
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9607/AL9607.pdf
Start Date	1996-06-03
End Date	1996-06-13
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9701

Website	https://www.bco-dmo.org/deployment/57378
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9701/cral9701.htm
Start Date	1997-01-13
End Date	1997-01-20
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9705

Website	https://www.bco-dmo.org/deployment/57379
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9705/al9705.html
Start Date	1997-05-19
End Date	1997-05-27
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9707

Website	https://www.bco-dmo.org/deployment/57380
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9707/al9707.html
Start Date	1997-06-18
End Date	1997-06-28
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

Website	https://www.bco-dmo.org/deployment/57401
Platform	R/V Endeavor
Start Date	1995-02-10
End Date	1995-02-20
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

EN263

Website	https://www.bco-dmo.org/deployment/57403
Platform	R/V Endeavor
Report	http://globec.who.edu/globec-dir/reports/en263/EN263.pdf
Start Date	1995-03-13
End Date	1995-03-24
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

EN265

Website	https://www.bco-dmo.org/deployment/57405
Platform	R/V Endeavor
Start Date	1995-04-11
End Date	1995-04-22
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

EN276

Website	https://www.bco-dmo.org/deployment/57413
Platform	R/V Endeavor
Report	http://globec.who.edu/globec-dir/reports/en276/EN276.pdf
Start Date	1996-01-10
End Date	1996-01-22
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

EN278

Website	https://www.bco-dmo.org/deployment/57414
Platform	R/V Endeavor
Start Date	1996-02-13
End Date	1996-02-25
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

Website	https://www.bco-dmo.org/deployment/57415
Platform	R/V Endeavor
Start Date	1996-04-08
End Date	1996-04-20
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC275

Website	https://www.bco-dmo.org/deployment/57440
Platform	R/V Oceanus
Start Date	1996-03-11
End Date	1996-03-22
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC298

Website	https://www.bco-dmo.org/deployment/57444
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc298/cruisereport.html
Start Date	1997-02-11
End Date	1997-02-23
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC300

Website	https://www.bco-dmo.org/deployment/57446
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc300/oc300rpt.mr7.html
Start Date	1997-03-16
End Date	1997-03-28
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC302

Website	https://www.bco-dmo.org/deployment/57448
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc302/oce302.html
Start Date	1997-04-22
End Date	1997-05-02
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

PAR95-010

Website	https://www.bco-dmo.org/deployment/57473
Platform	R/V Parizeau
Report	http://globec.who.edu/globec-dir/reports/par95010/PAR95-010.pdf
Start Date	1995-06-06
End Date	1995-06-13
Description	<p>long term mooring turn-around</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

PAR95-034

Website	https://www.bco-dmo.org/deployment/57474
Platform	R/V Parizeau
Report	http://globec.who.edu/globec-dir/reports/par95034/PAR95034.pdf
Start Date	1995-11-24
End Date	1995-12-03
Description	<p>long term mooring turn-around</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

PAR96-024

Website	https://www.bco-dmo.org/deployment/57475
Platform	R/V Parizeau
Report	http://globec.who.edu/globec-dir/reports/par9624/par9624.htm
Start Date	1996-09-23
End Date	1996-09-30
Description	<p>long term mooring turn-around</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

PAR97-025

Website	https://www.bco-dmo.org/deployment/57476
Platform	R/V Parizeau
Report	http://globec.who.edu/globec-dir/reports/par9725/par9725.htm
Start Date	1997-06-27
End Date	1997-07-04
Description	<p>long term mooring</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9801

Website	https://www.bco-dmo.org/deployment/57382
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9801/al9801.html
Start Date	1998-01-07
End Date	1998-01-19
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC317

Website	https://www.bco-dmo.org/deployment/57451
Platform	R/V Oceanus
Start Date	1998-02-06
End Date	1998-02-19
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC319

Website	https://www.bco-dmo.org/deployment/57452
Platform	R/V Oceanus
Report	http://globec.whoi.edu/globec-dir/reports/oc319/oc319new/oc319rpt.8april98.htm
Start Date	1998-03-15
End Date	1998-03-27
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC322

Website	https://www.bco-dmo.org/deployment/57454
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc322/oc322.html
Start Date	1998-04-15
End Date	1998-04-27
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9806

Website	https://www.bco-dmo.org/deployment/57384
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9806/al9806.html
Start Date	1998-05-13
End Date	1998-05-22
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9808

Website	https://www.bco-dmo.org/deployment/57385
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9808/al9808.html
Start Date	1998-06-16
End Date	1998-06-26
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9901

Website	https://www.bco-dmo.org/deployment/57386
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9901/al9901.html
Start Date	1999-01-12
End Date	1999-01-24
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC336

Website	https://www.bco-dmo.org/deployment/57459
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc336/oc336cruise-report.html
Start Date	1999-02-11
End Date	1999-02-23
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

Website	https://www.bco-dmo.org/deployment/57427
Platform	R/V Endeavor
Report	http://globec.who.edu/globec-dir/reports/en320new/en320mda.htm
Start Date	1999-03-10
End Date	1999-03-23
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

OC341

Website	https://www.bco-dmo.org/deployment/57464
Platform	R/V Oceanus
Report	http://globec.who.edu/globec-dir/reports/oc341/reptoc341.html
Start Date	1999-04-16
End Date	1999-04-27
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9904

Website	https://www.bco-dmo.org/deployment/57387
Platform	R/V Albatross IV
Start Date	1999-05-19
End Date	1999-05-27
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H²¹⁸O/H²¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

AL9906

Website	https://www.bco-dmo.org/deployment/57388
Platform	R/V Albatross IV
Report	http://globec.who.edu/globec-dir/reports/al9906/al9906rpt.html
Start Date	1999-06-14
End Date	1999-06-24
Description	<p>broad-scale</p> <p>Acquisition Description Oxygen isotope data are derived from water samples collected on GLOBEC broad-scale and contemporaneous Canadian cruises 1994-1998. The isotope ratio H₂¹⁸O/H₂¹⁶O is measured, using a VG Prism II mass-spectrometer operated by Rick Mortlock and maintained by Rick Fairbanks at LDEO.</p> <p>Processing Description Replicates of each water sample are analyzed. The data presented is the average of the replicates and their difference. If the difference exceeds 0.066 the analysis is repeated. The average standard deviation of the data is 0.033. The content of the data file varies according to source, degree of editing, and some arbitrary formatting changes. For broad-scale data the cruise is always given in the file name and sometimes the header. The station number is the broad-scale standard station with associated lat., long., and bottom depth. Temperature, salinity, bottle depth, and sometimes fluorescence and transmissivity data are copied from the broad-scale hydrographic data file. Canadian data are from C.S.S. Parizeau cruises. The station and bottle number are unique to a particular cruise. When depth is listed as pressure it is numerically equivalent to depth in meters. Hydrographic data are supplied by Gary Bugden at the Bedford Institute of Oceanography. The time for each file is denoted by month and year. More precise day and hour information is available in the original hydrographic file.</p>

PAR95-031

Website	https://www.bco-dmo.org/deployment/57957
Platform	R/V Parizeau
Start Date	1995-10-20
End Date	1995-11-09
Description	Canadian Cruise associated with the GLOBEC Georges Bank program, although not part of it.

PAR95-032

Website	https://www.bco-dmo.org/deployment/57958
Platform	R/V Parizeau
Start Date	1995-11-14
End Date	1995-11-23
Description	Canadian cruise associated with GLOBEC Georges Bank, but not part of it. Cruise dates are actually the start and end dates of the dataset.

[[table of contents](#) | [back to top](#)]

Project Information

U.S. GLOBEC Georges Bank (GB)

Website: http://globec.whoi.edu/globec_program.html

Coverage: Georges Bank, Gulf of Maine, Northwest Atlantic Ocean

The U.S. GLOBEC Georges Bank Program is a large multi-disciplinary multi-year oceanographic effort. The proximate goal is to understand the population dynamics of key species on the Bank - Cod, Haddock, and two species of zooplankton (*Calanus finmarchicus* and *Pseudocalanus*) - in terms of their coupling to the physical environment and in terms of their predators and prey. The ultimate goal is to be able to predict changes in the distribution and abundance of these species as a result of changes in their physical and biotic environment as well as to anticipate how their populations might respond to climate change. The effort is substantial, requiring broad-scale surveys of the entire Bank, and process studies which focus both on the links between the target species and their physical environment, and the determination of fundamental aspects of these species' life history (birth rates, growth rates, death rates, etc). Equally important are the modelling efforts that are ongoing which seek to provide realistic predictions of the flow field and which utilize the life history information to produce an integrated view of the dynamics of the populations. The U.S. GLOBEC Georges Bank Executive Committee (EXCO) provides program leadership and effective communication with the funding agencies.

[[table of contents](#) | [back to top](#)]

Program Information

U.S. GLOBal ocean ECosystems dynamics (U.S. GLOBEC)

Website: <http://www.usglobec.org/>

Coverage: Global

U.S. GLOBEC (GLOBal ocean ECosystems dynamics) is a research program organized by oceanographers and fisheries scientists to address the question of how global climate change may affect the abundance and production of animals in the sea. The U.S. GLOBEC Program currently had major research efforts underway in the Georges Bank / Northwest Atlantic Region, and the Northeast Pacific (with components in the California Current and in the Coastal Gulf of Alaska). U.S. GLOBEC was a major contributor to International GLOBEC efforts in the Southern Ocean and Western Antarctic Peninsula (WAP).

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
National Science Foundation (NSF)	unknown GB NSF
National Oceanic and Atmospheric Administration (NOAA)	unknown GB NOAA

[[table of contents](#) | [back to top](#)]