



Geochemistry, Geophysics, Geosystems

Supporting Information for

Emplacement and high-temperature evolution of gabbros of the 16.5 °N oceanic core complexes (Mid-Atlantic Ridge): insights into the compositional variability of the lower oceanic crust

Sanfilippo A.^{1,2*}, Dick H.J.B.², Marschall H.R.^{2,3}, Lissenberg C.J.⁴, Urann, B.²

1- Dipartimento di Scienze della Terra e dell'Ambiente - Università degli Studi di Pavia, Via Ferrata 1, Pavia, Italy

2-Department of Geology and Geophysics - Woods Hole Oceanographic Institution, Woods Hole, MA 02543, USA

3- Institut für Geowissenschaften, Goethe Universität Frankfurt, Altenhöferallee 1, 60438 Frankfurt am Main, Germany

4- School of Earth and Ocean Sciences, Cardiff University, Park Place, Cardiff CF10 3AT, UK

Table of contents *

Table S1- Geographic location and rock type for each dredge site

Table S2- Average major element olivine compositions (wt%)

Table S3- Average major element plagioclase compositions (wt%)

Table S4- Average major element clinopyroxene compositions (wt%)

Table S5- Average major element orthopyroxene compositions (wt%)

Table S6- Major element amphibole compositions (wt%)

Table S7- Mass and proportions of liquid and solid crystallized during a fractional crystallization process

Table S8- Mass and proportions of liquid and solid crystallized during the model at different mass crystallized/mass erupted ratios (Mc/Me)

Table S9- Composition of trapped melt at different mass crystallized/mass erupted ratios (Mc/Me)

**Supporting tables are uploaded separately*