

Denitrification and DNRA data from Little Lagoon, Alabama collected from 2012-2013

Website: <https://www.bco-dmo.org/dataset/723966>

Data Type: Other Field Results

Version: 1

Version Date: 2018-01-16

Project

» [Groundwater Discharge, Benthic Coupling and Microalgal Community Structure in a Shallow Coastal Lagoon](#) (LittleLagoonGroundwater)

Contributors	Affiliation	Role
Mortazavi, Behzad	National Science Foundation (NSF-DEB)	Principal Investigator, Contact
Burnett, William C.	Florida State University (FSU - EOAS)	Co-Principal Investigator
Ake, Hannah	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Denitrification and DNRA data from Little Lagoon, Alabama collected from 2012-2013

Table of Contents

- [Coverage](#)
- [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
- [Related Publications](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

Coverage

Spatial Extent: Lat:30.241929 Lon:-87.773756

Temporal Extent: 2012 - 2013

Dataset Description

Denitrification and DNRA data from Little Lagoon, Alabama.

Acquisition Description

Little Lagoon is a shallow coastal lagoon that is tidally connected to the Gulf of Mexico but has no riverine inputs. The water in the lagoon is replenished solely from precipitation and groundwater inputs primarily on the East end (Su et al. 2012). Because of the rapid development in Baldwin County, a large amount of NO₃⁻ enters the Little Lagoon system through SGD (Murgulet & Tick 2008). In this region, there can be rapid changes in the depth to groundwater (Fig. 4.1 inset) and episodic SGD inputs to the lagoon (Su et al. 2013). Within the lagoon, three sites were selected (East, Mouth, and West) to represent the gradient that exists across the lagoon from the input of groundwater. Sites were sampled on a near-monthly basis from February 2012 to February 2013.

DNRA

Approximately 1 L of outflow water was collected from the inflow water and each core for DNRA analysis. Appropriate sample volume was determined after NH₄⁺ nutrient analysis and expected atom % enrichment. δ¹⁵N-NH₄⁺ was measured in samples, constructed blanks, and standards that bracketed the NH₄⁺ concentration of the samples following a modified ammonium diffusion procedure (Holmes et al. 1998) that collects NH₄⁺ dissolved in water by converting NH₄⁺ to NH₃ under basic conditions and then traps the NH₃ on an acidified glass fiber filter. Non diffused standards were prepared according to Stark and Hart (1996) to account for blank corrections. After ¹⁵N analysis on a Europa Scientific SL-2020 system (Stable Isotope Lab, Utah State University), DNRA was calculated from the production rate of ¹⁵NH₄⁺ (p¹⁵NH₄⁺) during the incubation according to Christensen et. al (2000): (7) where ρ is the production of ¹⁵N-NH₄⁺ and D₁₄ and D₁₅ are the denitrification rates of ¹⁴N-NO₃⁻ and ¹⁵N-NO₃⁻, respectively. This assumes that DNRA takes place in the same sediment layers as denitrification and that the ¹⁵N labeling of NO₃⁻ being reduced to NH₄⁺ equals the ¹⁵N labeling of NO₃⁻ being reduced to N₂

(Christensen et al. 2000).

Denitrification and anammox from slurry assays

Volumetric rates of denitrification, anammox, and the relative contribution of anammox to gross N₂ production were determined from sediment slurry incubations. Slurry rates for depth-integrated sediments (0-50 mm) were prepared in Exetainers (Thamdrup & Dalsgaard 2002) with artificial seawater (ASW) (70.2g NaCl, 3.0g KCl, 49.4 g MgSO₄*7H₂O, 5.8g CaCl₂*2H₂O L⁻¹) constructed at a salinity of 52 and diluted with deionized water to match the salinity of each site. After dilution, homogenized sediment from 0 to 50 mm was added to the ASW and the incubation bottle was sparged with N₂ and amended with 100 μmol L⁻¹ Na¹⁵NO₃ - (99 atom %). Sediment slurry was dispensed to 12 ml Exetainers, yielding approximately 1 ml of sediment and 11 ml ASW with no headspace. For each site, 12 vials total were incubated with three vials stopped at time points 0 to 36 h. Incubations were stopped by adding 250 μL of ZnCl₂ and resealing the vials without headspace.

Denitrification and anammox rates in slurries were calculated according to equations 5 and 6 described below.

Excess ²⁹N₂ and ³⁰N₂ concentrations for intact core and slurry incubations were calculated from dissolved ²⁹N₂:²⁸N₂ and ³⁰N₂:²⁸N₂ measured using a MIMS. Rates of excess ²⁹N₂ (p₂₉) and ³⁰N₂ (p₃₀) production were calculated from the flux calculation described above. Rates of ambient ¹⁴N₂ production (p₁₄) in core incubations with ¹⁵NO₃ - tracer addition were determined as (Nielsen 1992, Risgaard-Petersen et al. 2003):

$$(1) p_{14} = 2 \times r_{14} \pm [p_{29} + p_{30} \pm (1 - r_{16})]$$

The ¹⁴N:¹⁵N ratio of NO₃ - undergoing reduction to N₂ (r₁₄) was determined as follows:

$$(2) r_{14} = [R_{29} \times (1 - r_a) - r_a] \times (2 - r_a)^{-1}$$

where R₂₉ was the ratio of p₂₉ to p₃₀ determined for the cores and r_a was the relative contribution of anammox to gross N₂ production determined in vial slurry incubations. Gross denitrification and anammox rates within intact sediment cores with ¹⁵NO₃ - tracer addition were calculated as follows:

$$(3) \text{denitrification} = p_{14} \pm (1 - r_a)$$

$$(4) \text{anammox} = p_{14} \pm r_a$$

Denitrification stimulated by the added ¹⁵N-NO₃ - (D₁₅) was calculated from the classical IPT (Nielsen 1992) and these amended rates are a measure of the denitrification capacity under field conditions when NO₃ - is not limiting.

Rates of denitrification and anammox in vial slurry incubations with ¹⁵NO₃ - amendments

were calculated from the equations of Thamdrup and Dalsgaard (2002): (5) (6) where FN was the fraction of ^{15}N in NO_3^- . For months when anammox slurry incubations were not performed (August and November 2012), p14 is calculated as D14 from the IPT (Nielsen 1992). Potential denitrification and anammox rates were converted to an areal basis using the wet weight of the sediment in the slurry. All rates and fluxes pertaining to N species in this study were normalized to one atom N.

Additional methodology can be found in:

Bernard, Rebecca & Mortazavi, Behzad & A. Kleinhuizen, Alice. (2015). Dissimilatory nitrate reduction to ammonium (DNRA) seasonally dominates NO_3^- reduction pathways in an anthropogenically impacted sub-tropical coastal lagoon. *Biogeochemistry*. 125. 47-64. [10.1007/s10533-015-0111-6](https://doi.org/10.1007/s10533-015-0111-6).

Processing Description

Data were flagged as below detection limits if no measurable rates were returned after calculations. See equations in methodology section of:

Bernard, Rebecca & Mortazavi, Behzad & A. Kleinhuizen, Alice. (2015). Dissimilatory nitrate reduction to ammonium (DNRA) seasonally dominates NO₃⁻ reduction pathways in an anthropogenically impacted sub-tropical coastal lagoon. *Biogeochemistry*. 125. 47-64. [10.1007/s10533-015-0111-6](https://doi.org/10.1007/s10533-015-0111-6).

Statistical Analysis

To test the seasonal flux variability between sites in Little Lagoon, two-way ANOVAs with site and date as independent variables were performed. When data could not be transformed to meet ANOVA assumptions, Wilcoxon/Kruskal-Wallis nonparametric tests were used. When significant differences occurred, Tukey HSD or Steel-Dwass post hoc tests were used to determine significant interactions. A Principal component analysis (PCA) was conducted on all biogeochemical parameters to identify underlying multivariate components that may be influencing N fluxes. Spearman's rho correlation analysis was used to examine the relationship between the principal components and fluxes. Statistical significance of the data set was determined at $\alpha=0.05$ and error is reported as standard error. All statistical analyses were performed in SAS JMP 10 (SAS Institute Inc.).

BCO-DMO Data Processing Notes:

- Data reorganized into one table under one set of column names from both original files
- Units removed from column names
- Column names reformatted to meet BCO-DMO standards
- Information captured in original columns entered under column "Value_Description" where units are also described
- Created column Year to describe to capture the metadata in the file name

[[table of contents](#) | [back to top](#)]

Related Publications

Bernard, R. J., Mortazavi, B., & Kleinhuizen, A. A. (2015). Dissimilatory nitrate reduction to ammonium (DNRA) seasonally dominates NO₃ – reduction pathways in an anthropogenically impacted sub-tropical coastal lagoon. *Biogeochemistry*, 125(1), 47–64. doi:[10.1007/s10533-015-0111-6](https://doi.org/10.1007/s10533-015-0111-6)

Murgulet, D., & Tick, G. R. (2008). Assessing the extent and sources of nitrate contamination in the aquifer system of southern Baldwin County, Alabama. *Environmental Geology*, 58(5), 1051–1065. doi:[10.1007/s00254-008-1585-5](https://doi.org/10.1007/s00254-008-1585-5)

Nielsen, L. P. (1992). Denitrification in sediment determined from nitrogen isotope pairing. *FEMS Microbiology Letters*, 86(4), 357–362. doi:[10.1111/j.1574-6968.1992.tb04828.x](https://doi.org/10.1111/j.1574-6968.1992.tb04828.x)

Stark, J. M., & Hart, S. C. (1996). Diffusion Technique for Preparing Salt Solutions, Kjeldahl Digests, and Persulfate Digests for Nitrogen-15 Analysis. *Soil Science Society of America Journal*, 60(6), 1846. doi:[10.2136/sssaj1996.03615995006000060033x](https://doi.org/10.2136/sssaj1996.03615995006000060033x)

Su, N., Burnett, W.C., Eller, K.T., MacIntyre, H.L., Mortazavi, B., Leifer, J., Novoveska, L. (2012). Radon and radium isotopes, groundwater discharge and harmful algal blooms in Little Lagoon, Alabama. *Interdisciplinary Studies on Environmental Chemistry*, 6, 329–337.

Su, N., Burnett, W.C., MacIntyre, H.L., Liefer, J.D., Peterson, R.N., Viso, R. (2013). Natural radon and radium isotopes for assessing groundwater discharge into Little Lagoon, AL: implications for harmful algal blooms. *Estuaries Coasts*, 1–18

Thamdrup, B., & Dalsgaard, T. (2002). Production of N₂ through Anaerobic Ammonium Oxidation Coupled to Nitrate Reduction in Marine Sediments. *Applied and Environmental Microbiology*, 68(3), 1312–1318. doi:[10.1128/AEM.68.3.1312-1318.2002](https://doi.org/10.1128/AEM.68.3.1312-1318.2002)

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
Year	Year ID that samples were taken	unitless
Value_Description	Description of the measurement taken; description includes relevant units for each sample taken; Descriptions include: DIN:DIP = ratio of dissolved inorganic nitrogen to dissolved inorganic phosphate; Denitrification = Denitrification; p14 ambient denitrification = ambient denitrification rates; DNRA = dissimilatory nitrate reduction to ammonium; D15 denitrification = denitrification from added heavy labeled isotope.	unitless
Date	Month and day that samples were taken; MMM-DD	unitless

East	Denitrification and DNRA values collected at the East site; location of site is 30.253347, -87.724729	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹
East_SE	Standard error of denitrification and DNRA values collected at the East site	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹
Mouth	Denitrification and DNRA values collected at the Mouth site; location of site is 30.243683, -87.738407	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹
Mouth_SE	Standard error of denitrification and DNRA values collected at the Mouth site	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹

West	Denitrification and DNRA values collected at the West site; location of site is 30.247181, -87.767856	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹
West_SE	Standard error of denitrification and DNRA values collected at the West site	umol N m ⁻² hr ⁻¹ ; umol N m ⁻² d ⁻¹ ; mmol NH ₄ ⁺ m ⁻² d ⁻¹

[[table of contents](#) | [back to top](#)]

Instruments

Dataset-specific Instrument Name	Europa Scientific SL-2020 system
Generic Instrument Name	Isotope-ratio Mass Spectrometer
Dataset-specific Description	Used for ¹⁵ N analysis
Generic Instrument Description	The Isotope-ratio Mass Spectrometer is a particular type of mass spectrometer used to measure the relative abundance of isotopes in a given sample (e.g. VG Prism II Isotope Ratio Mass-Spectrometer).

Dataset-specific Instrument Name	Multichannel proportioning pump
Generic Instrument Name	Pump
Dataset-specific Description	Used to filter sediment
Generic Instrument Description	A pump is a device that moves fluids (liquids or gases), or sometimes slurries, by mechanical action. Pumps can be classified into three major groups according to the method they use to move the fluid: direct lift, displacement, and gravity pumps

Dataset-specific Instrument Name	MIMS
Generic Instrument Name	Membrane Inlet Mass Spectrometer
Dataset-specific Description	Used to measure dissolved gas
Generic Instrument Description	Membrane-introduction mass spectrometry (MIMS) is a method of introducing analytes into the mass spectrometer's vacuum chamber via a semipermeable membrane.

Dataset-specific Instrument Name	Continuous Flow Analyzer
Generic Instrument Name	Continuous Flow Analyzer
Dataset-specific Description	Used to measure continuous flow rate
Generic Instrument Description	A sample is injected into a flowing carrier solution passing rapidly through small-bore tubing.

[[table of contents](#) | [back to top](#)]

Deployments

LittleLagoon

Website	https://www.bco-dmo.org/deployment/528089
Platform	SmallBoat_FSU
Start Date	2010-04-05
End Date	2013-08-17
Description	The sampling sites were all accessed from small boats, here amalgamated to one deployment called LittleLagoon.

[[table of contents](#) | [back to top](#)]

Project Information

Groundwater Discharge, Benthic Coupling and Microalgal Community Structure in a Shallow Coastal Lagoon (LittleLagoonGroundwater)

Coverage: southern Alabama, east of Mobile

This project investigated the link between submarine groundwater discharge (SGD) and microalgal dynamics in Little Lagoon, Alabama. In contrast to most near-shore environments, it is fully accessible; has no riverine inputs; and is large enough to display ecological diversity (c. 14x 0.75 km) yet small enough to be comprehensively sampled on appropriate temporal and spatial scales. The PIs have previously demonstrated that the lagoon is a hot-spot for toxic blooms of the diatom *Pseudo-nitzschia* spp. that are correlated with discharge from the surficial aquifer. This project assessed variability in SGD, the dependence of benthic nutrient fluxes on microphytobenthos (MPB) abundance and productivity, and the response of the phytoplankton to nutrient enrichment and dilution. The work integrated multiple temporal and spatial scales and demonstrated both the relative importance of SGD vs. benthic recycling as a source of nutrients, and the role of SGD in structuring the microalgal community. (paraphrased from Award abstract)

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0962008

[[table of contents](#) | [back to top](#)]