

Rates of grazing by parrotfishes and macroalgal browsing in Fiji during 2010-2012 (Killer Seaweeds project)

Website: <https://www.bco-dmo.org/dataset/674125>

Data Type: Other Field Results

Version: 1

Version Date: 2017-01-06

Project

» [Killer Seaweeds: Allelopathy against Fijian Corals](#) (Killer Seaweeds)

Contributors	Affiliation	Role
Hay, Mark	Georgia Institute of Technology (Georgia Tech)	Principal Investigator
Ake, Hannah	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Rates of grazing by parrotfishes and macroalgal browsing in Fiji during 2010-2012 (Killer Seaweeds project)

Table of Contents

- [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
 - [Parameters](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Temporal Extent: 2010 - 2012

Dataset Description

Fiji MPA vs non-MPA corals, fishes, and herbivory. This dataset contains rates of grazing by parrotfishes and macroalgal browsing.

Acquisition Description

The study was conducted from November 2010 through February 2011 and between November 2011 and January 2012 on shallow (~1 m below the surface at low tide, equal or shallower than 2 m at high tide), intertidal fringing reefs platforms (up to 800-m wide) along the Coral Coast (18° 13.05'S, 177° 42.97'E) of Viti Levu, Fiji's main island. Many of the owners of traditional fishing rights along the Coral Coast have established small, customary no-take MPAs to improve and sustain their adjacent fishing grounds. The MPAs in this region are delimited by surface markings and enforced by local villagers, and they have been closed to all fishing activities since their inception (about 10 years). The only exception to this closure was a small experimental hook and line fishing research project that was conducted in the MPAs of Votua and Namada. In the non-MPAs, the main fishing targets are species of Acanthuridae (Nasinae), Epinephelidae, Labridae, Mullidae, and Lutjanidae. Permission for the research was granted by the Fijian Ministry of Education, National Heritage, Culture & Arts, Youth & Sports, which is authorized to approve field studies in Fijian waters. No animal collection or experimental procedures involving animals were conducted during the study, and no endangered species were recorded during our assessments.

To assess the effects of MPAs on fish assemblages, fish feeding group composition, herbivory rates, benthic cover, and coral recruit density, we compared three spatially paired MPA and adjacent, fished, areas (non-MPAs) associated with the villages of Votua, Vatu-o-lalai and Namada. Comparisons of fish assemblages inside and outside of closures are widely used for determining the effects of reserves, but it should be acknowledged that this approach does not reveal the state of an MPA relative to an undisturbed baseline.

The studied MPAs were established in 2002 (Vatu-o-lalai, Namada) and 2003 (Votua), and shortly after establishment, coral cover was low (~7%), and macroalgal cover was high (~35–45%) in both the MPAs and non-MPAs. All surveys and assays were conducted during the same season (austral summer) to minimize seasonal variation in sampling. The reef extends approx. 1 km from shore within each MPA and non-MPA, and all data were collected between 30 and 700 m of the shore (i.e., shoreward of the reef crest) parallel to

the shoreline.

Herbivory rates: Rates of grazing by parrotfishes and macroalgal browsing were assessed across the six study sites using established techniques. The feeding rates of parrotfishes were estimated within each of the six study sites from December 2011–January 2012 using remote stationary video cameras; this method was selected as it has been shown to reduce observer effects on fish behavior. Underwater cameras (GoPro) attached to a small lead weight were randomly positioned next to areas covered by algal turfs within each study site, and all feeding on the benthos was recorded for 2 hours. At the start of each video, a length of chain was used to demarcate a 4-m² area and probablyvide a scale for estimating the length of any fishes in the video. The chain was removed after one minute, and the cameras were left to record all feeding activities in the absence of divers.

To ensure similar sampling effort among sites, sampling was conducted over 18 days, always during high tide. In the first week, during which high tide occurred in the morning, four cameras were distributed in the MPA and four in the non-MPA of a given village, and over the following two days, the same procedure was repeated for the remaining two villages. A few days later, when high tide occurred during the middle of the day, the same procedure was repeated and then repeated again for the afternoon period. This entire sampling scheme was performed twice, so we recorded a total of eight videos per study site per time period within each village. All videos were subsequently viewed, and all parrotfishes observed feeding on the reef substrata were identified to species, and their length estimated. Grazing rates were then calculated as the product of species-specific bite rates and bite areas, and expressed as the percentage of the 4m² area grazed per day. Species-specific bite areas were obtained from the literature, and where these were not available the bite area of a closely related species with a similar feeding type and body size was used.

Macroalgal browsing was assessed at each site using a series of macroalgal assays during December 2011. Five common macroalgal species in the non-MPAs (*Hormophysa triquetra*, *Padina boryana*, *Sargassum polycystum*, *Sargassum* sp., and *Turbinaria ornata*) were collected by hand, spun in a salad spinner for 20 revolutions to remove water and weighed . One thallus of each alga was randomly selected and attached at equal intervals along a 60-cm length of 3-ply rope by inserting the holdfast between the strands. The order of the algal species along the rope was randomized among replicates. Three replicate assays (or ropes) were exposed to herbivores, and three assays were placed in exclusion cages (60 x 20 x 20 cm, 1-cm square mesh) at each site and left on the reef for 5 h. Assays within each site were separated by 20-50 m. After 5h the assays were collected and each thallus was carefully removed from the rope, spun and weighed, and the reduction in algal biomass was calculated.

Processing Description

We used Generalized linear mixed models (GLMM) implemented under a Bayesian framework to test the effect of protection status (MPA vs. non-MPA) on the abundance and biomass of each subcategory of Herbivores and Non-herbivores at the three village sites. We used the same approach to test the effect of protection status on the number of individuals, biomass and diversity of species in the observed feeding groups. The models have a hierarchical structure where the protection status is nested within site. For abundance and biomass of Herbivores and Non-herbivores, we used a multi-response models where each subcategory is a separate response variable. Because we have several samples for the same site, transect and sampling day were included as random factors. For the richness and abundance model, we used a Poisson error structure given the nature of the data. For biomass and diversity data we used a Gaussian error structure. We performed separate analyses for Herbivores and Non-herbivores. We compared model fit against a benchmark model in which protection status was not included as a fixed effect using the Deviance information criterion (DIC). The MCMC used to sample the posterior distributions of effect sizes ran for 106 iterations and was sampled every 100 iterations (thinning = 100) after burn-in (5×10^5). We considered effect size significant when the 95% credible interval of the estimated posterior distributions of parameters did not include 0. We monitored chain mixing by checking the effective sample sizes (ESS) for fixed and random effects. We used inverse gamma priors for variance components. Exploratory analyses indicate that estimates for fixed effects were robust to prior selection. Outliers were removed prior to the GLMM analyses to reduce overdispersion, although analyses with and without the outliers yielded qualitatively similar results. We used the R package MCMCGLMM for all analyses based on GLMMs.

We compared benthic cover between MPAs and non-MPAs using three-way ANOVA, with village site (Votua, Vatu-o-lalai, and Namada), status (MPA and non-MPA) and year (2010/2011 and 2011/2012) as fixed factors. Separate ANOVAs were used to compare the percentage cover of four different substratum types (scleractinian corals, macroalgae, epilithic algal turfs and others). Benthic cover data were arcsine-transformed, and fish density and biomass data were log-transformed to meet assumptions of normality (frequency histograms). When differences were significant, the test was followed by specific planned comparisons between paired treatments (MPA vs non-MPA) at each village site. P-values were adjusted with the Holm-Sídák method, in which the adjusted p-value is equal to $\frac{\alpha}{k}$, where k refers to the number of comparisons.

The rates of grazing and browsing and the density of coral recruits were compared between

MPAs and adjacent non-MPAs using two-way ANOVA with status (MPA and non-MPA) and village site (Votua, Vatu-o-lalai, and Namada) as fixed factors. Separate analyses were used to compare (1) parrotfish grazing rates, (2) macroalgal browsing rates, and (3) the number of coral recruits per quadrat (log-transformed). Holm-Sídák-adjusted paired comparisons were also used when differences were significant. ANOVAs for benthic cover, grazing and browsing rates, and density of coral recruits, as well as all graph plots in this manuscript, were programmed in R 3.0.1 using base package functions.

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
village	The village where sampling was performed.	unitless
protection	Status of area where sampling was done; Marine Protected Area (MPA) or non-MPA (NON)	unitless
Turbinaria_before	Percentage of Turbinaria ornata in the 4m ² area.	percent
Padina_before	Percentage of Padina boryana in the 4m ² area.	percent
S_poly_before	Percentage of Sargassum polycystum in the 4m ² area.	percent
Sarg_sp_before	Percentage of Sargassum sp. in the 4m ² area.	percent
Hormo_before	Percentage of Hormophysa triquetra in the 4m ² area.	percent
Turbinaria_after	Percentage of Turbinaria ornata in the 4m ² area after browsing.	percent
Padina_after	Percentage of Padina boryana in the 4m ² area after browsing.	percent
S_poly_after	Percentage of Sargassum polycystum in the 4m ² area after browsing.	percent
Sarg_sp_after	Percentage of Sargassum sp. in the 4m ² area after browsing.	percent
Hormo_after	Percentage of Hormophysa triquetra in the 4m ² area after browsing.	percent
Turbinaria_loss	Percentage of Turbinaria ornata loss in the 4m ² area.	percent
Padina_loss	Percentage of Turbinaria ornata loss in the 4m ² area.	percent
S_poly_loss	Percentage of Turbinaria ornata loss in the 4m ² area.	percent
Sarg_sp_loss	Percentage of Turbinaria ornata loss in the 4m ² area.	percent
Hormo_loss	Percentage of Turbinaria ornata loss in the 4m ² area.	percent

Turbinaria_loss_g	Percentage of loss g error.	percent
Padina_loss_g	Percentage of loss g error.	percent
S_poly_loss_g	Percentage of loss g error.	percent
Sarg_sp_loss_g	Percentage of loss g error.	percent
Hormo_loss_g	Percentage of loss g error.	percent
Turbinaria_log	Log transformed percentage loss.	log
Padina_log	Log transformed percentage loss.	log
S_poly_log	Log transformed percentage loss.	log
Sarg_sp_log	Log transformed percentage loss.	log
Hormo_log	Log transformed percentage loss.	log

[[table of contents](#) | [back to top](#)]

Deployments

Fiji_2011

Website	https://www.bco-dmo.org/deployment/480730
Platform	Hay_GaTech
Start Date	2010-11-01
End Date	2012-01-01
Description	Studies for this deployment were conducted: November 2010 through February 2011 and between November 2011 and January 2012 on shallow (~1 m below the surface at low tide, equal or shallower than 2 m at high tide), intertidal fringing reefs platforms in Villages of Votua, Vatu-o-lalai and Namada, Coral Coast Viti Levu, Fiji. May–December 2011 on an approximately 1.5-2.5 m deep reef flat within a no-take marine reserve at Votua Village, Viti Levu, Fiji.

Project Information

Killer Seaweeds: Allelopathy against Fijian Corals (Killer Seaweeds)

Coverage: Viti Levu, Fiji (18°13.049'S, 177°42.968'E)

Extracted from the NSF award abstract: Coral reefs are in dramatic global decline, with reefs commonly converting from species-rich and topographically-complex communities dominated by corals to species-poor and topographically-simplified communities dominated by seaweeds. These phase-shifts result in fundamental loss of ecosystem function. Despite debate about whether coral-to-algal transitions are commonly a primary cause, or simply a consequence, of coral mortality, rigorous field investigation of seaweed-coral competition has received limited attention. There is limited information on how the outcome of seaweed-coral competition varies among species or the relative importance of different competitive mechanisms in facilitating seaweed dominance. In an effort to address this topic, the PI will conduct field experiments in the tropical South Pacific (Fiji) to determine the effects of seaweeds on corals when in direct contact, which seaweeds are most damaging to corals, the role allelopathic lipids that are transferred via contact in producing these effects, the identity and surface concentrations of these metabolites, and the dynamic nature of seaweed metabolite production and coral response following contact. The herbivorous fishes most responsible for controlling allelopathic seaweeds will be identified, the roles of seaweed metabolites in allelopathy vs herbivore deterrence will be studied, and the potential for better managing and conserving critical reef herbivores so as to slow or reverse conversion of coral reef to seaweed meadows will be examined. Preliminary results indicate that seaweeds may commonly damage corals via lipid-soluble allelochemicals. Such chemically-mediated interactions could kill or damage adult corals and produce the suppression of coral fecundity and recruitment noted by previous investigators and could precipitate positive feedback mechanisms making reef recovery increasingly unlikely as seaweed abundance increases. Chemically-mediated seaweed-coral competition may play a critical role in the degradation of present-day coral reefs. Increasing information on which seaweeds are most aggressive to corals and which herbivores best limit these seaweeds may prove useful in better managing reefs to facilitate resilience and possible recovery despite threats of global-scale stresses. Fiji is well positioned to rapidly use findings from this project for better management of reef resources because it has already erected >260 MPAs, Fijian villagers have already bought-in to the value of MPAs, and the Fiji Locally-Managed Marine Area (FLMMA) Network is well organized to get information to villagers in a culturally sensitive and useful manner. The

broader impacts of this project are far reaching. The project provides training opportunities for 2-2.5 Ph.D students and 1 undergraduate student each year in the interdisciplinary areas of marine ecology, marine conservation, and marine chemical ecology. Findings from this project will be immediately integrated into classes at Ga Tech and made available throughout Fiji via a foundation and web site that have already set-up to support marine conservation efforts in Fiji and marine education efforts both within Fiji and internationally. Business and community leaders from Atlanta (via Rotary International Service efforts) have been recruited to help organize and fund community service and outreach projects in Fiji -- several of which are likely to involve marine conservation and education based in part on these efforts there. Media outlets (National Geographic, NPR, Animal Planet, Audubon Magazine, etc.) and local Rotary clubs will be used to better disseminate these discoveries to the public. PUBLICATIONS PRODUCED AS A RESULT OF THIS RESEARCH Rasher DB, Stout EP, Engel S, Kubanek J, and ME Hay. "Macroalgal terpenes function as allelopathic agents against reef corals", Proceedings of the National Academy of Sciences, v. 108, 2011, p. 17726. Beattie AJ, ME Hay, B Magnusson, R de Nys, J Smeathers, JFV Vincent. "Ecology and bioprospecting," Austral Ecology, v.36, 2011, p. 341. Rasher DB and ME Hay. "Seaweed allelopathy degrades the resilience and function of coral reefs," Communicative and Integrative Biology, v.3, 2010. Hay ME, Rasher DB. "Corals in crisis," The Scientist, v.24, 2010, p. 42. Hay ME and DB Rasher. "Coral reefs in crisis: reversing the biotic death spiral," Faculty 1000 Biology Reports 2010, v.2, 2010. Rasher DB and ME Hay. "Chemically rich seaweeds poison corals when not controlled by herbivores", Proceedings of the National Academy of Sciences, v.107, 2010, p. 9683.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0929119
National Institutes of Health (NIH)	U01-TW007401

[[table of contents](#) | [back to top](#)]