

Counts of prey species found in harbor seal scat samples collected in the San Juan Islands from 2005-2008 (Seal_response_to_prey project)

Website: <https://www.bco-dmo.org/dataset/3817>

Data Type: Other Field Results

Version: 1

Version Date: 2012-12-13

Project

» [Responses of Seals and Sea Lions to Increased Rockfish Density](#) (Seal_response_to_prey)

Contributors	Affiliation	Role
Acevedo-Gutierrez, Alejandro	Western Washington University (WWU)	Lead Principal Investigator
Bromaghin, Jeffrey F	United States Geological Survey (USGS)	Co-Principal Investigator
Jeffries, Steven J	Washington Department of Fish and Wildlife	Co-Principal Investigator
Kennish, John M	University of Alaska, Anchorage (UAA)	Co-Principal Investigator
Lance, Monique M	Washington Department of Fish and Wildlife	Co-Principal Investigator
Levin, Philip S	National Marine Fisheries Service (NMFS)	Co-Principal Investigator
Rauch, Shannon	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Counts of prey species found in harbor seal scat samples collected in the San Juan Islands from 2005-2008.

Table of Contents

- [Dataset Description](#)
 - [Acquisition Description](#)

- [Processing Description](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Deployments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Temporal Extent: 2005-03-01 - 2008-02-01

Dataset Description

Counts of prey species identified in harbor seal scat samples collected from 2005 to 2008.

Also see the related data set, "seal_pre_y_by_region", in which frequencies of occurrence are provided for prey groupings for different sampling regions within the San Juan Islands.

Data and methods, including sample sizes, are described in:

Lance, M. M., Chang, W.-Y., Jeffries, S. J., Pearson, S. F. & Acevedo-Gutierrez, A. 2012. Harbor seal diet in northern Puget Sound: implications for the recovery of depressed fish stocks. Marine Ecology Progress Series 464:257-271. DOI:[10.3354/meps09880](https://doi.org/10.3354/meps09880)

Acquisition Description

From Lance et al.:

From 2005 to 2008, fecal samples (scats) were collected seasonally over 3 collection periods: March to early June ('spring'), late July to September ('summer/fall'), and January to February ('winter'). Scats were collected from 23 haul-out locations during daytime low tides. The sites were dispersed throughout the study area, represented various habitat types used by harbor seals, were the largest haul-outs where adequate sample sizes could be collected, and were easily accessible by boat. Two or three collection trips were made each season, with a target sample size of 60 scats per season per region.

Samples were stored frozen. In the lab, samples were enclosed in fine mesh paint-strainer bags and cleaned using a washing machine or nested sieves. Hard parts were cleaned and stored dry. Prey were identified to lowest possible taxon using a dissecting microscope, reference fish bone collections, and published keys.

Processing Description

BCO-DMO made the following modifications:

- Changed parameter names to conform with BCO-DMO convention.
- Replaced spaces with underscores.
- Replaced blanks with 'nd' to indicate 'no data'.

[[table of contents](#) | [back to top](#)]

Related Publications

Lance, M., Chang, W., Jeffries, S., Pearson, S., & Acevedo-Gutiérrez, A. (2012). Harbor seal diet in northern Puget Sound: implications for the recovery of depressed fish stocks. *Marine Ecology Progress Series*, 464, 257–271. doi:[10.3354/meps09880](https://doi.org/10.3354/meps09880)

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
species	Taxonomic name of the prey species.	text
common_name	Common name of the prey species.	text
family	Taxonomic family name.	text
season	Sampling season: Spring = March to early June, Summer/Fall = late July to September, Winter = January to February.	text
count	Total number of occurrences of the species identified in the seal scat samples.	integer
pcnt_unweighted	Unweighted occurrence frequency of the species.	%

[[table of contents](#) | [back to top](#)]

Deployments

lab_Acevedo

Website	https://www.bco-dmo.org/deployment/58897
Platform	WWU
Start Date	2005-03-01
End Date	2008-02-01
Description	Marine Behavior and Ecology Lab at Western Washington University, led by Alejandro Acevedo-Gutierrez, lead PI for the project "Responses of Seals and Sea Lions to Increased Rockfish Density" (NSF OCE-0550443). Address: Department of Biology Western Washington University 516 High St. Bellingham, WA 98225

[[table of contents](#) | [back to top](#)]

Project Information

Responses of Seals and Sea Lions to Increased Rockfish Density (Seal_response_to_pre)

Website: <http://biol.wwu.edu/mbel/?page=research>

Coverage: Salish Sea, USA and Canada

From NSF proposal: This project is a collaborative study of the responses of harbor seals and other mammalian predators to changes in prey density in Puget Sound. The general study approach will involve multi-year field estimates to observe the responses of predators to rockfish density in protected areas, candidate marine reserves, and unprotected sites. The collaborating investigators will estimate 1) rockfish density using visual and mark and recapture techniques; 2) predator abundance using aerials surveys and dedicated land observations; and 3) predator food consumption using scat to describe diet, tagging of harbor seals to describe individual foraging sites, and population-based and individual bioenergetics models to describe consumption of rockfish. The investigators will also take into account confounding factors that might explain predator behavior, such as environmental variables and alternative prey, by creating a GIS database from available information from the area. The different field observations and database estimates are explicitly linked through a common hypothesis and coordinated methodologies, and their results will be integrated into a model describing the impact of predation on rockfish populations. The responses of top predators to changes in prey density and their impact on fish populations of interest are unknown. This

study will evaluate the effectiveness of MPAs as fish refugia, offer a framework for the management and conservation of marine resources, and provide an exciting opportunity for students to participate in ecological and conservation research. Hypotheses: 1) Harbor seals and other pinniped species show aggregative responses to changes in prey density. Hence, their abundance will increase with fish density. 2) Harbor seals and other pinniped species show Type 2 or 3 functional responses to changes in prey density. Thus, their consumption rate of a particular prey type follows an asymptotic or sigmoidal curve relative to the prey's density, respectively. 3) Predation by harbor seals and other pinniped species is sufficiently intense that it impedes recovery of depleted fish populations. Objectives: 1) Quantify the number of harbor seals and other pinniped species in relation to rockfish density and other environmental (confounding) factors. 2) Estimate the consumption rate of harbor seals and other pinniped species in relation to rockfish density and other prey species. 3) Correlatively estimate the influence of predation by harbor seals and other pinniped species on survivorship and population size of rockfish. Publications resulting from this NSF award: Bjorland, R. H., Pearson, S. F., Jeffries, S. J., Lance, M. M., Acevedo-Gutiérrez, A. & Ward, E. J. 2015. Stable isotope mixing models elucidate sex and size effects on the diet of a generalist marine predator. *Marine Ecology Progress Series* 526: 213-225. DOI: 10.3354/meps11230Bromaghin, J. F., Lance, M. M., Elliott, E. W., Jeffries, S. J., Acevedo-Gutierrez, A. & Kennish, J. M. 2013. New insights into the diets of harbor seals in the Salish Sea of western North America revealed by quantitative fatty acid signature analysis. *Fishery Bulletin* 111: 13-26. DOI: 10.7755/FB.111.1.2 Buzzell, B.1, Lance, M. & Acevedo-Gutiérrez, A. 2014. Spatial and temporal variation in river otter (*Lontra canadensis*) diet and predation on rockfish (Genus *Sebastes*) in the San Juan Islands, Washington. *Aquatic Mammals* 40: 150- 161. DOI: 10.1578/AM.40.2.2014.150 Howard, S., Lance, M., Jeffries, S. & Acevedo-Gutierrez, A. 2013. Fish consumption by harbor seals (*Phoca vitulina*) in the San Juan Islands, WA. *Fishery Bulletin* 111: 27-41. DOI: 10.7755/FB.111.1.3 Lance, M. M., Chang, W.-Y., Jeffries, S. J., Pearson, S. F. & Acevedo-Gutierrez, A. 2012. Harbor seal diet in northern Puget Sound: implications for the recovery of depressed fish stocks. *Marine Ecology Progress Series* 464:257-271. DOI:10.3354/meps09880 Luxa, K. & Acevedo-Gutierrez, A. 2013. Food habits of harbor seals (*Phoca vitulina*) in two estuaries in the central Salish Sea. *Aquatic Mammals* 39: 10- 22. DOI: 10.1578/AM.39.1.2013.10 Peterson, S., Lance, M. M., Jeffries, S. J. & Acevedo-Gutierrez, A. 2012. Long distance movements and disjunct spatial use of harbor seals (*Phoca vitulina*) in the inland waters of the Pacific Northwest. *PLoS ONE* 7: e39046. DOI: 10.1371/journal.pone.0039046 Thomas, AC; Lance, MM; Jeffries, SJ; Miner, BG; Acevedo-Gutierrez, A. 2011. Harbor seal foraging response to a seasonal resource pulse, spawning Pacific herring. *Marine Ecology-Progress Series*, v.441. p. 225. DOI: 10.3354/meps09370 Ward, EJ; Levin, PS; Lance, MM; Jeffries, SJ; Acevedo-Gutierrez, A. 2012. Integrating diet and movement data to identify hot spots of predation risk and areas of conservation concern for endangered species. *Conservation Letters*, v.5, p. 37. DOI: 10.1111/j.1755-263X.2011.00210.x Wilson, K.2, Lance, M., Jeffries, S. & Acevedo-Gutiérrez, A.

2014. Fine-scale variability in harbor seal foraging behavior. PLoS ONE 9: e92838. DOI: 10.1371/journal.pone.0092838.

[[table of contents](#) | [back to top](#)]

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-0550443

[[table of contents](#) | [back to top](#)]