

Table 2: Suspended particulate matter (SPM) concentrations in lake water

Website: <https://www.bco-dmo.org/dataset/809857>

Data Type: Other Field Results

Version: 1

Version Date: 2020-04-22

Project

» [The measurement and use of bismuth-210 as a tracer of particle flux in aquatic systems](#) (Bi 210 measurement)

Contributors	Affiliation	Role
Waples, James T.	University of Wisconsin (UW-Milwaukee)	Principal Investigator
Copley, Nancy	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

Abstract

Table 2. Suspended particulate matter (SPM) concentrations in lake water, Lake Michigan, Nov. 2018

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
 - [Parameters](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: Lat:43.01784 Lon:-87.90352

Temporal Extent: 2018-11-08 - 2018-11-20

Dataset Description

Lake water samples (~ 50 L) for radionuclide analyses were collected from the slip (43.017835, -87.903521) with a submersible pump from a depth of 2 m on 08 November, 15 November, and 20 November 2018.

Acquisition Description

A detailed description of the methods for determining in situ ²¹⁰Pb, ²¹⁰Bi, and ²¹⁰Po activities is given by Waples (2020).

A second manuscript describing the use of Pb-210, Bi-210, and Po-210 as particle tracers in aquatic systems is currently in review for Limnology and Oceanography (March/2020).

Lake water samples (~ 50 L) for radionuclide analyses were collected from the slip (43.017835, -87.903521) with a submersible pump from a depth of 2 m on 08 November, 15 November, and 20 November 2018.

Water samples collected by submersible pump were weighed and filtered through nitrocellulose filters (0.45 μm , 293 mm, Millipore) within two hours of sample collection to separate dissolved and particle-bound nuclide fractions of ^{210}Pb , ^{210}Bi , and ^{210}Po . Yield monitors of ^{207}Bi and ^{209}Po (Eckert & Ziegler Isotope Products) were then added to both fractions. No lead monitor was required. Dissolved nuclide fractions ($< 0.45 \mu\text{m}$) of ^{210}Pb , ^{210}Bi , and ^{210}Po were co-precipitated onto newly formed ferric hydroxide and collected by filtration. Both particle-bound and dissolved nuclide fractions were then digested in concentrated nitric acid. Bismuth and polonium were completely separated from lead on EmporeTM anion solid phase extraction (SPE) disks (3M, 47 mm, product number 2252, now manufactured by CDS Analytical). Eluate from the particle-bound fraction was re-spiked with ^{207}Bi and set aside for ~one month for the determination of particle-bound ^{210}Pb (via ^{210}Bi in secular equilibrium with its parent). Anion SPE disks were then counted– first for ^{210}Bi via β -decay on a low background gas-flow proportional detector with anticoincidence circuitry (G542 System, Gamma Products), then for ^{207}Bi via γ -emission to determine yield. Polonium was then removed from the SPE disks, plated to copper, and α -counted for ^{210}Po and the yield monitor ^{209}Po . Total ^{210}Pb was determined in a separate water sample via ^{210}Bi that had grown into secular equilibrium with its parent. Activity calculations including uncertainty propagation and decay corrections are all fully described by Waples (2020).

Lake water samples ($\sim 50 \text{ L}$) for radionuclide analyses were collected from the slip (43.017835, -87.903521) with a submersible pump from a depth of 2 m on 08 November, 15 November, and 20 November 2018.

Software used:

- Maestro software (v. 6.04 and 6.06, Advanced Measurement Technology)
- SigmaPlot (v. 11, Systat Software)
- Excel (v. 2002, Microsoft)

Processing Description

BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- re-formatted date from m/d/yy to yyyy-mm-dd, added ISO_DateTime.UTC

[[table of contents](#) | [back to top](#)]

Parameters

Parameter	Description	Units
sample	sample identifier	unitless
ISO_DateTime_Local	sample collection local date and time; ISO formatted yyyy-mm-ddTHH:MM:SS (UTC- 6 hr)	unitless
ISO_DateTime_UTC	sample collection date and time; ISO formatted yyyy-mm-ddTHH:MM:SSZ (converted to UTC from local: UTC- 6 hr)	unitless
latitude	latitude; north is positive	decimal degrees
longitude	longitude; east is positive	decimal degrees
depth	sample depth	meters
SPM_b	suspended particulate matter ($>0.45 \mu\text{m}$)	grams/meter ³ (g m ⁻³)

Project Information

The measurement and use of bismuth-210 as a tracer of particle flux in aquatic systems (Bi 210 measurement)

Coverage: Lake Michigan 43.0 N 87.9 W

Naturally occurring radioactive nuclides provide invaluable "clocks" with which to investigate a wide range of processes in earth and environmental sciences. Nuclides with shorter half-lives are well suited to studying rapid processes, but are often present at very low concentrations and are particularly difficult to measure. The goal of this Early Concept Grant for Exploratory Research (EAGER) is to develop and demonstrate a new analytical method for measuring bismuth-210 (half life, 5 days) in lake water, seawater, and associated particles. Because bismuth tends to become associated with particles more than its parent isotope lead-210 (half life, 22 years), the bismuth-210/lead-210 daughter/parent pair can potentially be used to investigate very short time-scale processes associated with particle (sediment) formation and cycling in lakes and the coastal ocean. Knowledge about such processes is important to understanding carbon and nutrient cycling in these productive and dynamic areas. The specific goals of this exploratory project are to 1) demonstrate an improved method for measuring the lead-210/bismuth-210/polonium-210 trio in aquatic system samples; 2) measure lead-210/bismuth-210/polonium-210 in Lake Michigan; and 3) explore the feasibility of using bismuth-210/lead-210 disequilibrium as a proxy for particle flux. The project would support an undergraduate student researcher for the summer.

Funding

Funding Source	Award
NSF Division of Ocean Sciences (NSF OCE)	OCE-1639865