

Annual rainfall between Cabritte Horn and White Point on the south shore of St. John, US Virgin Islands, 1996-2018

Website: <https://www.bco-dmo.org/dataset/827806>

Data Type: Other Field Results

Version: 1

Version Date: 2020-10-28

Project

» [RUI-LTREB Renewal: Three decades of coral reef community dynamics in St. John, USVI: 2014-2019](#)
(RUI-LTREB)

| Contributors | Affiliation | Role |
|-----------------------------------|---|------------------------|
| Edmunds, Peter J. | California State University Northridge (CSU-Northridge) | Principal Investigator |
| Copley, Nancy | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

Annual rainfall between Cabritte Horn and White Point on the south shore of St. John, US Virgin Islands, July and August, 1996-2018, used in the associated mortality and growth analyses.

Table of Contents

- [Coverage](#)
 - [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
 - [Related Datasets](#)
 - [Parameters](#)
 - [Instruments](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Spatial Extent: N:18.31714 E:-64.7208 S:18.30702 W:-64.73152

Temporal Extent: 1996 - 2018

Acquisition Description

From publication Limnology and Oceanography

Environmental conditions:

The rainfall data integrated values between samplings, and it was used to provide a proxy for environmental conditions positively associated with rainfall, including reduced surface light, enhanced seawater turbidity, terrestrial runoff, sedimentation, and nutrients. Rainfall from August 1996 to 2011 was obtained from the Southeastern Regional Climate Centre (<http://www.sercc.com>), which reports data from a Standard Rain Gauge in Cruz Bay (Station 671980), 7 km west of Great Lameshur Bay. When this record was incomplete, data were obtained from East End (Station 672551, ~ 5 km east of Great Lameshur Bay, 21 of 185 records), or values for missing months were interpolated using a grand mean from all other values for the same month from 1972 to 2011 (16 of 185 records). From 2012 to the present, rainfall was measured using a Standard Rain Gauge deployed on the north shore of St. John (18.356°N, 64.766°W). The gauge was manually recorded and values summed by month. The rainfall over each sampling year

was calculated by summing monthly values from August to July of the subsequent year.

Processing Description

BCO-DMO Processing Notes:

- data extracted to .csv from file "Flor L&O Edmunds 2020 (matrix paper).zip/Data in L&O Paper_21_Oct_2020.xlsx", sheet "rainfall".
- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions

[[table of contents](#) | [back to top](#)]

Related Datasets

IsSupplementTo

Edmunds, P. J. (2020) **Growth and mortality rates of small corals in St. John, US Virgin Islands, 1996-2018**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2020-10-28 <http://lod.bco-dmo.org/id/dataset/827764> [[view at BCO-DMO](#)]

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|-------------|---------------------------------|------------------|
| Year | study year | unitless |
| rainfall_cm | total annual amount of rainfall | centimeters/year |

[[table of contents](#) | [back to top](#)]

Instruments

| | |
|---|-------------------------------------|
| Dataset-specific Instrument Name | Standard Rain Gauge |
| Generic Instrument Name | Precipitation Gauge |
| Generic Instrument Description | measures rain or snow precipitation |

[[table of contents](#) | [back to top](#)]

Project Information

RUI-LTREB Renewal: Three decades of coral reef community dynamics in St. John, USVI: 2014-2019 (RUI-LTREB)

Website: <http://coralreefs.csun.edu/>

Coverage: USVI

Describing how ecosystems like coral reefs are changing is at the forefront of efforts to evaluate the biological consequences of global climate change and ocean acidification. Coral reefs have become the poster child of these efforts. Amid concern that they could become ecologically extinct within a century, describing what has been lost, what is left, and what is at risk, is of paramount importance. This project exploits an unrivalled legacy of information beginning in 1987 to evaluate the form in which reefs will persist, and the extent to which they will be able to resist further onslaughts of environmental challenges. This long-term project continues a 27-year study of Caribbean coral reefs. The diverse data collected will allow the investigators to determine the roles of local and global disturbances in reef degradation. The data will also reveal the structure and function of reefs in a future with more human disturbances, when corals may no longer dominate tropical reefs. The broad societal impacts of this project include advancing understanding of an ecosystem that has long been held emblematic of the beauty, diversity, and delicacy of the biological world. Proposed research will expose new generations of undergraduate and graduate students to natural history and the quantitative assessment of the ways in which our planet is changing. This training will lead to a more profound understanding of contemporary ecology at the same time that it promotes excellence in STEM careers and supports technology infrastructure in the United States. Partnerships will be established between universities and high schools to bring university faculty and students in contact with k-12 educators and their students, allow teachers to carry out research in inspiring coral reef locations, and motivate children to pursue STEM careers. Open access to decades of legacy data will stimulate further research and teaching.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|---|-----------------------------|
| NSF Division of Environmental Biology (NSF DEB) | DEB-1350146 |

[[table of contents](#) | [back to top](#)]