

Trajectories and velocities of *Crepidula fornicata* larvae swimming under various food cues from experiments conducted at Woods Hole Oceanographic Institution in July, 2018

Website: <https://www.bco-dmo.org/dataset/834221>

Data Type: experimental

Version: 1

Version Date: 2020-12-17

Project

» [Larval Response to Turbulence During Dispersal and Settlement](#) (Larvae in turbulence)

| Contributors | Affiliation | Role |
|---------------------------------------|---|---------------------------|
| Mullineaux, Lauren | Woods Hole Oceanographic Institution (WHOI) | Principal Investigator |
| Helfrich, Karl R. | Woods Hole Oceanographic Institution (WHOI) | Co-Principal Investigator |
| DiBenedetto, Michelle | Woods Hole Oceanographic Institution (WHOI) | Contact |
| York, Amber D. | Woods Hole Oceanographic Institution (WHOI BCO-DMO) | BCO-DMO Data Manager |

Abstract

Trajectories and velocities of *Crepidula fornicata* larvae swimming under various food cues from experiments conducted at Woods Hole Oceanographic Institution in July, 2018. These data were published in DiBenedetto et al. (2020).

Table of Contents

- [Dataset Description](#)
 - [Acquisition Description](#)
 - [Processing Description](#)
 - [Related Publications](#)
 - [Parameters](#)
 - [Project Information](#)
 - [Funding](#)
-

Coverage

Temporal Extent: 2018-07

Acquisition Description

Trajectory data of *Crepidula fornicata* larvae swimming in 65mL still water flasks, analyzed from video footage. Two experimental controls were varied, resulting in four experiment types. Larvae were either previously fed or starved (Fed/St), and either in the presence of food or not in the presence of food (F/0). Five replicates (1-5) were conducted for each of the four conditions, resulting in 20 individual datasets. Further description and analysis of the data is published in DiBenedetto et al., 2020 (<https://doi.org/10.1242/jeb.239178>).

Processing Description

Video footage was processed in MATLAB. The centroids of the larvae were identified in each image. The centroids were linked to form trajectories using a predictive tracking algorithm. Linear and angular velocities were calculated along each trajectory, as described in DiBenedetto et al. 2020.

BCO-DMO Data Manager Processing notes:

- * 20 csv files containing larvae tracks combined into one data table with added column to capture information in the file name(treatment variables, replication information).
- * Numbers with 15 decimal places rounded to 5 decimal places.
- * Original missing data identifier "NaN" is converted to compatible missing data identifiers based on file type (matlab, netcdf, etc). 'nd' meaning "no data" is the default missing data identifier at BCO-DMO.

[[table of contents](#) | [back to top](#)]

Related Publications

DiBenedetto, M. H., Meyer-Kaiser, K. S., Torjman, B., Wheeler, J. D., & Mullineaux, L. S. (2020). Departures from isotropy: the kinematics of a larval snail in response to food. *The Journal of Experimental Biology*, jeb.239178. doi:[10.1242/jeb.239178](https://doi.org/10.1242/jeb.239178)
Results

[[table of contents](#) | [back to top](#)]

Parameters

| Parameter | Description | Units |
|----------------|--|-------------------------------|
| Exp_name | Name for the experimental run composed of treatment information (Previously_Fed, Food_Presence,Replicate). | unitless |
| Previously_Fed | Whether larvae were either previously fed ('Fed') or starved ('St'). | unitless |
| Food_Presence | Whether larvae were in the presence of food ('F') or not in the presence of food ('0'). | unitless |
| Replicate | Replicate number | unitless |
| track | Larva track number | unitless |
| T | Time (seconds) elapsed since the start of the experiment. | seconds (sec) |
| X | Horizontal position | centimeters (cm) |
| Y | Vertical position | centimeters (cm) |
| U | Horizontal velocity | centimeters per second (cm/s) |
| V | Vertical velocity | centimeters per second (cm/s) |
| Theta | Angular velocity | radians per second (rad/s) |

[[table of contents](#) | [back to top](#)]

Project Information

Larval Response to Turbulence During Dispersal and Settlement (Larvae in turbulence)

Coverage: Laboratory studies at Woods Hole Oceanographic Institution

Description from NSF award abstract:

The planktonic larval stage of benthic marine invertebrates provides a mechanism for exchange of individuals between remote populations. Dispersal is affected by swimming behaviors, particularly those that alter the larva's vertical position in the water. Larvae of some species change their vertical positions in response to turbulence by ceasing to swim and sinking downward (diving). By doing so, they can alter their horizontal transport in currents and increase their supply to the seafloor. The main objectives of this study are to investigate behavioral responses of oyster (*Crassostrea virginica*) larvae to turbulence in the water column and at the seafloor, and to determine how these behaviors affect settlement. The investigators hypothesize that diving behavior enhances settlement into suitable habitat, even where mean bed shear stress is high. They expect that once larvae approach the bottom, they can take advantage of temporal and spatial refuges (such as turbulent lulls in the lee of roughness elements) to settle in otherwise harsh conditions. Investigating larval responses to turbulence is a challenge because it requires simultaneous measurement of time-variant flows and larval behaviors. The investigators will modify a conventional particle image velocimetry (PIV) approach so it can be used to track larval motions and fluid velocities simultaneously. PIV provides information on flow kinematics (e.g., rotation and strain rate) in the immediate vicinity of a larva, as well as bulk dissipation rates and measures of Taylor and integral length scales that likely influence larval acceleration. When these measurements are coupled with a larval trajectory, they provide a history of the fluid environment a larva experiences, and can be used to determine what characteristic of turbulence triggers the diving behavior. They also make it possible to calculate the bottom shear stress an individual larva experiences when it encounters the bottom and attempts to settle. The investigators will examine turbulence effects on larval behaviors in the water column using a grid-stirred tank. They will use a racetrack flume to test the hypothesis that larval settlement success depends on the frequency of lulls of sufficient duration for larval attachment.

Laboratory experiments will provide a mechanistic understanding of larval behavior that can be used in general theoretical models exploring how behavior influences dispersal and population connectivity. The quantified swimming responses of oysters are critical input for coupled bio-physical models of dispersal in the field. An understanding of larval behavior contributes to our ability to predict the effects of natural and anthropogenic perturbations (some of which are linked to global climate change) on benthic communities in coastal ecosystems where turbulence and habitat suitability vary spatially. This information is critical for informed decision making on shellfish management and design of marine reserves. The technique developed for simultaneous PIV and larval tracking will open new questions in larval ecology and be broadly applicable to studies of plankton interactions with turbulence.

[[table of contents](#) | [back to top](#)]

Funding

| Funding Source | Award |
|--|-----------------------------|
| NSF Division of Ocean Sciences (NSF OCE) | OCE-0850419 |

[[table of contents](#) | [back to top](#)]