

# Multilocus microsatellite genotypes of *Acropora palmata* from the Caribbean and North-West Atlantic from 2015-2016 (Coral Hybridization project)

**Website:** <https://www.bco-dmo.org/dataset/666350>

**Data Type:** Other Field Results

**Version:** 1

**Version Date:** 2016-11-30

## Project

» [Collaborative research: Is hybridization among threatened Caribbean coral species the key to their survival or the harbinger of their extinction?](#) (Coral Hybridization)

Contributors	Affiliation	Role
<a href="#">Baums, Iliana B.</a>	Pennsylvania State University (PSU)	Principal Investigator
<a href="#">Copley, Nancy</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

Genotypes for the coral, *Acropora palmata* collected throughout the Caribbean from five microsatellite loci. *A. palmata* is a diploid species, however, somatic mutations in the form of third and fourth alleles per locus are occasionally detected. Data are in columns of region, database\_id, genet\_id, mutant\_id, locus, allele, and size\_bp. These data were deposited in DRYAD: <http://dx.doi.org/10.5061/dryad.f6600> and published in the paper Devlin-Durante et al, Mol Ecol. (2016).

## Table of Contents

- [Coverage](#)
- [Dataset Description](#)
  - [Acquisition Description](#)
  - [Processing Description](#)
- [Related Publications](#)
- [Related Datasets](#)
- [Parameters](#)
- [Instruments](#)
- [Deployments](#)
- [Project Information](#)
- [Funding](#)

## Coverage

**Spatial Extent:** N:26.7075 E:-64.677 S:12.0831 W:-88.199

## Acquisition Description

Adult *Acropora palmata* colonies were sampled (1 cm<sup>2</sup>) with hammer and chisel and tissues were preserved in 96% non-denature Ethanol. DNA was extracted using the DNEasy tissue kit (Qiagen) following manufacturer's instructions. Two multiplex Polymerase Chain Reactions (PCR) were performed per sample using fluorescently labeled primers to assay five loci containing AAT repeats. These five microsatellite loci have previously been demonstrated to be mendelian and coral-specific using controlled crosses (Baums et al., 2005). PCR products were visualized with an automated sequencer (ABI 3730). An internal size standard (Gene Scan 500-Liz, Applied Biosystems CA) ensured accurate sizing. Electropherograms were analyzed with GeneMapper Software 3.0 (Applied Biosystems, CA). Alleles were

scored as PCR product size.

## References:

Baums IB, Devlin-Durante MK, LaJeunesse TC (2014) New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population genetic studies. *Molecular Ecology* 23, 4203-4215.

Baums IB, Hughes CR, Hellberg MH (2005) Mendelian microsatellite loci for the Caribbean coral *Acropora palmata*. *Marine Ecology - Progress Series* 288, 115-127 for details on PCR amplification.

Devlin-Durante MK, Miller MW, Caribbean Acropora Research G, Precht WF, Baums IB. How old are you? Genet age estimates in a clonal animal. *Mol Ecol*. 2016. doi: 10.1111/mec.13865 for details on how mutations were scored.

## Processing Description

Electropherograms of microsatellite alleles were scored for allele sizes (basepairs) in Genemapper vers. (Applied Biosystems) and transferred to spreadsheets.

### BCO-DMO Processing Notes:

- added conventional header with dataset name, PI name, version date
- modified parameter names to conform with BCO-DMO naming conventions
- nd (no data) was entered into all blank cells
- replaced spaces and / with underscores
- converted the submitted multilevel table to a flat table

VERSIONS: 2017-21-26: Replaced version 1 (2016-11-30) with version 2 (2017-10-19). There was a shift in 1 basepair at locus 181 for some of the data.

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Baums, I. B., Devlin-Durante, M. K., & LaJeunesse, T. C. (2014). New insights into the dynamics between reef corals and their associated dinoflagellate endosymbionts from population genetic studies. *Molecular Ecology*, 23(17), 4203–4215. doi:[10.1111/mec.12788](https://doi.org/10.1111/mec.12788)

*General*

Baums, I., Hughes, C., & Hellberg, M. (2005). Mendelian microsatellite loci for the Caribbean coral *Acropora palmata*. *Marine Ecology Progress Series*, 288, 115–127. doi:[10.3354/meps288115](https://doi.org/10.3354/meps288115)

*Methods*

Devlin-Durante, M. K., Miller, M. W., Precht, W. F., & Baums, I. B. (2016). How old are you? Genet age estimates in a clonal animal. *Molecular Ecology*, 25(22), 5628–5646. doi:[10.1111/mec.13865](https://doi.org/10.1111/mec.13865)

*Results*

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### IsRelatedTo

Baums, I. B. (2017) **Mutations at microsatellite loci of *Acropora palmata* in the Caribbean and North-West Atlantic from 2015-2016 (Coral Hybridization project)**. Biological and Chemical

## Different Version

Devlin-Durante, M. K., Miller, M. W., Caribbean Acropora Research Group, Precht, W. F., and Baums, I. B. (2017). Data from: How old are you? Genet age estimates in a clonal animal (Version 2) [Data set]. Dryad Digital Repository. <https://doi.org/10.5061/dryad.f6600>

[ [table of contents](#) | [back to top](#) ]

## Parameters

Parameter	Description	Units
region	sampling region	unitless
database_id	Baums lab sample Database ID (unique for each sample)	unitless
genet_id	Genets are identified by their Genet ID. All samples that share all ancestral alleles at five microsatellite loci were assigned the same genet ID. See methods for how ancestral alleles were determined.	unitless
mutant_id	A Unique ID given to each unique multilocus genotype. This multilocus genotype includes all alleles (diploid ancestral and additional mutations).	unitless
locus	the microsatellite locus that had a mutated allele (166; 181; 182; 192; 207)	basepairs
size_bp	size of allele?	basepairs
allele	allele identifier	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	automated sequencer (ABI 3730)
<b>Generic Instrument Name</b>	Automated DNA Sequencer
<b>Generic Instrument Description</b>	General term for a laboratory instrument used for deciphering the order of bases in a strand of DNA. Sanger sequencers detect fluorescence from different dyes that are used to identify the A, C, G, and T extension reactions. Contemporary or Pyrosequencer methods are based on detecting the activity of DNA polymerase (a DNA synthesizing enzyme) with another chemoluminescent enzyme. Essentially, the method allows sequencing of a single strand of DNA by synthesizing the complementary strand along it, one base pair at a time, and detecting which base was actually added at each step.

<b>Dataset-specific Instrument Name</b>	ABI 3730
<b>Generic Instrument Name</b>	PCR Thermal Cycler
<b>Generic Instrument Description</b>	General term for a laboratory apparatus commonly used for performing polymerase chain reaction (PCR). The device has a thermal block with holes where tubes with the PCR reaction mixtures can be inserted. The cycler then raises and lowers the temperature of the block in discrete, pre-programmed steps. (adapted from <a href="http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html">http://serc.carleton.edu/microbelife/research_methods/genomics/pcr.html</a> )

[ [table of contents](#) | [back to top](#) ]

---

## Deployments

### Baums\_Carib

<b>Website</b>	<a href="https://www.bco-dmo.org/deployment/666348">https://www.bco-dmo.org/deployment/666348</a>
<b>Platform</b>	Penn_State_U
<b>Start Date</b>	2015-10-01
<b>End Date</b>	2016-10-26
<b>Description</b>	genetics studies

[ [table of contents](#) | [back to top](#) ]

---

## Project Information

### **Collaborative research: Is hybridization among threatened Caribbean coral species the key to their survival or the harbinger of their extinction? (Coral Hybridization)**

**Coverage:** Caribbean and North-West Atlantic

#### *NSF Award Abstract:*

Reef-building acroporid corals form the foundation of shallow tropical coral communities throughout the Caribbean. Yet, the once dominant staghorn coral (*Acropora cervicornis*) and the elkhorn coral (*A. palmata*) have decreased by more than 90% since the 1980s, primarily from disease. Their continuing decline jeopardizes the ability of coral reefs to provide numerous societal and ecological benefits, including economic revenue from seafood harvesting and tourism and shoreline protection from extreme wave events caused by storms and hurricanes. Despite their protection under the U.S. Endangered Species Act since 2006, threats to the survival of reef-building acroporid corals remain pervasive and include disease and warming ocean temperatures that may lead to further large-scale mortality. However, hybridization among these closely related species is increasing and may provide an avenue for adaptation to a changing environment. While hybrids were rare in the past, they are now thriving in shallow habitats with extreme temperatures and irradiance and are expanding into the parental species habitats. Additional evidence suggests that the hybrid is more disease resistant than at least one of the parental species. Hybridization may therefore have the potential to rescue the threatened parental species from extinction through the transfer of adapted genes via hybrids mating with both parental species, but extensive gene flow may

alter the evolutionary trajectory of the parental species and drive one or both to extinction. This collaborative project is to collect genetic and ecological data in order to understand the mechanisms underlying increasing hybrid abundance. The knowledge gained from this research will help facilitate more strategic management of coral populations under current and emerging threats to their survival. This project includes integrated research and educational opportunities for high school, undergraduate and graduate students, and a postdoctoral researcher. Students in the United States Virgin Islands will take part in coral spawning research and resource managers will receive training on acroporid reproduction to apply to coral restoration techniques.

Current models predict the demise of reefs in the next 200 years due to increasing sea surface temperatures and ocean acidification. It is thus essential to identify habitats, taxa and evolutionary mechanisms that will allow some coral species to maintain their role as foundation fauna. Hybridization can provide an avenue for adaptation to changing conditions. Corals hybridize with some frequency and results may range from the introduction of a few alleles into existing parent species via introgression, to the birth of a new, perhaps better adapted genetic lineage. The only widely accepted coral hybrid system consists of the once dominant but now threatened Caribbean species, *Acropora cervicornis* and *A. palmata*. In the past, hybrid colonies originating from natural crosses between elkhorn and staghorn corals were rare, and evidence of hybrid reproduction was limited to infrequent matings with the staghorn coral. Recent field observations suggest that the hybrid is increasing and its ecological role is changing throughout the Caribbean. These hybrids appear to be less affected by the disease that led to the mass mortality of their parental species in recent decades. Hybrids are also found thriving in shallow habitats with high temperatures and irradiance suggesting they may be less susceptible to future warming scenarios. At the same time, they are expanding into the deeper parental species habitats. Preliminary genetic data indicate that hybrids are now mating with each other, demonstrating the potential for the formation of a new species. Further, hybrids appear to be capable of mating with both staghorn and elkhorn coral, perhaps leading to gene flow between the parent species via the hybrid. Research is proposed to address how the increase in hybridization and perhaps subsequent introgression will affect the current ecological role and the future evolutionary trajectory of Caribbean acroporids. Specifically, this collaborative project aims to answer the following questions: 1) What is the historic rate, direction, and degree of introgression across species ranges and genomes? Linkage block analysis based on genome-wide SNP genotyping across three replicate hybrid zones will answer this question. 2) What is the current extent and future potential of later generation hybrid formation? Morphometric and genetic analyses combined with in vitro fertilization assays will be used. 3) What mechanisms allow hybrids to thrive in hot, shallow waters? A series of manipulative in situ and ex situ experiments will determine whether biotic or abiotic factors favor hybrid survival in shallow waters. 4) Are hybrids more disease resistant than the parentals species? Disease transmission assays in reciprocal transplant experiments and histological analysis to determine the extent of disease will be conducted. A multidisciplinary approach will be taken that combines traditional and cutting edge technology to provide a detailed analysis of the evolutionary ecology of Caribbean corals.

*Note:* PI Nicole Fogarty's original award OCE-1538469 was issued while at Nova Southeastern University. This was replaced by OCE-1929979 upon moving to the University of North Carolina Wilmington.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1537959</a>

[ [table of contents](#) | [back to top](#) ]