

# Time course experiment related data to validate incubation duration for methane oxidation rate measurement of the deep Santa Barbara Basin water column from September 2019 (BASIN project)

**Website:** <https://www.bco-dmo.org/dataset/872687>

**Data Type:** Cruise Results, experimental

**Version:** 1

**Version Date:** 2022-04-05

## Project

» [Collaborative Research: Do benthic feedbacks couple sulfur, nitrogen and carbon biogeochemistry during transient deoxygenation?](#) (BASIN)

Contributors	Affiliation	Role
<a href="#">Valentine, David L.</a>	University of California-Santa Barbara (UCSB)	Principal Investigator
<a href="#">Treude, Tina</a>	University of California-Los Angeles (UCLA)	Co-Principal Investigator
<a href="#">Kinnaman, Franklin S</a>	University of California-Santa Barbara (UCSB)	Scientist
<a href="#">Gosselin, Kelsey M</a>	University of California-Santa Barbara (UCSB)	Student
<a href="#">Liu, Na</a>	University of California-Santa Barbara (UCSB)	Student
<a href="#">Qin, Qianhui</a>	University of California-Santa Barbara (UCSB)	Student, Contact
<a href="#">Newman, Sawyer</a>	Woods Hole Oceanographic Institution (WHOI BCO-DMO)	BCO-DMO Data Manager

## Abstract

This time course experiment was performed to test if the uptake rate of 3H-CH<sub>4</sub> is linear over the chosen incubation time (3 days) for the seawater samples.

---

## Table of Contents

- [Coverage](#)
  - [Dataset Description](#)
    - [Acquisition Description](#)
    - [Processing Description](#)
  - [Related Publications](#)
  - [Related Datasets](#)
  - [Parameters](#)
  - [Instruments](#)
  - [Project Information](#)
  - [Funding](#)
- 

## Coverage

**Spatial Extent:** Lat:34 Lon:-120

**Temporal Extent:** 2019-09-24

## Acquisition Description

## Methodology:

## Sampling and analytical procedures:

Methane concentrations were measured by headspace equilibration method, modified from Kinnaman et al., 2007.

Fractional methane turnover rates were analyzed based on the  $^3\text{H}$ -labeled methane incubation protocol by Bussmann et al., 2015.

Methane oxidation rates were calculated assuming adherence to the first-order rate law, by multiplying fractional methane turnover rate and ambient methane concentration (Valentine et al., 2001).

Samples were collected from R/V Connell.

[ [table of contents](#) | [back to top](#) ]

---

## Related Publications

Bussmann, I., Matousu, A., Osudar, R., & Mau, S. (2015). Assessment of the radio  $^3\text{H}$ -CH<sub>4</sub> tracer technique to measure aerobic methane oxidation in the water column. *Limnology and Oceanography: Methods*, 13(6), 312–327. doi:[10.1002/lom3.10027](https://doi.org/10.1002/lom3.10027)  
*Methods*

Kinnaman, F. S., Valentine, D. L., & Tyler, S. C. (2007). Carbon and hydrogen isotope fractionation associated with the aerobic microbial oxidation of methane, ethane, propane and butane. *Geochimica et Cosmochimica Acta*, 71(2), 271–283. doi:[10.1016/j.gca.2006.09.007](https://doi.org/10.1016/j.gca.2006.09.007)  
*Related Research*

Qin Q., Kinnaman F. S., Gosselin K.M., Liu N., Treude T. and Valentine D.L. (2022) Seasonality of Water Column Methane Oxidation and Deoxygenation in a Dynamic Marine Environment. In review with *Geochimica et Cosmochimica Acta*.  
*Results*

Valentine, D. L., Blanton, D. C., Reeburgh, W. S., & Kastner, M. (2001). Water column methane oxidation adjacent to an area of active hydrate dissociation, Eel river Basin. *Geochimica et Cosmochimica Acta*, 65(16), 2633–2640. doi:[10.1016/S0016-7037\(01\)00625-1](https://doi.org/10.1016/S0016-7037(01)00625-1) [https://doi.org/10.1016/S0016-7037\(01\)00625-1](https://doi.org/10.1016/S0016-7037(01)00625-1)  
*Methods*

[ [table of contents](#) | [back to top](#) ]

---

## Related Datasets

### IsRelatedTo

Qin, Q., Valentine, D. L., Treude, T., Kinnaman, F. S., Gosselin, K. M., Liu, N. (2022) **Initial Methane Concentration Alteration Experiment Data of the Deep Santa Barbara Basin Water Column from October 2019 (BASIN project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-04-05 <http://lod.bco-dmo.org/id/dataset/872652> [[view at BCO-DMO](#)]  
*Relationship Description: Time course experiment, together with initial oxygen and methane alteration experiments, serve as supports of the assumptions of the method.*

Qin, Q., Valentine, D. L., Treude, T., Kinnaman, F. S., Gosselin, K. M., Liu, N. (2022) **Initial Oxygen Concentration Alteration Experiment Data of the Deep Santa Barbara Basin Water Column from October 2019 (BASIN project)**. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 1) Version Date 2022-04-05 <http://lod.bco-dmo.org/id/dataset/872665> [[view at BCO-DMO](#)]  
*Relationship Description: Time course experiment, together with initial oxygen and methane alteration experiments, serve as supports of the assumptions of the method.*

[ [table of contents](#) | [back to top](#) ]

---

## Parameters

Parameter	Description	Units
Incubation_time_day	incubation time of seawater samples; harvest sample days are either 1 (day 1), 2 (day 2), or 2.85 (day 3 at an earlier collection time than days 1 and 2)	unitless
Depth_m	sampling depths of sea water samples	m
Percentage_of_tracer_converted	the percentage of injected tracer 3H-CH4 that is converted to 3H-H2O	unitless

[ [table of contents](#) | [back to top](#) ]

## Instruments

<b>Dataset-specific Instrument Name</b>	6 4-liter Niskin bottles
<b>Generic Instrument Name</b>	Niskin bottle
<b>Dataset-specific Description</b>	Seawater sampling was performed using a rosette equipped with 6 4-liter Niskin bottles, the in-situ temperatures of the water samples were recorded by the conductivity-temperature-depth recorder (CTD), a Seabird SBE 19plusV2 Seacat Profiler system attached to the rosette.
<b>Generic Instrument Description</b>	A Niskin bottle (a next generation water sampler based on the Nansen bottle) is a cylindrical, non-metallic water collection device with stoppers at both ends. The bottles can be attached individually on a hydrowire or deployed in 12, 24, or 36 bottle Rosette systems mounted on a frame and combined with a CTD. Niskin bottles are used to collect discrete water samples for a range of measurements including pigments, nutrients, plankton, etc.

<b>Dataset-specific Instrument Name</b>	Seabird SBE 19plusV2 Seacat Profiler System
<b>Generic Instrument Name</b>	CTD Sea-Bird SBE SEACAT 19plus
<b>Dataset-specific Description</b>	Seawater sampling was performed using a rosette equipped with 6 4-liter Niskin bottles, the in-situ temperatures of the water samples were recorded by the conductivity-temperature-depth recorder (CTD), a Seabird SBE 19plusV2 Seacat Profiler system attached to the rosette.
<b>Generic Instrument Description</b>	Self contained self powered CTD profiler. Measures conductivity, temperature and pressure in both profiling (samples at 4 scans/sec) and moored (sample rates of once every 5 seconds to once every 9 hours) mode. Available in plastic or titanium housing with depth ranges of 600m and 7000m respectively. Miniature submersible pump provides water to conductivity cell.

[ [table of contents](#) | [back to top](#) ]

## Project Information

**Collaborative Research: Do benthic feedbacks couple sulfur, nitrogen and carbon biogeochemistry during transient deoxygenation? (BASIN)**

**Coverage:** Santa Barbara Basin, California, USA

## NSF abstract:

This study focuses on chemical processes that occur in oxygen-limited waters along the world's continental margins. These processes are influenced by the activities of microbes and control the fate of key elements that are deposited to sediments in these areas including carbon, nitrogen and sulfur. As a result, they are key to the health and function of the ocean. The intellectual merit of this research is to study the coupled chemical and microbial processes that occur in these environments by combining robotic technology with experiments that will be conducted at the ocean floor and in the shipboard laboratory. The broader impacts of this project will provide at-sea training and educational opportunities to undergraduate and graduate students and the results will be broadly distributed to stakeholders and interested parties. Results from this research promise to identify and quantify rates for key processes that couple carbon, nitrogen and sulfur in marine environments adjacent to the continents. The project addresses an important aspect of environmental change in the ocean (i.e., decreased oxygen due to warming and nutrient enrichment) and its influence on chemical and biological cycles and ocean ecosystems.

The dynamics of oxygen minimum zones along continental margins, and their potential for future expansion, are important because of their intersection with global biogeochemical cycles and because of their far-reaching impacts on ocean ecosystems. However, the impacts of transient deoxygenation on biogeochemical cycles of carbon, nitrogen and sulfur at the sea floor are not well established and are the focus of this study. This study will test the overarching hypothesis that deoxygenation triggers a positive feedback loop between bacterial mats at the sea floor that consume hydrogen sulfide, a sulfur species that can be toxic to higher organisms, and an underlying community of bacteria that produce hydrogen sulfide. By this hypothesis, the establishment of sea floor mats, which depend on inorganic nitrogen sources to run their sulfur metabolism, accelerates nitrogen cycling in the uppermost sediment horizon following deoxygenation. The accelerated nitrogen cycling allows for upward expansion of the sulfide-producing bacteria, which in-turn provide a shallow source of sulfide as substrate to further support nitrogen cycling in the sea floor mat. The results of this study will enable understanding of the relationship between oxygen dynamics in the water column and the biogeochemical processes at the sea floor that link the transformations of carbon, nitrogen and sulfur. The results of this study promise to define the environmental conditions under which the sulfur and nitrogen cycles are coupled and subject to strong positive feedbacks at the seafloor, as well as the conditions under which they are decoupled. This study provides training in research and innovative analytical and experimental techniques to four graduate students and several undergraduates. Undergraduates will be engaged in research at two institutions, one of which has recently been designated as a Hispanic serving institution. Approximately 10 undergraduate students (20 in total) will participate in each of the two proposed oceanographic expeditions, through an established course entitled: Field Studies in Marine Biogeochemistry. This course provides an opportunity for students to develop an independent research project in advance of the expedition, to participate on the expedition, and to conduct research projects while at sea.

This award reflects NSF's statutory mission and has been deemed worthy of support through evaluation using the Foundation's intellectual merit and broader impacts review criteria.

[ [table of contents](#) | [back to top](#) ]

---

## Funding

Funding Source	Award
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1829981</a>
<a href="#">NSF Division of Ocean Sciences (NSF OCE)</a>	<a href="#">OCE-1830033</a>

[ [table of contents](#) | [back to top](#) ]