

WHOI-97-12

Copy 2

**Woods Hole
Oceanographic
Institution**



**Report on the Office of Naval Research Shallow-Water
Acoustic Workshop, 1-3 October 1996**

by

James F. Lynch

June 1997

Technical Report

Funding was provided by the Office of Naval Research through Contract No. N00014-96-1-1031.
Supported by ONR and DARPA

Approved for public release; distribution unlimited.

WHOI-97-12

**Report on the Office of Naval Research Shallow-Water
Acoustic Workshop, 1-3 October 1996**

by

James F. Lynch

**Woods Hole Oceanographic Institution
Woods Hole, Massachusetts 02543**

June 1997

Technical Report

Funding was provided by the Office of Naval Research through Contract No.
N00014-96-1-1031. Supported by ONR and DARPA

Reproduction in whole or in part is permitted for any purpose of the United States
Government. This report should be cited as Woods Hole Oceanog. Inst. Tech. Rept.,
WHOI-97-12.

Approved for public release; distribution unlimited.

Approved for Distribution:



Timothy K. Stanton, Chair

Department of Applied Ocean Physics and Engineering

ACKNOWLEDGEMENTS

As moderator of this workshop, I would like to thank a number of people for their efforts, cooperation, and goodwill for without them the Shallow-Water Acoustics Workshop would not have been possible. First, to Dr. Ellen Livingston for giving me the opportunity to serve as moderator; and for all her hard work in making it a success. Secondly, appreciation to the four group leaders for their wonderful jobs in leading the panels and ultimately providing the distilled results of their group's deliberations. They are: Drs. Paul Vidmar, Michael Buckingham, and William Siegmann, and Mr. Ed Chaika. I very gratefully acknowledge the co-sponsorship of this workshop by Dr. Bill Carey and DARPA. Bill's enthusiasm for shallow-water acoustics over the years has led to his making numerous contributions to the topic and his sponsorship of and participation in this workshop certainly is another significant contribution. Thanks to Ms. Bev Kuhn and the NRL staff people who took care of the paperwork, logistics, and other details involved in organizing this meeting. The workshop participants also deserve a big thank you -- being moderator of a meeting of an enthusiastic and talented group like the ocean acoustics community is very easy, indeed, as it is their minds and efforts that propelled the meeting. Finally, I'd like to thank my family for tolerating the sometimes extensive amount of traveling that the job of oceanographer entails.

TABLE OF CONTENTS

ACKNOWLEDGEMENTS.....	i
ABSTRACT	iv
INTRODUCTION	v
MODERATOR'S COMMENTARY.....	vii
<i>Introduction</i>	vii
<i>A Brief Revisionist History of Shallow Water Research (1977-1996)</i>	vii
<i>Comments on the Working Group Topics</i>	x
<i>Bottom Acoustics</i>	x
<i>Water Column Acoustics</i>	xi
<i>Modeling</i>	xiv
<i>Postscript</i>	xvi
1. THE BOTTOM ACOUSTICS GROUP REPORT	1
1.1. Introduction.....	1
1.2. Scattering	2
1.3. Geoacoustic Ground Truth	5
1.4. Extrapolation	7
1.5. Sediment Geoacoustics.....	8
1.6. Issues for Experiment Design.....	8
1.7. Recommendations	9
1.8. Navy Relevance.....	10
2. THE WATER COLUMN GROUP REPORT.....	10
2.1. Introduction.....	10
2.2. Acoustic Fluctuations In Shallow Water	11
2.3. Ambient Noise	13
2.4. Concluding Remarks.....	16
3. THE MODELING AND SIGNAL PROCESSING GROUP REPORT.....	17
3.1. Introduction.....	17
3.2. Deterministic Modeling.....	17
3.3. Stochastic Modeling.....	19
3.4. Signal Processing	20
3.5. Last Words	22
APPENDIX A	23
APPENDIX B.....	29

ABSTRACT

The results of an unclassified workshop on Shallow Water Acoustics, jointly sponsored by ONR and DARPA, are presented. The workshop was held on October 1-3, 1996 at the Naval Research Laboratory, Stennis Space Center, and included 83 participants specializing in ocean acoustics, geology and geophysics, physical oceanography, and other disciplines relevant to shallow water research. The goal of the workshop was to help determine the current status of and future directions for shallow water acoustics research. The report summarizes the deliberations and recommendations of the workshop, and includes detailed reports from the three working groups (bottom, water column, and modeling and signal processing) as well as from the workshop moderator (Dr. James Lynch, WHOI).

INTRODUCTION

On October 1-3, 1996 the Office of Naval Research, Code 3210A, and the DARPA ASW Shallow Water Program sponsored an unclassified workshop on Shallow-Water Acoustics at the Naval Research Laboratory at Stennis Space Center, Mississippi. Overall, the intent of this workshop was to have a variety of ocean acoustics investigators interact constructively in order to promote better vertical and horizontal integration among the basic research and the exploratory development research projects. The most significant objective of the workshop was to identify the strengths, needs, and goals of the present ONR Shallow-Water Acoustics program. Finally, the workshop provided an opportunity to discuss the Navy relevance of current research, especially identifying Navy exploratory development projects that are dependent on Shallow-Water Acoustics.

The workshop opened with several programmatic overview talks. These were followed by a full day of short project review and tutorial talks. On the second day, there were four consecutive open panel discussions on topics selected for their importance to current shallow-water acoustic research. These panels were: Bottom Acoustics (Group Leader, Paul Vidmar), Water Column Acoustics (Group Leader, Michael Buckingham), Acoustics Modeling /Signal Processing (Group Leader, William Siegmann), and Vertical and Horizontal Integration Issues (Group Leader, Ed Chaika). In the afternoon, the workshop participants broke into parallel working groups headed by the group leaders on the three technical research topics: Bottom Acoustics, Water Column Acoustics, and Acoustics Modeling/Signal Processing. On the final day of the workshop, the working groups reconvened to generate summary reports of their discussions which were presented at the concluding full session of the workshop. The written report which follows brings together and documents the results and recommendations from this workshop.

MODERATOR'S COMMENTARY ON THE OCT 1-3, 1996 ONR-DARPA SHALLOW WATER WORKSHOP

Jim Lynch, Moderator

Introduction

At my request, Drs. Ellen Livingston and Jeff Simmen accorded me the privilege of writing a small commentary on shallow water acoustics research, and in particular on the Oct. 1-3 Shallow Water Workshop held this autumn at NRL-SSC. Before embarking on this, however, I will make a very strong disclaimer that the statements and opinions contained in *this* section of the report are *strictly my own* and are *not* to be construed as representing either the opinion of the community or of the sponsors of the workshop.

I have written this section *not* because I have any claim to superior expertise in this field (I don't!), but rather because I think these comments may have some small added value in further focusing our results and adding a few more ideas for the future. If even one point I make is a good one that produces a positive result, it will have been worth the effort.

A Brief Revisionist History of Shallow Water Research (1977-1996)

I've now been in the field of underwater acoustic/ acoustical oceanography for 20 years, which gives me just enough of a time series of observations of the field of shallow water acoustics to say that there *is* such a thing as long term progress! When I first looked at shallow water acoustic propagation problems at ARL:UT some 20 years ago, there were decent 2-D ray and mode codes available. (PE was in its infancy and 3-D was a distant dream.) These codes were slow by modern standards, and most were range independent, but they worked. The characteristics of the bottom were known only in a very broad sense, and Hamilton's work was the roadmap for most uses. (His 1980 JASA paper is still highly regarded today, which indicates just how basic his contribution was.) The

water column was commonly regarded as something that could cavalierly be tossed off -- an XBT record in the general vicinity of interest was regarded as perfectly adequate information about the oceanographic field. By and large, "blue water" dominated people's attention anyway, so that the finer points of shallow water oceanography and geology were not first order worries.

In the decade of the 1980's, blue water still dominated the Navy's interest. However, shallow water was not entirely ignored, and research in the area proceeded at a goodly pace. Bottom acoustics was an important topic in the 80's, with much effort being put into measuring and computing what the effects of bottom and subbottom properties were on acoustic propagation. Inverse techniques for bottom properties, mainly linear, began to proliferate. Computer codes for propagation improved dramatically, with the PE and wavenumber integration techniques taking their places along with the standard ray and mode picture codes. Signal processing techniques, notably adaptive techniques and matched field processing (MFP), made substantial strides. Scattering from rough surfaces, particularly the water surface and the water bottom interface, was also explored extensively. The water column, however, remained largely ignored in the context of shallow water/ coastal acoustics.

The first half of the 1990's has also been a fruitful period for shallow water acoustics research. With the dissolution of the Soviet Union around 1990, and the tensions in the Middle East, the Navy's focus turned rather abruptly from "blue water" to "brown water," i.e. shallow water, as a top priority. In line with this, *two* shallow water workshops have been held by ONR this decade (1992 at WHOI and 1996 at NRL-SSC). The nineties have been the "era of the computer." The increase in computing power has made a great impact on modeling and analysis, whereas the decrease in computer size, the low power requirements, and the resistance to high g-forces that have enabled laptop computers to emerge have

made at-sea experimental observations at high bandwidths and over long durations practical. Computer codes have become sophisticated enough to handle much of the important propagation physics in 2-D slices, and 3-D and broadband codes now exist, though not all of them are as “user friendly” and efficient as one would desire them to be eventually. Thanks to these computer advances, bottom inversions using non-linear techniques such as simulated annealing and genetic algorithms have become possible, allowing the inversion of diverse data types for almost any geoacoustic or porous medium variable. Rapid bottom survey techniques are beginning to be developed, with chirp sonar being a prime example, but these also still have a long way to go.

Perhaps the biggest surprise in the 1990's has been the emergence of the water column as an important piece of the shallow water problem in the 25-2500 Hz range. The argument that the water column soundspeed fluctuations were small compared to the bottom property variations, and that in shallow water the downward refracting (or at least isovelocity) water column profile gave the bottom dominance more or less held sway until very recently. The fact that the low mode energy, which is the dominant energy propagated to longer ranges, travels mainly in the water column and thus is more strongly affected by its variability was largely overlooked. However, Zhou's 1991 Yellow Sea experiment paper in JASA started changing opinions about the water column's relative importance, and gave impetus to the current work on soliton scattering. The finding by the Barents Sea Polar Front group that the low modes were coupled far more strongly by a coastal front than by steep bottom bathymetry showed that these features, too, needed to be carefully considered. There have been numerous experiments recently which have focused on water column effects, and which have already shown them to be important determinants of shallow water acoustic propagation and scattering.

At this point, I am up to the present, so let me discontinue this brief “pseudo-historical” aside, and look at the present. I will do this by looking at the topics covered by the working groups.

Comments on the Working Group Topics

Before getting into the individual areas, I would first like to thank the panel and working group leaders for the wonderful jobs they have done. Ellen Livingston and I picked the leaders, Paul Vidmar, Mike Buckingham, Bill Siegmann, and Ed Chaika, because we knew that they all knew the fields they represented quite well and also were extremely dependable. Each group leader produced a fine report, and I thank them very much! The comments I'm adding here are, as I stated, just made in the hope of adding a little more value to their already excellent reports.

Bottom Acoustics

In doing *any* shallow water acoustics experiment, the bottom properties need to be determined to some extent for analysis purposes. However, it can be confidently stated that the bottom properties are the hardest quantities to obtain experimentally. *Rapid* survey techniques for getting at bottom properties are relatively rare (chirp sonar and shots/ airguns with sonobuoy receivers are two common ones I am aware of). More needs to be done, I feel, to make some of the other experimental techniques which have been developed for research into survey techniques, e.g. Hankel transform techniques, impact drop probes, etc., into common survey instruments. (As I type this in Hawaii, I have just learned from Bob Stoll that Sippecan *is* developing a commercial drop probe with him!) Without such rapid, “user friendly” survey techniques, 1) we will never map large areas adequately on a routine basis and 2) experiments that are not specifically “bottom mapping” experiments will be hampered in their analyses due to lack of bottom property information. This might be a 6.1-6.2 transition

issue, but its resolution means a lot to 6.1 research. This also sounds like an area of development where ONR Marine Geology and Geophysics (G&G) participation would be beneficial; G&G contributed heavily to the development of the chirp sonar, and so this would be right in line with their historical interest in developing geological survey tools.

I would also like to see what I will call, for lack of a better phrase, a “better interface” between the poro-elastic (“Biot-Stoll”) descriptions of the bottom and the geoacoustic (complex soundspeed, shearspeed, and density) descriptions. It was clear at the workshop that some people were more comfortable with one picture than another, and that the prescription for going from one picture to another was not clear to many participants. Perhaps a good review article in the Journal of the Acoustical Society by one of the more senior people in bottom acoustics would be helpful to the community as regards that issue. Also, it would perhaps be desirable to have software available for translating from one picture to the other.

Water Column Acoustics

While the water column was ignored for many years (to my mind, anyway), the past few years have seen a number of experiments performed addressing water column issues, which has allowed us to make up significant ground. Since 1992, the following experiments have probed shallow water column issues: 1) the 1992 Barents Sea Polar Front experiment (fronts and internal waves), 2) the 1995 SWARM experiment (nonlinear and linear internal waves), 3) the *three* 1996-97 PRIMER experiments (fronts, eddies, internal waves, and fine structure), 4) the 1996 SESAME experiment (fronts, eddies, internal waves), 5) the Intimate '96 experiment (internal waves), and 6) the 1996 Yellow Sea preliminary experiment (internal waves). I've probably missed a few experiments, as well as slightly misstated the goals of some experiments, for

which I offer my apologies. But that list gives you the basic idea that *something* is definitely happening as far as measuring the water column with good acoustics coupled to good, simultaneous oceanography measurements. I would make two comments on the above group of experiments. First, it might be useful for the PI's of all these experiments to gather for a "mini-conference" to compare analysis results and directions. (Note: there is a special session at the PSU ASA meeting in Spring 97 that will address at least the oceanographic part of many of these experiments.) And second, there are still a lot of measurements and issues that the above experiments did not get to that need to be carefully addressed. On my list of desirable *further* items would be: 1) measurements at a lot more frequencies, going in octaves (at minimum) from 25 Hz to 3200 Hz, 2) longer continuous samples of the acoustic transmissions (10-20 minutes) combined with 2-4 weeks of time series overall, 3) good, 360 degree *azimuthal* measurements of scattering from fronts, internal waves, etc. and 4) horizontal and vertical array deployments for coherence studies.

One interesting point to make concerning the water column experiments and modeling of the water column structure is that shallow water acoustics efforts like the ones above have been taking "ancillary" or "supporting" physical oceanographic (PO) data that have actually pushed the state of the art in PO somewhat. I think some acousticians were surprised at the meeting to find out that the PO community did *not* know about the coastal oceanography on all the time and space scales we needed for our work, and that we were breaking some new ground for PO as well. This came about because, historically, coastal PO did not look so hard at the fine space and time scales we need to know about for acoustics purposes; they generally looked at slower, broader features. Moreover, their measurement techniques were previously somewhat limited compared to modern technology. However, with ADCP's, Sea Soar, and other modern technologies, the PO community *can* help us get the measurements we need,

which are also things they need, as they study finer scales in oceanography. (And conversely, we have helped them!)

The second point to make about the PO is that we should try to link strongly to the PO *modeling* efforts in providing oceanographic input to acoustic propagation and scattering codes. The constraints that ocean dynamical equations can provide give us more sensible inputs to the acoustic models; data assimilating ocean models might produce the best input available. However, these models are not yet as fully developed as would desire for acoustics purposes, and again we are pushing the state of the art in PO. In particular, the PO models do not combine the large scale and fine scale oceanography well yet, i.e. the large scale models will handle “kilometer-ish” resolution and subtidal frequencies, but they won’t handle the $N_{max}(z)$ buoyancy frequencies and 10’s of meters scales needed to handle solitons. Fine scale soliton models (e.g. John Apel’s recent “dnoidal” wave model) on the other hand do not relate to subtidal frequency processes. The link needs to be made, in order to aid both acoustics *and* oceanographic efforts. This may be 6.1, 6.2, or even higher research, but whatever the designation, it will be *useful* work.

One more note on water column phenomena, specifically as regards deterministic versus stochastic descriptions. One of the livelier discussions of the workshop addressed whether one should model propagation through solitons deterministically or stochastically. This is not a trivial question. Given the exact bathymetry, water column stratification, tidal forcing, and wind/ thermohaline forcing of the ocean in the region of generation and propagation of solitons, one could *in theory* predict what we would see for a soliton field. *However*, we will never know all this information, so that in reality we will wind up with an estimation of the field. The field we estimate will have the correct properties for solitons, but its parameters will be different from the real ocean field. We are thus

looking at a deterministic process but with unknown parameters, the typical scenario for internal wave work! Perhaps the biggest difference between solitons and linear waves is that we know the generation sites and mechanisms a lot better for solitons, so that these do have some potential for partial predictability. Given this situation, do we use deterministic models, “conditional probabilities” as McCoy has suggested, or fully random approaches, as Tappert has suggested? This issue will be a lively one to address in the near future, not just for solitons, but for eddies, fronts, and other ocean phenomena which have some dynamical structure and order (though they are also turbulent/ chaotic due to the nonlinearities in the Navier-Stokes equations) but whose exact structure we will never be able to know exactly due to limited measurement resources. I think an indication of how interesting this problem will be is seen in the oceanographic data on solitons taken by the many experiments listed above. To my eye, there are significant *differences* as well as similarities in the soliton/ internal tide fields seen by various investigators, indicating that we will *not* be looking at a trivially simple ocean. Even the linear internal wave field in shallow water is unknown at this time; there is an ONR PO initiative called LIWI investigating this field, but there is much work to do here, and results will probably be few years in coming.

All in all, the coastal water column is providing some extremely good research questions to the science community, and the efforts of that community promise to provide both good science results and vital information to the Navy in the next few years.

Modeling

Modeling of acoustic propagation and scattering is a pretty highly evolved art form at this point in time, and Fred Tappert’s semi-facetious remark that “it has all been done” is not totally amiss. I think modelers *have* explored many of the physics, 3-D, broadband, etc., issues so that they know pretty much what to do

for most cases of interest. However, in practice, the community of model *users* (as opposed to modelers) who employ codes in day-to-day analysis work *do not* have access to all the types of codes that they need. OALLIB (<http://oalib.njit.edu/>) which I regard as one of the best things since sliced bread, provides the basic “flavors” of code, i.e. ray, mode and PE, but these are generally (though not totally) 2-D and narrowband. To do broadband calculations, one generally has to resort to Fourier synthesis, and it would be nice if all the codes available had “hooks” for this in place. 3-D is also a serious issue in shallow water, particularly due to topographic steering, and real 3-D codes (as opposed to Nx2D, which of course is still quite useful) are still relatively rare and a bit hard to use. FOR3D (PE), HARPO (ray) and Ching-Sang Chiu’s 3-D mode code seem to be the front runners in the area. I would personally suggest that ONR, which already supports OALLIB, go even further with its model support and make sure that a very complete suite of models is available to the U.S. underwater acoustics community.

Another issue in modeling is that one uses very accurate models but very inaccurate ocean and ocean bottom input to make predictions of things like propagation loss, travel times, etc. As the models generally take in deterministic data (i.e. either one realization of a random process or the mean of a range of parameter values), we get out a perfect, deterministic prediction *with no error bars*. This is unrealistic, as we all know. Yet there seems to be precious little effort made to quantify the effects of input error in codes. It is tacitly assumed that the user of a code can iterate the model over the full space of possible input parameters to get such answers, but this is generally neither possible nor convenient. David Rubenstein offered a few references to work on getting error/predictability estimates from “smart” algorithms, and I think this is a good topic for the community to pursue.

Postscript

Since I don't have anything much to add to the vertical/ horizontal resolution or signal processing discussions, let me stop here. Again, let me state that the opinions I expressed above are my own. I hope they are not all completely off base, and that some might even be useful.

1. THE BOTTOM ACOUSTICS GROUP REPORT

Paul Vidmar, Group Leader

1.1. Introduction

Scattering from the seabed is one of the most challenging and important problems in understanding acoustic propagation in shallow water. Reverberation, time spreading, and angle spreading of acoustic signals are regularly observed in shallow water experiments and are thought to be caused by scattering from the sea floor and subbottom. Our current inability to make reliable predictions of these phenomena emphasizes our lack of knowledge of the physics of scattering and underscores the importance of the scattering component of bottom acoustics research.

While many individual scientific issues related to scattering were raised during the panel discussion and the meetings of the working group, one recurring topic was the lack of knowledge of the features of the subbottom producing scattering. Geoacoustic “ground truth” is currently lacking and is sorely needed to guide theoretical and computational research, facilitate design of successful field experiments, provide the basis for analysis and interpretation of acoustic data, and for eventual application to Navy problems. The working group also identified some issues concerning the basic geoacoustic properties of sediments. Below we summarize discussions of the working group. Following that, we make some recommendations for future research directions and identify the relevance of this work to the Navy.

1.2. Scattering

Although theoretical and computational work has made progress toward understanding individual scattering mechanisms, it is not clear how this work can be applied to the problem of scattering from the seabed in shallow water. One difficulty is the fact that the sea floor topography and subbottom geoaoustic structure have a complicated mixture of length scales, orientations and physical properties. The scattered field is thus inherently three dimensional, highly frequency dependent, and is produced by many mechanisms acting in concert. Another, possibly more serious, difficulty is that there are very few measurements of subbottom variability on scales important to scattering - and almost none in areas where acoustic data were collected. A knowledge of the subbottom features that cause scattering is crucial for providing direction to theoretical and computational research and for making meaningful interpretation of acoustic data possible. While some progress has been made in developing techniques for measuring subbottom properties at scales of interest, notably applications of borehole tomography (Yamamoto, U. of Miami), chirp sonar (Turgut, NRL), and x-ray tomography (Anderson, Texas A&M, and Turgut, NRL) there is still disagreement in the scientific community concerning the interpretation and accuracy of data collected with these approaches. Below, we have organized and summarized these and other scientific issues raised by the working group.

Statistical Characterization of the Environment

The working group identified three different aspects of the sea floor and subbottom variability that need a statistical description because the scale of variability is so short that a deterministic description is not practical. They are: surface roughness, subbottom fluctuations, and reflecting objects. We distinguish between more continuous variations in properties (fluctuations) and reflecting

objects (such as layers, gas bubbles and shells) because the physics of scattering is different and they may require a different statistical characterization. The key to developing a statistical characterization of these features is availability of measurements of their geoacoustic properties.

Surface roughness: The statistical characterization of surface roughness is a fairly mature subject with many data sets available. The power law spectral characterization appears to work well and has been used by several researchers. The remaining issues involve extrapolation in terms of scale and geographic area (See Extrapolation section below). The application of the same techniques applies to characterizing the small scale roughness of subbottom layers.

Subbottom Inhomogeneities: There are very few measurements of subbottom fluctuations. Yamamoto (U of Miami) uses borehole tomography to measure subbottom fluctuations with resolution of about 1 m or less. He processes the data to obtain a two dimensional power spectral representation and has achieved success in applying this characterization to the prediction of backscattering from the sea floor. On a much smaller scale (1 mm resolution), Anderson (Texas A&M) and researchers at NRL use x-ray tomography to measure the three dimensional structure of very small scale fluctuations. Developing an acceptable three dimensional statistical characterization and will require even more data to be collected at various resolutions.

Reflecting objects: Several classes of reflecting objects are currently being studied. Turgut (NRL) uses chirp sonar data to obtain the three dimensional structure of layers in the subbottom. Anderson (Texas A&M) and NRL researchers use x-ray tomography to obtain the characteristics of gas bubbles, shell fragments, etc. found in the upper part of the sea floor. These data may be sufficient to begin developing statistical characterizations of these objects. The

main issues here involve extrapolation and examining additional scattering objects.

Scattering Mechanisms

Working group discussions identified several scattering mechanisms that need additional theoretical, computational, and experimental work. These mechanisms are: Volume inhomogeneities: Although several researchers have made progress, more work is needed to develop theories that are not empirical but driven by measurable environmental parameters. Work by Yamamoto (U of Miami) is proceeding in this direction using his borehole tomography measurements of velocity and density fluctuations as input to his scattering theory.

Finite size layers: Work needs to be done to describe scattering from a single finite size layer, such as a sand lens in a clay sediment, and from an ensemble of such layers. Edge effects, resonances, and multiple scattering may be important aspects of scattering from finite layers. Work by Turgut (NRL) and Badiey (U of Delaware) is making progress toward providing experimentally derived parameters for such layers in shallow water sediments. Shear coupling: While shear wave effects in marine sediments are negligible under most circumstances, it is possible that they have a role in the scattering process. Two factors make this possible: (a) the gradients in volume inhomogeneities may be two orders of magnitude higher than the average gradients in the sediment, leading to significant gradient driven coupling, and (b) shear waves may travel only a short distance to encounter a scatterer and, hence, may not be totally absorbed. Computational work by Stephen (WHOI) is the only work currently addressing this topic.

Orientation relative to the horizontal: Layers, striations on surfaces (sand waves) or elongated inhomogeneities that are oriented at an angle to the horizontal induce an azimuthal dependence to the scattering process. Borehole tomography carried out by Yamamoto (U of Miami) has measured the non-horizontal orientation of volume inhomogeneities and he has included them in his calculations of backscatter. More work is needed to extend this work and to develop an understanding of scattering from other non-horizontal structures.

Single and multiple scattering: Almost all theoretical work deals with scattering in the single scatter approximation. Work needs to be done to deal with multiple scattering and examine the validity of the single scatter assumption.

Theoretical and Computational Directions

The working group felt that theoretical and computational research should include the following aspects of the scattering problem:

Frequency dependence: Research should describe a broadband process to handle the expected strong frequency dependence of scattering.

Coherence: Research should move beyond predicting mean levels and come to grips with the moments of the acoustic field such as coherence (spatial, temporal, frequency, angular).

Azimuthal dependence: The scattering process is expected to be inherently three dimensional and needs to be treated as such. Out-of-plane scatter due to scatterer geometry or orientation needs to be addressed.

1.3. Geoacoustic Ground Truth

While the working group agreed that we needed to know the structure of the subbottom at scales from some fraction of an acoustic wavelength (maybe as

small as 1/10) to the typical size of an experimental region, no agreement was reached on what to measure or how to obtain that characterization. This state-of-the-art is truly astounding, given the critical need for “ground truth” as a basis for theoretical and computational work, design of experiments, and interpretation of data. Many methods for obtaining data about the subbottom were discussed (see list below), but each had its detractors as well as supporters. Clearly, the problem of characterizing the subbottom must be dealt with before any meaningful progress can be made toward understanding the physics of scattering in shallow water environments. In contrast, measuring and describing bathymetry at the required scales is in fairly good shape. There are recognized methods of collecting data (multibeam bathymetry systems, stereo photography, laser line scan systems) and models for characterizing bathymetry (power spectral models).

Some Methods for Measuring Subbottom Variability

- Borehole tomography for fluctuations and layers
- Chirp sonar for layers
- X-ray tomography of cores for high resolution structure
- Seismic profile for large scale features
- Analysis of cores for vertical structure and fields of cores for horizontal structure
- Extracting P wave velocity from S wave measurements taking advantage of the higher resolution of S wave velocity due to smaller wavelength
 - Penetrometer for shear strength, assuming a relationship between shear strength and shear velocity
 - Interface wave inversion
 - Gravity wave inversion

Electromagnetic measurements, assuming a link between electrical and acoustical properties.

1.4. Extrapolation

This topic deals with the often neglected problem of extending measurements outside their initial domain of validity. There are two aspects of the problem, geographic extrapolation and extrapolation of scale. Geographic extrapolation deals with the problem of extending a few localized measurements of sea floor and subbottom structure to characterize an entire region. Extrapolation of scale deals with the problem of using measurements of structure on one scale to predict structure at larger or smaller scales. The key to making progress in these areas is research to link geological processes and stochastic descriptions relevant to acoustics. Once this link is known, geological data and interpretation can be used to develop sampling criteria, establish limits on extrapolation to higher resolution, and provide guidance for interpolation and smoothing. Thus, extrapolation is a combined geology and acoustics problem that needs coordinated interdisciplinary research for progress to be made. The working group identified research issues that apply to extrapolation of both surface roughness and subbottom variability:

Geographic extrapolation:

- How far from a measurement location does a stochastic description of roughness or subbottom variability apply?
- How far apart do stochastic descriptions need to be measured to characterize an area?

Extrapolation of scale:

- How robust are power law spectral distributions? What are the limits in extending distribution to smaller scale features?
- What procedure should be used to interpolate or smooth data for input to models?

1.5. Sediment Geoacoustics

There are still several outstanding research issues involving the geoacoustic properties of shallow water sediments. Most notable is the problem of attenuation. We are still in the situation where estimates of attenuation based on measured physical properties are not reliable and inversion of acoustic data is needed to obtain attenuation values for an area. We do not know what portion of the measured attenuation is due to scattering and what is due to the absorption of energy. There are also questions concerning the importance of velocity dispersion (which must necessarily accompany absorption), the role of small quantities of gas in sediments, and the mechanism for the penetration of low angle energy into the sea floor. Progress on these issues may well require an interdisciplinary approach involving physicists, chemists, geologists, and acousticians to piece together a porous media theory based on an understanding of the acoustics of the micro-structure of sediments. An additional issue raised by the working group is the possibility of the temporal variability of sediment properties in shallow water due to biological activity, storms, and seasonal or daily heating. We do not know the characteristic spatial or temporal scales for these changes as they affect the geoacoustic properties of the sediment or how they affect the statistical characterization of the sea floor or subbottom.

1.6. Issues for Experiment Design

In addition to emphasizing that experiment design should ensure the collection of all data (including sea floor and subbottom data) needed to test the hypotheses of an experiment, the working group identified several specific issues related to bottom interaction that need attention in designing an experiment.

- The goal of an experiment should be to understand acoustical mechanisms (scattering from roughness, from volume inhomogeneities, from layers, from gas bubbles, etc.) rather than

producing a single number (scattering strength) that combines the effects of all mechanisms. Experiments should be designed to isolate mechanisms.

- Experiments should start with simple stochastic environments and move to more complicated environments as our understanding of scattering physics increases.
- Experiments should be located in existing natural laboratories to take advantage of geological and geoacoustic work that has already been done. Additional characterization of sea floor roughness or subbottom variability needed for an acoustic scattering experiment would extend and complement the existing environmental characterizations of the natural laboratories.
- An area characterization methodology needs to be developed to specify the number and location of detailed sea floor roughness and subbottom variability measurements required in an experiment.
- Calibration of sensors (geophones or hydrophones) on or beneath the sea floor is not well understood.
- Tank and laboratory experiments are valuable and should be continued. They provide very controlled conditions (often impossible to achieve in an at-sea experiment) that can examine the physics of individual scattering mechanisms and convincingly validate theoretical predictions.

1.7. Recommendations

The most important issue for bottom acoustics in shallow water is the problem of characterizing the structure of the subbottom on scales needed to describe scattering mechanisms. The working group could come to no consensus about what constitutes ground truth nor could they identify any measurement techniques that were universally accepted as accurate and reliable. This is clearly a fundamental issue that must be dealt with. Without reliable measurements of subbottom variability, theoretical and computational efforts are hindered and it is impossible to unambiguously interpret acoustic data from at-sea experiments. Other important research issues are those dealing with stochastic characterization of subbottom variability, the physics of individual scattering mechanisms,

extrapolation in scale and location, porous media theories based on the micro-physics of sediments, azimuthal and frequency dependence of scattering, and the coherence of the sound field.

1.8. Navy Relevance

Understanding the scattering process and the ability to predict coherence and environmental spreading of acoustic energy has direct application to improving models used to predict the performance of Navy acoustic systems. Developing a technology for measuring subbottom variability and a methodology for extrapolation would find application in the development and maintenance of geoacoustic databases used to provide input to performance prediction models. It is also possible that new signal processing algorithms and system concepts would flow from an understanding of the effect of the sea floor on active and passive acoustic signals.

2. THE WATER COLUMN GROUP REPORT

Michael J. Buckingham, Group Leader

2.1. Introduction

It is a truism to say that the water column plays the central role in underwater acoustics, for it is here that sounds are created, transmitted and received. The boundaries may exert their influence on the spatial, temporal and even spectral properties of the field, but the water column supports its very existence.

Clearly, the question of acoustic propagation through the ocean medium is an important one. However, ocean-acoustic propagation modeling has been the subject of intensive research over the past decade or more, with the result that a number of highly efficient numerical propagation codes now exist that provide an

adequate prediction capability in many circumstances of interest to the underwater acoustics community. This is not to say that the problem of predicting acoustic propagation in the ocean has been entirely solved, since there are still some areas that need attention, notably in connection with environments that support strong 3-dimensional effects. But it is now felt that existing propagation models are sufficiently well developed for emphasis to be placed elsewhere, on areas of ocean-acoustics research that are important to future navy needs.

Two specific topics, both stochastic in nature, are considered to be of immediate interest: acoustic fluctuations induced by various oceanographic processes; and ambient noise, including air entrainment processes and ambient noise inversions to obtain information about the ocean environment. It is recognized that a research program designed to address these issues must be based on four tightly coupled components: a) experimental planning; b) measurements; c) theory; and d) modeling.

2.2. Acoustic Fluctuations In Shallow Water

Internal waves in deep water are reasonably well represented by the Garrett-Munk spectrum (Flatté, 1979), enabling fluctuations in long-range acoustic transmissions to be characterized accurately (Colosi, 1994) . The same cannot be said of internal waves and their effects on acoustic signals in shallow water. There is a clear need to develop a theoretical model or models of internal waves in shallow water, since this is fundamentally important to interpreting acoustic fluctuations in continental shelf and slope regions. In fact, a number of oceanographic fluctuation mechanisms need to be characterized, with a view to identifying their effects on acoustic fields in shallow water, including the following:

- internal waves
- thermal microstructure
- solitons
- variability in surface bubble plumes
- variability associated with the shelf-break front
- seasonal variability in ocean structure
- tidal variability

Stochastic modeling of acoustic fields in a variable shallow water environment is also required. Some inroads have already been made into this problem in a recently published analysis of scintillation in a shallow-water waveguide by Creamer (Creamer, 1996) . This treatment is based on averaged equations for intensities and fluctuations (second- and fourth-order moments), and provides insights into the fundamental physics underlying the fluctuating field. Further theoretical developments are clearly essential if a more complete understanding of the links between oceanographic and acoustic fluctuations is to be achieved. A parallel experimental effort is also required, aimed at investigating the coupling between the oceanographic driving fluctuations and the resultant acoustic fluctuations.

However well the fluctuating environment is understood, it will still only be possible to predict the properties of a transmitted acoustic signal in an average sense, in terms of its second- and higher-order moments. An alternative approach is found in adaptive techniques, such as phase conjugation, that are currently under investigation. In effect, the variability in the acoustic travel path is sampled and backed-out of the received signals, leaving them unaffected by the oceanographic fluctuations. There are some parallels here to active noise and active vibration control systems that have received such close attention over the

past two decades. An interesting aspect of phase-conjugate systems is that when the receiver array is not focused an inversion can be performed to determine the properties of the fluctuating field. Phase-conjugation, used adaptively for eliminating travel-path fluctuations and in an inverse mode for characterizing the stochastic field, is regarded as a very important technique in connection with future shallow-water propagation studies.

Shallow-water tomography is seen as another area of interest, and one which has received little attention compared to its deep-water counterpart. There seems to be a good case for a tomographic experiment designed to provide data on shallow-water acoustic fluctuations in support of the theoretical studies mentioned above.

In summary, the research topics identified in this section are listed below.

- Development of shallow-water internal-wave spectra (theory & experiment)
- Theoretical development of acoustic fluctuations in shallow water
- Experimental investigations of links between oceanographic and acoustic fluctuations in shallow water
- Development of phase conjugation and other adaptive propagation schemes
- Shallow-water tomography

2.3. *Ambient Noise*

Ambient noise has been of interest to the ocean acoustics community for many years, mainly with a view to reducing its effect on signal-detection capability. Recently, it has come to be recognized that ambient noise itself contains information on the ocean environment, which can be extracted to advantage. It is interesting, however, that the classical Knudsen spectrum of noise due to breaking waves, which has been known for the best part of fifty years, still

has no entirely satisfactory explanation. This is a fundamental problem of ocean physics, which needs to be addressed fairly urgently since it has a bearing on several aspects of ambient noise research, including inversions of the noise to determine ocean processes such as gas fluxes across the air-sea interface.

Wave noise arises from the creation of bubbles just beneath the sea surface. An important feature of the bubble field below a breaking wave is the bubble size distribution, for which there is no satisfactory theoretical model although it is known from measurements that it scales approximately as a^{-n} , where a is bubble radius and $2.5 < n < 6$. Another aspect of the bubble plumes, and one that has received very a little attention, is the bubble creation rate. This is particularly important when passive-acoustic techniques are used for interrogating surface processes, since the signal originates with the newly formed, acoustically active bubbles. That is to say, passive systems respond to the rate of creation, whereas active systems, for instance, upward looking sonars, yield information on the quiescent bubble field. The basic physics of the bubble size distribution and the bubble creation rate are both considered to be very important research issues in support of future ambient noise inversions.

Such inversions include the use of ambient noise to determine gas transfer rates across the air-sea interface, which is relevant to absorption of greenhouse gases by the ocean, which in turn has a bearing on global warming estimates. The noise may also be used to establish the statistical distribution of surface waves; and measurements of noise in the water column can provide information on sea floor parameters and to some extent stratification. All such inversion techniques are relatively new and involve various novel ideas. This raises several questions, including robustness and repeatability of the techniques, and performance limitations. There is a recognized need to examine the new methods of stochastic

signal inversion, with the objective of answering some of these questions and identifying limitations on performance.

A region of accelerating interest is the surf zone. Until very recently, little effort had been directed towards the acoustics of the littoral environment and even now almost everything still needs to be done. Complexity is perhaps the characteristic feature of surf zone processes, in terms of the fluid mechanics, the acoustics, and the coupling between the two. Even the simple question of whether sound propagates out from the breaking region to open water and if so, what is the coupling mechanism, cannot be answered with certainty. The acoustics of the surf zone is identified as a particularly challenging and interesting topic of research, which needs to be addressed systematically, with a view to unraveling the essential physics of the environment. From the experimental point of view, several techniques will be needed to characterize the acoustics and oceanography of the surf zone, including Doppler sonars, upward-looking sonars, and bubble counting and sizing devices (acoustic and optical).

A feature of near-shore ambient noise in temperate and tropical climates is sound from colonies of snapping shrimp. Each pulse is extremely intensive (probably highly non-linear) and very brief, lasting no more than 10^{-5} seconds. The overall noise field from the creatures is impulsive in nature, very non-Gaussian, and originates from a random spatial distribution of sources. The bandwidth of the snapping shrimp sounds is approximately 5 kHz to beyond 100 kHz, spanning the performance range of many active sonars. For such systems, the snapping shrimp may be a cause of serious performance degradation. On the plus side, the shrimp provide a form of acoustic illumination in the water column, which can be used to advantage by the ambient noise imaging systems (Acoustic Daylight) that are currently under development. There is a pressing requirement to

identify the spatial and temporal statistics of snapping shrimp noise fields, since so little is known of their properties.

In summary, the research topics identified in this section are listed below.

- Origin of the breaking-wave noise spectrum
- Develop physics of the bubble size distribution from wave breaking
- Develop physics of the bubble creation rate from wave breaking
- Develop stochastic-field inversion techniques
- Assess performance limitations of stochastic-field inversion techniques
- Establish fundamental coupling between fluid mechanics and acoustics in the surf zone
- Quantify the spatial and temporal distributions of snapping shrimp noise in near-shore locations

2.4. Concluding Remarks

In the above discussions of the current and future status of water column acoustics, the effects of the sea floor have been excluded, although clearly topography and other factors influence shallow-water propagation and ambient noise. But bottom effects are under consideration by another group.

On another issue, the ocean acoustics community has accumulated many high-quality data sets over the years, and it is recommended that, whenever possible, these be used to advantage rather than perform repeat experiments. It is also recommended that existing assets, for instance in the form of a planned research cruise, be exploited to the full by the community to maximize the return on the investment.

3. THE MODELING AND SIGNAL PROCESSING GROUP REPORT

William Siegmann, Group Leader

3.1. Introduction

High-quality modeling and processing tools are essential for fundamental understanding of shallow-water acoustics and for detection and localization problems. The geoacoustic and water-column issues that are relevant for scientific progress and for Navy applications are discussed in previous sections. New capabilities for propagation models and signal processing methods that are required to meet those needs are described here.

3.2. Deterministic Modeling

Much progress has been made in developing and validating deterministic propagation models for shallow water. Minor refinements or small accuracy improvements for these models are not needed. The focus of research should be on incorporating new and relevant physical mechanisms and capabilities. It is not enough to construct and test implementations that require supercomputers to run current problems. The models must be practical and efficient on today's high-end workstations, so that they will have a significant shelf life as computational power continues to increase. The goal is a suite of efficient and benchmarked shallow-water "super-models" which subsume earlier versions and allow easy upgrading.

The primary objective is to incorporate the correct physics, or more precisely the best available understanding of the correct physics for the applications of interest, into the propagation models. Example physical mechanisms in earlier sections range from thermal microstructure to poro-elastic sediments with dispersion and anisotropy. A complete picture of the frequency

dependence of all results is crucial. A related role for the models is to demonstrate the connectivity between micro-mechanical modeling (for example, of highly variable sediments using homogenization techniques) and commonly used effective-media theories.

Another major requirement is fully capable three-dimensional models. The role of 3-D scattering (forward and backward) in shallow water cannot be appraised or treated effectively with existing models. Elastic and poro-elastic sediments need incorporation into 3-D models which are suitable for near-shore and other coastal environments. Broadband 3-D models are considered essential. With advances in physical understanding and computational hardware, the time is ripe for attaining practical 3-D capabilities. No suggestions are made for particular approaches, since each has vigorous advocates; hybrid methods may be the strongest candidates.

Even for the best available propagation models, the sensitivity and predictability pictures are incomplete. For 3-D and other new versions, the levels of field uncertainties need to be specified in terms of environmental uncertainty levels. Another modeling issue which might lead to significantly increased efficiency is the possibility of using impedance conditions at the seabed interface, rather than performing calculations throughout the sediment volume. This entails construction of non-local boundary conditions and useful localized approximations to them, over the range of frequencies and geoacoustic sediment properties.

An unresolved problem is modeling effects of scattering from objects in the ocean volume or the (elastic or poro-elastic) sediment. Previous efforts, emphasizing cases with fluid bottoms, concentrated on acoustic interaction with the object in isolation. The challenge is to treat the fully coupled propagation

problem with objects imbedded in a physically reasonable model of a shallow waveguide.

3.3. *Stochastic Modeling*

Stochastic propagation modeling is critically important for shallow waveguides with substantial and widespread variabilities that are best treated randomly. Some types of stochastic problems for explaining experimental results or testing theoretical predictions are well suited for Monte Carlo treatment. Available well-refined deterministic models can then be used, providing the simulations are feasible. Other types of stochastic problems are better suited for examination by perturbation, moment, or related methods, particularly when behaviors of higher field moments are required. For these it is not enough to develop approximate moment equations; their validity and practicality must be demonstrated.

A principal objective should be determining field coherences (vertical, horizontal, and temporal) in concert with investigating physical mechanisms described previously; a few examples include interface roughness, sediment heterogeneities, gas bubbles, and microstructure. There are corresponding problems for modal and cross-modal coherences. Examining the intermittency of acoustic signals and correlating them with environmental intermittencies is a related issue. The time spreading of broadband signals also should be specified in terms of environmental variabilities.

Scattering and reverberation are persistent modeling problems that have special significance in shallow waveguides. The importance of a practical capability for treating 3-D scattering cannot be overemphasized. This fundamental problem is linked to several issues in previous sections, where relevant physical mechanisms are detailed. Particular attention should be given to using scattering

models to determine scattering cross sections; to discriminate between absorption, attenuation, and scattering losses; to find interface scattering matrix statistics; to specify surface scattering effects (from bubbles, wakes, ice) on signal statistics; and similarly for volumetric (fish schools, microstructure) and bottom (from layering, roughness, inhomogeneities, gas bubbles, and shells) scattering effects. Characteristics of discrete reverberation arising from both isotropic and anisotropic features should be determined, along with 3-D variability from elastic or poro-elastic sediments. Diffuse reverberation, particularly its coherence, from 3-D roughness and inhomogeneities should be treated.

Models of 3-D shallow-water noise statistics, including both natural and man-made sources, are important for fundamental understanding and applications. The evolution of the noise statistics in shallow waveguides needs determined, along with the noise scintillation index. Another issue is developing stochastic propagation tools and procedures for non-stationary fluctuations, one example of which may be the noise field but environmental volume and surface variabilities may be others.

It is important to emphasize that for any of the problem areas, uncertainties in predicted moments are needed in terms of environmental uncertainty estimates. Whether the stochastic propagation is treated by Monte Carlo simulations or averaged moment equations, this requirement holds for signal processing and applications.

3.4. Signal Processing

Results from modeling efforts not only support experimental analyses and enable new theoretical predictions but also impact directly on signal processing and associated inversion techniques. Matched field processing (MFP) is a shining star from the interactions between modeling and signal processing, although

significant issues remain for its optimal and efficient use in shallow-water applications. Such interactions should be strongly encouraged to continue and to extend in other directions as well, particularly for the prediction of field components and statistics that are relevant for signal processing. The following problem areas are focused on processing techniques for source detections and localizations but also concern inversion methods for oceanic and geoacoustic properties. The fundamental ocean science in these two classes of problems are so intertwined that they are not separated here.

Priorities are to specify the quality of matched field processing results and to increase MFP capabilities. The achievable gain using MFP in shallow waveguides has not yet been definitively determined. Statistics should be developed for inversions obtained by MFP. More research is needed for MFP localizations and inversions using horizontal and volumetric arrays. Effects of source, receiver, and target motion on MFP, including reciprocity, integration time, and Doppler, should be determined. A related problem deserving investigation is the influence of non-uniform currents on MFP. Another area suggested for attention is target effects on active MFP applications.

Other research issues are certain extensions for inversion and localization procedures. Among these are capabilities for broadband, coupled modes, and 3-D. Optimum array geometries for inversions should be determined, and emphasis should be placed on specifying parameter resolutions. Target strength inversion, both in the ocean and the (elastic or poro-elastic) sediment, should be investigated using broadband signals.

Efforts should be directed toward promising processing and inversion techniques. One is the collection of phase space methods that include Wigner distributions and wavelets. Another is the set of phase conjugate methods, in

particular their performance in shallow waveguides and with currents. The robustness of mismatch-tolerant processors should be evaluated.

Additional research areas are considered useful for shallow water applications. Noise statistics for signal processing need to be obtained, and opportunistic noise sources should be exploited. The possible advantages of field scintillation characteristics for detection and localization should be explored. The long-term need is to determine optimal array geometries, frequencies, bandwidths, and sensor speeds for shallow waveguide processing.

3.5. *Last Words*

For coping with shallow water challenges, modeling and signal processing tools have perforce become sophisticated physically, mathematically, and computationally. They will surely continue to grow even more so. With the many approaches and implementations, it is essential for users to have confidence in the available codes. Since a variety of benchmark types and problems currently exist, developers are urged to use appropriate ones for testing and validating their implementations. There is a need for new benchmarks, in particular for assessing 3-D, stochastic, broadband, elastic/poro-elastic, and scattering models. Developers should make their codes as widely accessible as possible, taking advantage especially of the web and ftp sites. Thorough documentation of the code, test cases, and operational guidelines earn such great user appreciation as to mitigate the developer's unsung hours in preparing them.

Appendix A

ONR Shallow-Water Acoustics Workshop

Workshop Attendees

Abawi, Ahmad
 NCCOSC RDTE Div D881
 53560 Hull Street
 San Diego, CA 92152-5001
 619-553-3101
 abawi@nosc.mil

Apel, John
 Global Ocean Associates
 PO Box 12131
 Silver Spring, MD 20908
 301-460-7018
 globocen@erols.com

Badiey, Mohsen
 College of Marine Studies
 University of Delaware
 Newark, DE 19716-3500
 302-831-3687
 badiey@ultima.cms.udel.edu

Barbera, Jim
 jbarbera@vrt.com

Berman, David H
 Dept of Physics & Astronomy
 University of Iowa
 Iowa City, IO 52242-1479
 319-335-1223
 berman@iowa.physics.uiowa.edu

Bishop, Judy
 Code 3111
 Naval Undersea Warfare Center
 Newport, RI 02841
 203-432-1207
 lee%veamf1@npt.nuwc.navy.mil

Biondo, Albert
 Johns Hopkins University
 Applied Physics Lab
 Johns Hopkins Road
 Bldg 7 Room 354
 Laurel, MD 20723
 301-953-5000x4286
 albert_biondo@jhuapl.edu

Buckingham, Michael
 Marine Physics Lab
 Scripps Institute of Oceanography
 9500 Gilman Drive
 La Jolla, CA 92093-0213
 619-534-7977
 mjb@mpl.ucsd.edu

Cable, Peter
 BBN Corp
 Union Station
 New London, CT 06320-6147
 203-447-3261
 pcable@bbn.com

Carey, William M
 Massachusetts Institute of Technology
 Ocean Engineering Dept
 Room 5-204
 Cambridge, MA 02139
 617-253-7639
 wcarey@mit.edu

Caruthers, Jerry
 Code 7170
 Ocean Acoustics
 Naval Research Lab/SSC
 Stennis Space Center, MS 39529-5004
 601-688-5438
 jwc@milo.nrlssc.navy.mil

Chaika, Ed
 Naval Meteorology & Oceanography
 Command (CNMOC)
 1020 Balch Blvd
 Stennis Space Ctr, MS 39529-5055
 601-688-4677
 chaikae@cnmoc.navy.mil

Chapman, N. Ross
 School of Earth & Ocean Science
 University of Victoria
 Victoria BC V8W2Y2
 604-472-4340
 chapman@uvic.ca

Chiu, Ching-Sang
 OC/CI
 Naval Post Graduate School
 Monterey, CA 93943
 408-656-3239
 chiu@usw.nps.navy.mil

Collins, Michael D
 Code 7140
 Naval Research Laboratory/DC
 4555 Overlook Avenue, SW
 Washington, DC 20375-5350
 202-404-4823
 collins@abyss.nrl.navy.mil

Colosi, John
 Woods Hole Oceanographic Institute
 Mail Stop 11
 Woods Hole, MA 02543
 508-289-2317
 jcolosi@whoi.edu

Connor, Laurence
 gfrisk@cliff.whoi.edu

Creamer, Dennis B
 Code 5583
 Naval Research Laboratory/DC
 4555 Overlook Avenue, SW
 Washington, DC 20375-5320
 202-767-6949
 creamer@wave23i.nrl.navy.mil

Deane, Grant
 Scripps Institute of Oceanography
 9500 Gilman Drive
 La Jolla, CA 92093-0238
 619-534-0536
 grant@mpl.ucsd.edu

DeFerrari, Prof Harry A
 RSMAS-University of Miami
 4600 Rickenbacker Causeway
 Miami, FL 33149-1098
 305-361-4644/4160
 hdeferrari@rsmas.miami.edu

Denner, Warren
 EOS Research Associates
 200 Camino Aguajito #202
 Monterey, CA 93940
 408-373-1576

Dowling, David R
 Dept Mech Eng and Appl Mech
 University of Michigan
 317 WE Lay Automotive Lab
 Ann Arbor, MI 48109-2121
 313-936-0423
 drd@engin.umich.edu

Duda, Timothy
 Dept Appl Ocean Phys & Eng
 Woods Hole Oceanographic Institute
 Bigelow Lab 2
 Woods Hole, MA 02543
 508-457-2000x2495
 tduda@whoi.edu

Estalote, Eddie
 Code 3210A
 Office of Naval Research
 800 North Quincy Street
 Arlington, VA 22217-5660
 703-696-6940
 estalote@onr.navy.mil

Finette, Steven
 Code 7120
 Naval Research Lab/DC
 Washington, DC 20375
 finette@wave.nrl.navy.mil

Fitch, Robert
 MANDEX
 4001 N 9th Street #106
 Arlington, VA 22203
 703-243-1160
 fitchr@onr.navy.mil

Frazer, L Neil
 School of Ocean & Earth Science
 University of Hawaii at Manoa
 2525 Correa Road
 Honolulu, HI 96822
 808-956-7873
 neil@mano.soest.hawaii.edu

Frisk, George V
 Dept Appl Ocean Phys & Eng
 Woods Hole Oceanographic Institute
 Woods Hole, MA 02543
 508-457-2000x2283
 gfrisk@whoi.edu

Gilbert, Robert
University of Delaware
Newark, DE 19716-3500
302-831-2315
gilbert@math.udel.edu

Gilbert, Kenneth E
Applied Research Laboratory
Pennsylvania State University
PO Box 30
State College, PA 16804
814-863-8291
gilbert@seaair.acs.psu.edu

Glegg, Stewart A L
Dept of Ocean Engineering
Florida Atlantic University
PO Box 3091
Boca Raton, FL 33431-0991
407-367-2633
glegg@oe.fau.edu

Goff, John A
Institute for Geophysics
University of TX at Austin
8701 North MoPac Expresswy
Austin, TX 78759-8397
512-471-0476
goff@utig.ig.utexas.edu

Goodman, Ralph R.
Applied Research Laboratory
Pennsylvania State University
PO Box 30
State College, PA 16804
814-863-8140
gilbert@seaair.acs.psu.edu

Harned, Nancy
Code 321
Office Of Naval Research
Ballston Centre Tower One
800 North Quincy St #407-19
Arlington, VA 22217-5660
703-696-4758
harnedn@onr.navy.mil

Hodgkiss, William S
Marine Physical Lab
Scripps Institute of Oceanography
9500 Gilman Drive
LaJolla, CA 92093-0701
619-534-1798
wsh@mpl.ucsd.edu

Holland, Charles
Planning Systems, Inc
McLean, VA
cholland@plansys.com

Jackson, Pam
Code 7176
Naval Research Lab/SSC
Stennis Space Center, MS 39529-7050
601-688-4782

Koch, Robert

Lee, Ding
Code 3111
Naval Undersea Warfare Center
Newport, RI 02841
203-432-1207
lee%veamf1@npt.nuwc.navy.mil

Lindwall, Dennis
Code 7432
Naval Research Lab/SSC
Stennis Space Center, MS 39529-7050
lindwell@nrlssc.navy.mil

Livingston, Ellen
Code 321OA
Office Of Naval Research
800 North Quincy Street
Arlington, VA 22203-5660
703-696-4203
livinge@onr.navy.mil

Love, Rick
Naval Research Lab/SSC
Stennis Space Center MS 39529-7050

Lu, I-Tai
 Dept of Electrical Engineering
 Polytechnic University
 Route 110
 Farmingdale, NY 11735
 516-755-4226/4215
 itailu@stealth.poly.edu

Lynch, James F
 Dept Appl Ocean Phys and Engr
 Woods Hole Oceanographic Institute
 Woods Hole, MA 02543
 508-548-1400x2230
 jim@vaquero.whoi.edu

McCoy, John J
 School of Engineering
 Catholic University of America
 102 Pangborn Hall
 Washington, DC 20064
 202-319-5160
 mccoy@cua.edu

Mire, Christine
 Naval Research Lab/SSC
 Stennis Space Center, MS 39529-7050

Oba, Roger
 Code 7120
 Naval Research Lab/DC
 4555 Overlook Ave, SW
 Washington, DC 20375-5320

Odom, Robert I
 Applied Physics Lab
 University of Washington
 1013 NE 40th Street
 Seattle, WA 98105
 206-685-3788
 odom@apl.washington.edu

Orr, Marshall H
 Code 7120
 Naval Research Laboratory/DC
 4555 Overlook Avenue, SW
 Washington, DC 20375-5320
 202-767-3359/2192
 orr@wave.nrl.navy.mil

Owsley, Norm
 owsleynl@nl.nuwc.navy.mil

Pasewark, Bruce
 Code 7120
 Naval Research Lab/DC
 4555 Overlook Ave, SW
 Washington, DC 20375-5320
 pasewark@wave.nrl.navy.mil

Porter, Michael B
 Dept of Mathematics
 New Jersey Institute of Technology
 Newark, NJ 07102-1982
 201-596-5782
 miporter@alba.njit.edu

Preston, John
 Applied Research Laboratory
 Pennsylvania State University
 110 Tech Center Bldg
 University Park, PA 16802
 814-863-1310
 preston@ciao.arl.psu.edu

Rajan, Subramaniam D
 Scientific Solutions, Inc
 5907 106th Avenue NE
 Kirkland, WA 98033
 206-828-4866
 rajan@gte.net

Ramsdale, Dan
 Code 7170
 Naval Research Laboratory/SSC
 Stennis Space Center, MS 39529-5004
 601-688-4788
 ramsdale@mistics.nrlssc.navy.mil

Rouseff, Daniel
 Applied Physics Lab, HN-10
 University of Washington
 1013 NE 40th Street
 Seattle, WA 98105-6698
 206-685-3078
 rouseff@apl.washington.edu

Rubenstein, David
 Mail Stop 1-3-2
 Ocean Sciences Operation
 SAIC/Maritime Sciences Group
 1710 Goodridge Dr. POBx 1303
 McLean, VA 22102
 703-827-4748
 davidr@osg.saic.com

Schmidt, Henrik
 Dept of Ocean Engineering
 Massachusetts Institute of Technology
 Cambridge, MA 02139
 617-253-5727
 henrik@keel.mit.edu

Schneider, John
 University of Washington
 1013 NE 40th Street
 Seattle, WA 98105
 509-335-4655
 schneidj@eecs.wsu.edu

Sidorovskaia, Natalia
 Physics Dept
 University of New Orleans
 New Orleans, LA 70148
 nataly@ampsgr.rsmas.miami.edu

Siegmann, William L
 Dept of Mathematical Sciences
 Rensselaer Polytechnic Institute
 Troy, NY 12180-3590
 518-276-6905
 siegmw@euler.math.rpi.edu

Simmen, Jeffrey
 Code 321 OA
 Office of Naval Research
 800 North Quincy St
 Arlington, VA 22217-5660
 703-696-4204
 simmenj@onr.navy.mil

Smith, George
 Naval Research Lab/SSC
 Stennis Space Center, MS 39529-7050

Stanic, Steve
 Code 7174
 Bldg 1005
 Naval Research Laboratory/SSC
 Stennis Space Center MS 39529-5004
 601-688-5235

Stephen, Ralph A
 Dept of Geology & Geophysics
 Woods Hole Oceanographic Institute
 Woods Hole, MA 02543
 508-548-1400x2583
 rstephen@whoi.edu

Stoll, Robert
 Lamont-Doherty Earth Observ
 Columbia University
 PO Box 100
 Palisades, NY 10964
 914-365-8392
 stoll@ldeo.columbia.edu

Tague, John
 Code 321UA
 Office of Naval Research
 800 North Quincy Street 407
 Arlington, VA 22217-5660
 taguej@onr.navy.mil

Tang, Xin
 University of Miami
 4600 Rickenbacker Causeway
 Miami, FL 33149

Tang, D J
 Applied Physics Laboratory
 University of Washington
 1013 NE 40th Street
 Seattle, WA 98105
 206-543-1290
 djtang@apl.washington.edu

Tappert, Frederick
 Div of Applied Mar Phys
 University of Miami, RSMAS
 4600 Rickenbacker Causeway
 Miami, FL 33149
 305-361-4643
 tappert@amp.rsmas.miami.edu

Tolstoy, Alex
 Integrated Perf Decisions, Inc
 2314 Halekoa Drive
 Honolulu, HI 96821
 808-735-8070
 atolstoy@atinc.com

Turgut, Altan
 Code 7120
 Naval Research Lab/DC
 4555 Overlook Ave, SW
 Washington, DC 20375-5320

Vidmar, Paul J
SAIC
920 Andres Avenue
Coral Gables, FL 33134
305-445-4831
pvidmar@shadow.net

Werby, Michael F
Code 7181
Naval Research Laboratory /SSC
Stennis Space Center, MS 39529-5004
601-688-4835
nataly@ampsgi.rsmas.miami.edu

Westwood, Evan
Applied Research Lab
The University of Texas at Austin
P.O. Box 8029
Austin, TX 78713-8029
512-835-3454
westwood@arlut.utexas.edu

Wheatley, Bobby
Stennis Space Center, MS
601-688-4128

Wilkens, Roy H
Institute of Geophysics & Planet.
U of Hawaii
2525 Correa Road
Honolulu, HI 96822
808-956-5228
wilkens@soest.hawaii.edu

Wolf, Stephen N
Code 7120
Naval Research Laboratory/DC
4555 Overlook Avenue, SW
Washington, DC 20375-5320
202-767-3079
swolf@wave.nrl.navy.mil

Wood, Warren
Code 7432
Naval Research Lab/SSC
Stennis Space Center, MS 39529-7050
warren.wood@nrlssc.navy.mil

Worcester, Peter
Institute of Geo & Planetary Phys
Scripps Institute of Oceanography
9500 Gilman Drive
La Jolla, CA 92093-0213
619-534-4688
pworcester@ucsd.edu

Xie, Xiao-B
Institute of Tectonics
Earth Sciences, UCSC
Santa Cruz, CA
408-45-5094
xie@earthsci.ucsc.edu

Yamamoto, Tokuo
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33176
305-361-4637
tok@amp.rsmas.miami.edu

Zhou, Ji-Xun
School of Mech Engineering
Georgia Institute of Technology
Atlanta, GA 30332-0405
404-894-6793
jixun.zhou@me.gatech.edu

Appendix B

ONR Shallow-Water Acoustics Workshop

Workshop Agenda

Moderator: Jim Lynch

----- Agenda and Topics -----

Tuesday, 1 October 1996 8:00 AM - 5:30 PM
--

Introduction	7:55 AM
Programmatic Overview Talks (15 min each)	8:00 AM
ONR Ocean Acoustics Program and Shallow Water Thrust	E. Livingston
ONR 6.3 AEAS Modeling Program	E. Estalote
ARPA Shallow Water Multistatic Active Program	W. Carey
NRL 6.2 Shallow Water Signal Processing	S. Wolf
NRL 6.2 Active Acoustics	R. Love
ONR Active and Passive Signal Processing Programs	N. Harned
Shallow Water Active Detection and Classification	J. Bishop
Tactical Passive-Active Detection and Classification	N. Owsley
ONR Broadband Workshop Report	S. Wolf
General Academic Community Talks (10 min each)	10:30 AM
<i>Bottom Acoustics</i>	
1. Active Bottom Loss Database Development	P. Vidmar
2. Matched Field Tomography for Estimation of Geoacoustic Properties in Shallow Water	R. Chapman
3. Ambient Noise Coherence in Shallow Water	M. Buckingham
4. Modal Mapping in a Complex Shallow Water Environment	G. Frisk
5. Reverberation Derived Low Frequency Shallow Water Bottom Scattering Strength Estimates	P. Cable, M. Steele, J. O'Connor
6. Modeling Elastic Wave Propagation Using a Complex Screen Method	Xiao-Bi Xie, Ru-Shan Wu

- | | |
|--|------------------------------|
| 7. Time Domain Finite Difference Methods in Shallow Water Acoustics | R. Stephen |
| 8. Measurements of the Sound Speed and Attenuation Structure within the Seabed and its Effects on the Propagation and Scattering of Low Frequency Acoustics Waves in Shallow Water | T. Yamamoto |
| 9. In situ Compressional Wave Velocity and Attenuation Measurement Using the Acoustic Lance | R. Wilkins, L. Frazer, S. Fu |

Lunch in Cafeteria Atrium	12:00 Noon
----------------------------------	-------------------

General Academic Community Talks (10 min each)	1:00 PM
---	----------------

Bottom Acoustics - cont'd

- | | |
|---|-------------------|
| 10. 1996 Progress Report for MFP Geoacoustic Tomographic Inversion | A. Tolstoy |
| 11. Computational Shallow Water Acoustics --Shear Waves and Backscattering | D. Lee, J. Bishop |
| 12. Forward and Inverse Problems in Shallow Water | M. Collins |
| 13. Seafloor Characterization within the New Jersey and Northern California STRATAFORM Natural Laboratories: Statistical Analysis from High Resolution Swath Mapping Data | J. Goff |
| 14. Modal Inverses for Bottom Properties | S. Rajan |
| 15. Frequency Dependent Sound Transmission in Shallow Water Regions | M. Badiay |
| 16. Direct and Inverse Acoustic Problems in Shallow Oceans | R. Gilbert |
| 17. Inversion Techniques for Characterizing a Coupled Mode Acoustic Waveguide | D. Rouseff |

General Academic Community Talks (10 min each)	2:30 PM
---	----------------

Water Column Acoustics

- | | |
|--|--|
| 18. Preliminary Acoustic Results from the Intimate 96 Shallow Water Tomography Experiment | E. Coelho, S. Jesus, M. Porter, Y. Stephan |
| 19. The New Jersey Shallow Water Acoustic Random Media Propagation Experiment – SWARM (<i>SWARM Group: J. Apel, M. Badiay, J. Berkson, K.P. Bongiovanni, J. Bouthillette, E. Carey, C.S. Chiu, T. Duda, C. Eck, S. Finette, R. Headrick, J. Irish, J. Kemp, J. Lynch, A. Newhall, M. Orr, B. Pasewark, J. Preisig, B. Racine, S. Rosenblad, A. Shaw, D. Taube, D. Tielbuerger, A. Turgut, K. von der Heydt, W. Witzell, S. Wolf</i>) | SWARM Group |

- | | |
|---|---|
| 20. The SESAME Experiments-the Effects of Internal Solitons on Acoustics Propagation at the Malin Shelf | R. Field, C. Mire,
H. Chandler,
M. Broadhead |
| 21. Shelfbreak PRIMER-an Integrated Acoustic and Oceanographic Field Study in the Middle Atlantic Bight (<i>PRIMER GROUP: R.C. Beardsley, K.H. Brink, M.J. Caruso, C.S. Chiu, G.G. Gawarkiewicz, J. F. Lynch, J.H. Miller, R. Pickart, A.R. Robinson, K.B. Smith</i>) | PRIMER Group |
| 22. Time-Reversing and Phase-Conjugate Arrays | D. Dowling,
S. Khosla |
| 23. Variability of Acoustic Transmissions in the Strait of Gibraltar | P. Worcester,
B. Cornuelle,
C. Tiemann,
U. Send |
| 24. Internal Tides and Solitons on the Continental Shelf: the "cnoidal" Wave Solution to the KDV Equation | J. Apel, M. Orr,
S. Finette, J. Lynch |
| 25. Coupled Acoustic Mode Propagation through Continental Shelf Internal Solitary Waves | T. Duda, J. Preisig |
| 26. Observations of cnoidal Internal Waves and their Effect on Acoustic Propagation in Shallow Water | D. Rubenstein |
| 27. Fluctuations, Coherence, and Predictability of Long Range Acoustic Propagation in Shallow Water | H. DeFerrari,
C. Monjo |
| 28. Effects of Internal Waves on Sound Pulse Propagation in the Straits of Florida | X. Tang, F. Tappert |
| 29. Large Acoustic Scintillations in the Straits of Florida | F. Tappert, X. Tang,
D. Creamer |
| 30. Environmental Adaptive Acoustics in Shallow Water: An Acoustic Demonstration of an Acoustic Time Reversal Mirror | W. Kuperman,
W. Hodgkiss, H.
Song, T. Akal,
C. Ferla, D. Jackson |
| 31. Robustness of a Ray Travel Time Inversion Approach | I-Tai Lu |
| 32. A Model to Understand the Biological Sonars of Dolphins | P. Moore |
| 33. Reverberation and Internal Wave Studies | J.X. Zhou |

Wednesday, 2 October 1996 8:00 AM - 5:30 PM
--

- | | |
|--|---|
| General Academic Community Talks (10 min each) | 8:00 AM |
| <i>Modeling and Signal Processing</i> | |
| 34. Moderately Broadband Shallow Water Acoustics | J. McCoy |
| 35. Coupled Mode Representations for Propagation in Deterministic and Stochastic Range Dependent Shallow Water | R. Odom, M. Park,
V. Peyton, J. Mercer |
| 36. Broadband Model Predictions for Shallow Water Pulse Spreading | J. Preston |
| 37. Shallow Water Propagation Effects and Predictability | W. Siegmann |
| 38. The Effects of Rescattering by Rough Interfaces in Waveguides | D. Berman |
| 39. Three Dimensional Sound Propagation in Shallow Water | S. Glegg, J. Riley |
| 40. Pulse Propagation With And Without Range Dependence | M. Werby,
N. Sidorovskaia |
| 41. Seismo-Acoustic Field Statistics in Shallow Water | H. Schmidt |

Open Panel Discussions (2.5 hours total)	9:30 AM
--	----------------

Bottom Acoustics (Group Leader: Paul Vidmar)
Water Column Acoustics (Group Leader: Michael Buckingham)

WORKING LUNCH BREAK	12:00 Noon
----------------------------	-------------------

Open Panel Discussions (2.5 hours total)	12:30 PM
--	-----------------

Acoustics Modeling and Signal Processing (Group Leader: William Siegmann)
Vertical and Horizontal Integration Issues (Group Leader: Ed Chaika)

Working Groups Meet in Parallel on Major Topics (2.5 hours)	3:00 PM
---	----------------

Bottom Acoustics (Group Leader: Paul Vidmar)
Water Column Acoustics (Group Leader: Michael Buckingham)
Acoustics Modeling and Signal Processing (Group Leader: William Siegmann)

Thursday, 3 October 1996 8:00 AM - 1200 Noon

- | | |
|---|-------------------|
| Working Groups Meet on Major Topics - cont'd (3 hours) | 8:00 AM |
| Reports of Working Groups | 11:00 AM |
| Meeting Adjourned EXCEPT FOR GROUP LEADERS | 12:00 Noon |
| Group Leaders Combine to Organize Final Report (2 hours) | 12:00 Noon |

Resource person: Bev Kuhn (703) 696-6998 kuhnb@onrhq.onr.navy.mil

DOCUMENT LIBRARY

Distribution List for Technical Report Exchange – February 1996

University of California, San Diego
SIO Library 0175C
9500 Gilman Drive
La Jolla, CA 92093-0175

Hancock Library of Biology & Oceanography
Alan Hancock Laboratory
University of Southern California
University Park
Los Angeles, CA 90089-0371

Gifts & Exchanges
Library
Bedford Institute of Oceanography
P.O. Box 1006
Dartmouth, NS, B2Y 4A2, CANADA

Commander
International Ice Patrol
1082 Shennecossett Road
Groton, CT 06340-6095

NOAA/EDIS Miami Library Center
4301 Rickenbacker Causeway
Miami, FL 33149

Research Library
U.S. Army Corps of Engineers
Waterways Experiment Station
3909 Halls Ferry Road
Vicksburg, MS 39180-6199

Institute of Geophysics
University of Hawaii
Library Room 252
2525 Correa Road
Honolulu, HI 96822

Marine Resources Information Center
Building E38-320
MIT
Cambridge, MA 02139

Library
Lamont-Doherty Geological Observatory
Columbia University
Palisades, NY 10964

Library
Serials Department
Oregon State University
Corvallis, OR 97331

Pell Marine Science Library
University of Rhode Island
Narragansett Bay Campus
Narragansett, RI 02882

Working Collection
Texas A&M University
Dept. of Oceanography
College Station, TX 77843

Fisheries-Oceanography Library
151 Oceanography Teaching Bldg.
University of Washington
Seattle, WA 98195

Library
R.S.M.A.S.
University of Miami
4600 Rickenbacker Causeway
Miami, FL 33149

Maury Oceanographic Library
Naval Oceanographic Office
Building 1003 South
1002 Balch Blvd.
Stennis Space Center, MS, 39522-5001

Library
Institute of Ocean Sciences
P.O. Box 6000
Sidney, B.C. V8L 4B2
CANADA

National Oceanographic Library
Southampton Oceanography Centre
European Way
Southampton SO14 3ZH
UK

The Librarian
CSIRO Marine Laboratories
G.P.O. Box 1538
Hobart, Tasmania
AUSTRALIA 7001

Library
Proudman Oceanographic Laboratory
Bidston Observatory
Birkenhead
Merseyside L43 7 RA
UNITED KINGDOM

IFREMER
Centre de Brest
Service Documentation - Publications
BP 70 29280 PLOUZANE
FRANCE

REPORT DOCUMENTATION PAGE	1. REPORT NO. WHOI-97-12	2.	3. Recipient's Accession No.
4. Title and Subtitle Report on the Office of Naval Research Shallow-Water Acoustic Workshop, 1-3 October 1996			5. Report Date June 1997
7. Author(s) James F. Lynch			6.
9. Performing Organization Name and Address Woods Hole Oceanographic Institution Woods Hole, Massachusetts 02543			8. Performing Organization Rept. No. WHOI-97-12
12. Sponsoring Organization Name and Address Office of Naval Research and DARPA			10. Project/Task/Work Unit No.
			11. Contract(C) or Grant(G) No. (C) N00014-96-1-1031 (G)
15. Supplementary Notes This report should be cited as: Woods Hole Oceanog. Inst. Tech. Rept., WHOI-97-12.			13. Type of Report & Period Covered Technical Report
16. Abstract (Limit: 200 words)			14.
<p>The results of an unclassified workshop on Shallow Water Acoustics, jointly sponsored by ONR and DARPA, are presented. The workshop was held on October 1-3, 1996 at the Naval Research Laboratory, Stennis Space Center, and included 83 participants specializing in ocean acoustics, geology and geophysics, physical oceanography, and other disciplines relevant to shallow water research.</p> <p>The goal of the workshop was to help determine the current status of and future directions for shallow water acoustics research. The report summarizes the deliberations and recommendations of the workshop, and includes detailed reports from the three working groups (bottom, water column, and modeling and signal processing) as well as from the workshop moderator (Dr. James Lynch, WHOI).</p>			
17. Document Analysis a. Descriptors shallow water acoustics underwater acoustics basic research recommendations b. Identifiers/Open-Ended Terms c. COSATI Field/Group			
18. Availability Statement Approved for public release; distribution unlimited.		19. Security Class (This Report) UNCLASSIFIED	21. No. of Pages 55
		20. Security Class (This Page)	22. Price