

Soil warming alters nitrogen cycling in a New England forest: implications for ecosystem function and structure

S.M. Butler¹, J.M. Melillo¹, J.E. Johnson^{1,2}, J. Mohan^{1,3}, P.A. Steudler¹, H. Lux^{1,4}, E. Burrows^{1,5}, R.M. Smith^{1,6}, C.L. Vario^{1,7}, L. Scott¹, T.D. Hill^{1,8}, N. Aponte^{1,9} and F. Bowles¹⁰

¹The Ecosystem Center, Marine Biological Laboratory, Woods Hole, MA, ²Biology Department, Stanford University, Palo Alto, CA, ³School of Ecology, University of Georgia, Athens, GA, ⁴Harvard Forest, Petersham, MA, Department of Chemistry and Chemical Biology, ⁵Rutgers University, Piscataway, NJ, ⁶Geology Department, University of Maryland, College Park, MD, ⁷Department of Biological Sciences, Dartmouth College, Hanover, NH, ⁸Yale School of Forestry and Environmental Studies, New Haven, CT, ⁹University of Puerto Rico at Mayagüez, ¹⁰Research Designs, Lyme, NH

Electronic Supplementary Material 1




Figure S1. Concentration of (a) ammonium and (b) nitrate in lysimeter samples collected at 50cm depth, in response to soil warming. Each circle represents the mean of n=7 lysimeter samples \pm 1 SE. Sample months varied by year.




Figure S2. Concentration of nitrous oxide measured in response to soil warming. Each bar represents the mean of n=6 gas samples \pm 1 SE.